

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS	PA Required	Criteria
----------	--------------------	------------	-----------------	--------------------	------------	---------------------	-------------	----------

PDL Effective February 2, 2024

***PLEASE NOTE: For a search box hit Ctrl F**

*** PLEASE NOTE: All cost effective generics applicable to DEL are considered PREFERRED Drugs. "BASIC" Covered Drugs are bolded with the Coverage Indicator of "MC / DEL".**

General Criteria for all PDL categories- For more information or help using the PDL, providers may call 1-888-445-0497; members should call 1-866-796-2463. To access PDL and PA materials via the internet: www.mainearepdl.org

A: Preferred Drugs- Unless otherwise specified, preferred drugs are available without prior authorization. Step order may apply for preferred drugs in some drug categories as indicated on the PDL. (See item "D" below for explanation of step order.)

C: Adequate Drug Trials- 1. The minimum trial period for each preferred and step order drug is two weeks, unless otherwise stated within specific PDL drug categories; trials with less than a two week duration will be reviewed on a case-by-case basis; 2. A trial will not be considered valid if preferred or non-preferred products were readily available (by override, individual purchase, samples, etc.); 3. Certain drug trials, such as with controlled substances, may require evidence that the preferred drugs were actually tried (example: with random pill counts and with random urine drug tests, using the methods of GC/MS with no lower threshold); 4. Adequate trials require documentation of attempts to titrate dose of preferred agents toward desired clinical response. 5. Adequate trials include prevention/treatment of common adverse effects associated with preferred agents (example: antinausea, antipruritics, etc.)

D: Step Order- When numbers appear in the "step order" column, it means drugs in this category must be used in the order specified, with the lower numbers having preference over the higher numbers. Chart notes should be provided to confirm drug trials that do not appear in the member's MaineCare drug profile.

E. The Department will institute strategies to ensure cost effectiveness through the use of an enhanced Drug Benefit Preferred brand drugs will no longer be preferred in any PDL drug category where preferred generic drugs are also available. It is expected that preferred generics will be used prior to any preferred brands. This will be operated as a form of step care. Preferred brands in these categories will require prior authorization for these high utilization / high cost members.

F: Brand Name Medication Requests- (Must be submitted on the Brand Name PA request form)- According to MaineCare Benefits Manual Chapter II (80.07-5), when medically necessary covered brand-name drugs have an A-rated generic equivalent available, the most cost effective medically necessary version will be approved and reimbursed, since the brand-name and A-rated generic drugs have been determined by the FDA to be chemically and therapeutically equivalent. The Bureau does not make determinations as to whether or not a generic drug is clinically inferior or inequivalent to its brand version. This is the proper role of the FDA. Physicians should submit their reports of generic inequivalence directly to the FDA via the MEDWATCH.

G: PA requests for non- FDA Approved Indications- Decisions will be made on a case-by-case basis until the DUR committee is able to review the evidence and make a recommendation. Interim approvals and DUR recommendations for approval of a drug for a non- FDA approved indication will require a minimum of two published, peer reviewed, non contradicted, double- blind, placebo-controlled randomized clinical studies establishing both safety and efficacy.

H: Dose Consolidation Requirements- Some drugs may also be affected by dose consolidation requirements. Please see Dose Consolidation List and/or Splitting Tables provided in the PDL.

I. Trials from Multiple Drug Classes - Trial/failure/intolerance to preferred agents from multiple classes within the same category or other categories of drugs may be required prior to the approval of non-preferred agents (e.g., Cymbalta, Zofran, Elidel and others).

J. Drug-specific PA Forms- Drug-specific PA forms contain medical necessity documentation requirements and/or criteria that may not be repeated in the PDL. Drug-specific PA forms may be obtained on the web at www.mainearepdl.org.

K. PA Exemptions for Prescribers- According to MaineCare Benefits Manual Chapter II (80.07-4), providers may receive a three (3) month exemption from prior authorization requirement for certain categories of drugs when they demonstrate high compliance with the Department's PDL. The Department will notify providers in writing which drug categories are included and what dates apply to the exemption. If a provider loses his/ her exemption, members who previously were not required to obtain a PA while the prescriber was exempt will be required to do so, and criteria for approval of that medication will need to be met.

L: Drug-Drug Interactions (DDI)- The DUR Committee has implemented new drug-drug interaction edits requiring prior authorization. Several drug-drug combinations and PDL drug categories are affected by new PA requirements. These will be indicated in the PDL with DDI notation. Please see the DDI document provided in the PDL.

ASSORTED ANTIBIOTICS

BETA-LACTAMS / CLAVULANATE COMBO'S	MC/DEL		AMOXICILLIN	MC/DEL		AUGMENTIN ³	3. Chewable 125mg & 250mg and Solution 125mg/5ml and 250mg/5ml available without PA. 4. Use preferred generic amoxicillin/clavulanate potassium alternatives. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Ampicillin will now be non-preferred and require prior authorization if it is currently being used in combination with either Prevacid, pantoprazole, Prilosec, or any currently non preferred PPI.
	MC/DEL		AMOXICILLIN/POTASSIUM CLA CHEW	MC/DEL		AUGMENTIN XR TB12 ⁴		
	MC/DEL		AMOXICILLIN/POTASSIUM CLA SUSR					
	MC/DEL		AMOXICILLIN/POTASSIUM CLA TABS					
	MC/DEL		AMPICILLIN					
	MC		BICILLIN L-A SUSP					
	MC/DEL		DICLOXACILLIN SODIUM CAPS					
	MC		OXACILLIN SODIUM SOLR					
	MC/DEL		PENICILLIN V POTASSIUM					
	MC		TIMENTIN SOLR					
MC		UNASYN SOLR						
MC/DEL		ZOSYN						
CEPHALOSPORINS	MC/DEL		CEFADROXIL HEMIHYDRATE	MC		CEDAX	1. Both brand and generic are clinically non-preferred. 2. Dosing limits apply, please see Dosage Consolidation List. 3. Approvals will only be considered for patients 18 years of age or older who have limited or no alternative	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Vantin will now be non-preferred and require prior authorization if it is currently being used in combination with either Prevacid, pantoprazole, Prilosec, or any currently non
	MC/DEL		CEFAZOLIN SODIUM SOLR	MC/DEL		CEFACTOR ¹		
	MC/DEL		CEFDINIR	MC/DEL		CEFADROXIL MONOHYDRATE TABS		
	MC/DEL		CEFEPIME	MC/DEL		CEFIXIME SUS		
	MC/DEL		CEFPODOXIME	MC/DEL		CEPHALEXIN TABS		
	MC/DEL		CEFPODOXIME PROXETIL SUS	MC		CEPHALEXIN 750MG CAPS		
	MC/DEL		CEFPODOXIME PROXETIL TAB	MC/DEL		CEFTIN		
	MC/DEL		CEFIXIME 400MG ² CAP	MC		DAXBIA		
	MC/DEL		CEFPROZIL	MC		FETROJA ³		

	<p>MC/DEL</p> <p>MC</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC</p> <p>MC/DEL</p> <p>MC</p>	<p>CEPHALEXIN 250MG & 500MG CAPS</p> <p>CEFTAZIDIME 6MG</p> <p>CEFTIN SUSP</p> <p>CEFTRIAZONE</p> <p>CEFUROXIME AXETIL TABS</p> <p>CEPHALEXIN MONOHYDRATE</p> <p>FORTAZ SOLR</p> <p>SUPRAX CHEWABLE</p> <p>TAZICEF 6GM</p>	<p>MC/DEL</p> <p>MC/DEL</p> <p>MC</p> <p>MC</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC</p>	<p>FORTAZ</p> <p>FORTAZ SOLN</p> <p>KEFLEX CAPS</p> <p>OMNICEF</p> <p>ROCEPHIN</p> <p>SUPRAX²</p> <p>TAZICEF SOLR</p> <p>TEFLARO</p>	<p>treatment options for the treatment of complicated urinary tract infections (cUTIs)</p> <p>Use PA Form# 20420</p>	<p>preferred PPI.</p> <p>As outlined in the US CDC Guidance on the Use of Expedited Partner Therapy (EPT) in the Treatment of Gonorrhea, MaineCare will cover a single 800 mg dose of cefixime for the treatment of gonorrhea as part of EPT.</p>
MACROLIDES / ERYTHROMYCIN'S	<p>MC/DEL</p> <p>MC/DEL</p> <p>MC</p> <p>MC</p> <p>MC</p> <p>MC</p> <p>MC</p> <p>MC/DEL</p>	<p>AZITHROMYCIN TABS</p> <p>AZITHROMYCIN SUSP</p> <p>E.E.S.</p> <p>ERYPED 200 SUSR</p> <p>ERYPED 400 SUSR</p> <p>ERY-TAB TBEC</p> <p>ERYTHROCIN STEARATE TABS</p> <p>ERYTHROMYCIN</p>	<p>MC/DEL</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC</p> <p>MC</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC/DEL</p>	<p>AZITHROMYCIN POW</p> <p>CLARITHROMYCIN SUSP</p> <p>CLARITHROMYCIN TABS</p> <p>DIFICID</p> <p>PCE TBEC</p> <p>ZITHROMAX TABS</p> <p>ZITHROMAX 1GM PAK</p> <p>ZITHROMAX TRI-PAK</p> <p>ZITHROMAX SUSP</p> <p>ZMAX</p> <p>ZINPLAVA</p>	<p>1. 7- Day supply per month without PA.</p> <p>Use PA Form# 20420</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>DDI: Preferred erythromycin will now be non-preferred and require prior authorization if it is currently being used in combination with either Carbamazepine, Enblex 15mg or Vesicare 10mg. Any non preferred formulation of erythromycin will require prior authorization and the member's drug profile will also be monitored for concurrent use with either Carbamazepine, Enblex 15mg or Vesicare 10mg.</p> <p>DDI: Preferred clarithromycin formulations (clarithromycin tablets) will now be non-preferred and require prior authorization if they are currently being used in combination with either Carbamazepine, Onglyza 5mg, Enblex 15mg or Vesicare 10mg. Any non preferred formulation of clarithromycin will require prior authorization and the member's drug profile will also be monitored for concurrent use with either Carbamazepine, Onglyza 5mg, Enblex 15mg or Vesicare 10mg.</p> <p>Zinplava® will be non-preferred and require clinical prior authorization to verify it is prescribed or consulted by GI or ID specialist, diagnosis, and concurrent use of an antibacterial agent as well as limiting its use to those who have recurrent C. diff disease that has recurred despite use of guideline recommended vancomycin taper or for whom this would be contraindicated.</p>
TETRACYCLINES	<p>MC/DEL</p> <p>MC/DEL</p> <p>MC/DEL</p>	<p>DOXYCYCLINE MONOHYDRATE 100mg & 50mg CAPS</p> <p>MINOCYCLINE HCL CAPS</p> <p>TETRACYCLINE HCL CAPS</p>	<p>MC</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC</p> <p>MC/DEL</p> <p>MC</p> <p>MC/DEL</p> <p>MC</p>	<p>DECLOMYCIN TABS</p> <p>DORYX CPEP</p> <p>DOXYCYCLINE HYCLATE</p> <p>DOXYCYCLINE MONOHYDRATE 150mg & 75mg CAPS</p> <p>DYNACIN CAPS</p> <p>MINOLIRA ER</p> <p>NUZYRA¹</p> <p>ORACEA</p> <p>PERIOSTAT</p> <p>SEYSARA²</p> <p>SOLODYN ER</p> <p>XIMINO</p>	<p>Use PA Form# 20420</p> <p>1. For the treatment of patients ≥ 8 years of age.</p> <p>2. For the treatment of patients ≥ 9 years of age.</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>
FLUOROQUINOLONES	<p>MC/DEL</p> <p>MC/DEL</p> <p>MC/DEL</p>	<p>CIPROFLOXACIN</p> <p>LEVOFLOXACIN</p> <p>OFLOXACIN</p>	<p>MC</p> <p>MC</p> <p>MC</p> <p>MC</p> <p>MC</p> <p>MC</p> <p>MC</p> <p>MC</p> <p>MC</p>	<p>AVELOX SOLN</p> <p>AVELOX ABC PACK TABS</p> <p>BAXDELA</p> <p>CIPRO</p> <p>FACTIVE</p> <p>LEVAQUIN TABS SOLN/INJ</p> <p>LEVAQUIN TABS¹</p> <p>NOROXIN TABS</p> <p>PROQUIN XR</p>	<p>Use PA Form# 20420</p> <p>1. Dosing limits apply, see Dosage Consolidation List.</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>DDI: Preferred ofloxacin will now be non-preferred and require prior authorization if they are currently being used in combination with amiodarone.</p> <p>DDI: Preferred levofloxacin will now be non-preferred and require prior authorization if they are currently being used in combination with amiodarone.</p> <p>DDI: Preferred Avelox will now be non-preferred and require prior authorization if they are currently being used in combination with amiodarone.</p> <p>DDI: All preferred fluoroquinolones will require clinical PA for patients over 60 that are currently on immunosuppressants or steroid therapy.</p> <p>DDI: Factive is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with amiodarone.</p>
AMINO GLYCOSIDES	<p>MC</p> <p>MC</p> <p>MC/DEL</p> <p>MC/DEL</p>	<p>GENTAMICIN</p> <p>KITABIS PAK</p> <p>NEOMYCIN SULFATE TABS</p> <p>TOBRAMYCIN AMPUL-NEB</p>	<p>MC/DEL</p> <p>MC</p> <p>MC/DEL</p> <p>MC</p> <p>MC/DEL</p> <p>MC/DEL</p>	<p>ARIKAYCE^{1,2}</p> <p>BETHKIS¹</p> <p>TOBI PODHALER¹</p> <p>TOBI NEBU²</p> <p>TOBRAMYCIN SULFATE SOLN²</p> <p>ZEMDRI²</p>	<p>Use PA Form# 20420</p> <p>1. Clinical PA to verify appropriate diag</p> <p>2. See criteria section</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>TOBI Podhaler is limited to patients with significant impairment from using nebulized version of medication</p> <p>Current users of Tobi Nebu and Tobramycin Soln will be allowed a grace period until 10/1/15 to transition to preferred Kitabis.</p>

							Arikayce will require clinical PA to confirm MAC lung disease and for use in adults who have limited or no alternative treatment options. Zemdri will be reserved for patients with limited or no alternative treatment of care.
ANTI-MYCOBACTERIALS / ANTI-TUBERCULOSIS	MC/DEL MC/DEL MC/DEL MC/DEL		ETHAMBUTOL HCL TABS MYAMBUTOL TABS RIFABUTIN CAPS RIFAMPIN	MC/DEL MC/DEL MC		MYCOBUTIN CAPS PRETOMANID RIFADIN CAPS	Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Pretomanid is indicated as part of a combination regimen with bedaquiline and linezolid for the treatment of adults with pulmonary extensively drug resistant (XDR) or treatment-intolerant or non-responsive multidrug-resistant (MDR) tuberculosis (TB). Approval of this indication is based in limited clinical safety and efficacy data. This drug is indicated for use in a limited and specific population of patients. DDI: Preferred rifampin will be non-preferred and require prior authorization if it is currently being used in combination with either Pradaxa or Latuda.
ANTIMALARIAL AGENTS	MC/DEL MC MC/DEL MC/DEL		DARAPRIM TABS KRINTAFEL ² MEFLOQUINE HCL TABS QUININE SULFATE	MC MC/DEL MC/DEL MC MC MC/DEL		ARALEN TABS CHLOROQUINE PHOSPHATE TABS ³ HYDROXYCHLOROQUINE TABS ³ ISONARIF ¹ MALARONE TABS PLAQUENIL TABS	Use PA Form# 20420 1. Ingredients available as preferred without PA. 2. Krintafel is preferred for ≥ 16 years of age. 3. Established users will be grandfathered Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Avoid coadministration of Krintafel® with Organic Cation Transporter 2 (OCT2) and Multidrug and Toxin Extrusion (MATE) substrates (e.g. dofetilide, metformin).
ANTHELMINTICS	MC/DEL MC/DEL MC/DEL		ALBENDAZOLE PRAZIQUANTEL TAB STROMECTOL TABS	MC MC MC/DEL		ALBENZA TABS EMVERM BILTRICIDE TABS	Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIBIOTICS - MISC.	MC MC MC MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC		AZACTAM SOLR COLY-MYCIN-M SOLR COLISTIMETHATE SODIUM SOLR FIRVANQ ⁴ FUROXONE TABS METRONIDAZOLE ¹ PENTAMIDINE ISETHIONATE SOLR SOLOSEC TRIMETHOPRIM TABS VANCOMYCIN 5GM INJ. VANCOMYCIN CAPS XIFAXAN 200mg	MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC MC		AEMCOLO COLISTIMETHATE SODIUM SOLR CAYSTON ³ FLAGYL CAPS FLAGYL TABS FLAGYL ER TBCR KETEK METRONIDAZOLE 375MG CAPS ¹ METRONIDAZOLE 750MG TABS ¹ NEBUPENT SOLR REBYOTA ⁵ TINDAMAX VANCOMYCIN 10GM INJ. ² XENLETA XIFAXAN VOWST ⁵	1. 375mg caps and 750mg tabs are non-preferred. Please use available preferred strengths(250mg & 500mg tabs) to obtain required dose without PA. 2. Please use multiple 5gm which are preferred to obtain dose without PA. 3. Clinical PA is required to establish CF diagnosis and medical necessity. Prior trial and failure of preferred Tobi before approval will be granted. 4. Quantity limit of one per 150ml bottle. 5. For the treatment of patients 18 years of age and older. Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. 1. For macrolide resistant infections when quinolones inappropriate DDI: Ketek is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with either Enblex 15mg or Vesicare 10mg or carbamazepine. Cayston is only indicated to improve respiratory symptoms in CF patients with Pseudomonas aeruginosa. Dosing limits, as should be given TID X28 days (followed by 28 days OFF Cayston therapy). A bronshodilator should be used before administration of Cayston. Xenleta will be considered for the treatment of adults with community-acquired bacterial pneumonia (CABP) caused by the following susceptible microorganisms: Streptococcus pneumoniae, Staphylococcus aureus (methicillin-susceptible isolates), Hemophilus influenzae, Legionella pneumophila, Mycoplasma pneumoniae, and Chlamydia pneumoniae. Vowst: To prevent the recurrence of Clostridioides difficile infection (CDI) in individuals 18 years of age and older following antibacterial treatment for recurrent CDI (rCDI). Rebyota: For the prevention of recurrence of Clostridioides difficile infection (CDI) in individuals 18 years of age and older following antibiotic treatment for recurrent CDI. The limitation of use is that Rebyota® is not indicated for treatment of CDI.
CARBAPENEMS				MC MC MC/DEL MC/DEL		INVANZ SOLR MERREM SOLR PRIMAXIN RECARBRIO	Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

LINCOSAMIDES / OXAZOLIDINONES / LEPROSTATICS	MC/DEL MC/DEL MC/DEL MC	CLEOCIN SOLN CLEOCIN SUSR CLINDAMYCIN HCL 150CAPS DAPSONE TABS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL	8 8 8 8 8 9 9	CLEOCIN CAPS CLINDAMYCIN HCL 300CAPS ¹ SIVEXTRO VIBATIV LINEZOLID TABS ZYVOX SUSR ZYVOX TABS	1. Use multiple 150's for Clindamycin instead of 300's. Use PA Form# 30820 for Zyvox & Vibativ Use PA Form# 20420 for all others	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. For Zyvox or Vibativ, please see the criteria listed in the Antibacterial Antibiotics PA form.
ANTI INFECTIVE COMBO'S - MISC.	MC/DEL MC/DEL MC/DEL MC/DEL	ERYTHROMYCIN/SULF SUSR SEPTRA/DS TABS SULFAMETHOXAZOLE/TRIMETH TRIMETHOPRIM/SULFAMETHOXA	MC MC		BACTRIM DS TABS VABOMERE ¹	Use PA Form# 20420 1. For the treatment of patients ≥ 18 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIPROTOZOALS	MC/DEL MC/DEL	BENZNIDAZOLE ² LAMPIT ²	MC		ALINIA ¹	1. Alina is preferred for children less than 12 years of age. 2. Clinical PA required for appropriate diagnosis. Use PA Form# 20420	Benznidazole is indicated for pediatric patients 2 to 12 years of age for the treatment of Chagas disease (American trypanosomiasis) caused by Trypanosoma cruzi.
ANTI - FUNGALS							
ANTIFUNGALS - ASSORTED	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	ANCOBON CAPS FLUCONAZOLE ¹ KETOCONAZOLE TABS ⁷ NYSTATIN TERBINAFINE TABS ⁴ VORICONAZOLE TABS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC	6 6 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	LAMISIL TABS ⁴ ITRACONAZOLE BREXAFEMME CRESEMBA ⁹ GRIFULVIN V TABS GRISEOFULVIN SUSP GRISEOFULVIN ULTRAMICROSI TABS GRIS-PEG TABS REZZAYO ⁹ SPORANOX SOLN ² SPORANOX PULSEPAK CAPS ³ SPORANOX CAPS ³ DIFLUCAN ERAXIS INJ ⁶ GRIFULVIN SUSP ONMEL NOXAFIL ⁵ TOLSURA VFEND TABS VIVJOA	See quantity limit table. Non-preferred products must be used in specified step order. Continue to use Anti-Fungal PA form for non-preferred products. 1. QL--1/every 7-day period (150mg only). 2. Sporanox QL 300cc/month with PA. See quantity limit table. 3. Sporanox QL 30/month with PA. 4. Quantity limit of one tablet daily. Please see dosage consolidation list. 5. Approved if immuno suppressed/ HIV or if the member has failed a 7 day trial of a preferred antifungal therapy. 6. Eraxis will be approved if submitting with documentation that it was initiated during a hospitalization and this request is to finish the hospital course. 7. Quantity limits allowing 30 day supply without PA. PA will be required if using > 30 days.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. The other criteria are listed on the Antifungal PA form including the required proof of a non-cosmetic fungal infection. DDI: Any Griseofulvin will now be non-preferred and require prior authorization if it is currently being used in combination with either Prevacid, pantoprazole, Prilosec, or any currently non preferred PPI. DDI: Sporanox is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for current use with Enablex 15mg, Vesicare 10mg, Prandin, Prevacid, pantoprazole, Prilosec, or any currently non preferred PPI, due to a significant drug-drug interaction. DDI: Vfend is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with Warfarin. DDI: Fluconazole (except 150mg strength) will now be non-preferred and require prior authorization if it is currently being used with glimepiride (Amaryl), Enablex 15mg, or Vesicare 10mg. Diflucan is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with either glimepiride (Amaryl), Enablex 15mg, or Vesicare 10mg. DDI: Fluconazole will require prior authorization if being used in combination with Plavix or Warfarin. DDI: Ketoconazole will now be non-preferred and require prior authorization if they are currently being used in combination with any of the following medications: Prevacid, Pantoprazole, Plavix, Onglyza, Enablex 15mg, Vesicare 10mg, Latuda, Cometriq, Tafenlar or Omeprazole. Rezzayo: In patients 18 years of age or older who have limited or no alternative options for the treatment of candidemia and invasive candidiasis.

8. For children < 18, quantity limits allows 8 weeks supply without PA. PA will be required if using > than 8 weeks. If 18 and older PA will be required for any quantity. Not approving for Onychomycosis indication.

9. For patients ≥ 18years of age

[Use PA Form# 10120](#)

ANTI - VIRALS

ANTI - VIRALS							
ANTIRETROVIRALS	MC/DEL	ABACAVIR TABS	MC/DEL	8	ABACAVIR SOL	<p>Use PA Form# 20420</p> <p>1. Quantity limit of one per day</p> <p>2. Only preferred if Norvir script is in member's profile within the past 30 days of filling Prezista</p> <p>3. Isentress Chewable will only be approved if between the age of 2-12 years old</p> <p>4. Request will require use of the individual</p> <p>5. Clinical PA required.</p> <p>6. Only preferred for post-exposure prophylaxis.</p>	<p>Fuzeon: Prescriber is either an HIV specialist provider or has consulted with one. Documentation of genotype testing issued and shows that there is no other potent, appropriate two or three drug oral regimen available, AND patient has a positive HIV viral load within past 6 months while on his/her current antiretroviral regimen AND the drug will be prescribed with at least two other drugs that are likely to be active based on the genotype testing.</p> <p>DDI: Reyataz requires prior authorization if it is currently being used in combination with either Prevacid, pantoprazole, Prilosec, or any currently non preferred PPI .</p> <p>DDI: Norvir requires prior authorization if it is currently being used in combination with either Enablex 15mg or Vesicare 10mg.</p> <p>DDI: Preferred Crixivan caps requires prior authorization if it is currently being used in combination with either Enablex 15mg or Vesicare 10mg.</p> <p>DDI: The concomitant use of the following drugs with Descovy® is not recommended: tipranavir/ritonavir, St. John's wort, and the antimycobacterials rifabutin, rifampin, or rifapentine.</p> <p>DDI: Administration with the following drugs: the anticonvulsants carbamazepine, oxcarbazepine, phenobarbital, and phenytoin; the antimycobacterials rifampin and rifapentine; proton pump inhibitors such as dexlansoprazole, esomeprazole, lansoprazole, omeprazole, pantoprazole, rabeprazole; systemic dexamethasone (more than a single dose); and St. John's wort with Odefsey is contraindicated.</p> <p>Stribild: PA required; must provider rationale as to why the member's medical need cannot be met with preferred agents, particularly Genvoya or combinations of preferred and agents AND must be antiretroviral treatment-naïve or virologically controlled on current therapy (HIV-1RNA < copies/ml) AND be HBV negative AND not be combined with other anti-retroviral agents.</p> <p>DDI: Tivicay will require prior authorization is used with nevirapine, oxcarbazepine, phenytoin, phenobarbital, carbamazepine, and St. John's wort.</p> <p>DDI: Aatazanavir or darunavir and the following drugs are contraindicated (due to potential for serious and/or life-threatening events or loss of therapeutic effect): alfuzosin, dronedarone, rifampin, irinotecan, dihydroergotamine, ergotamine, methylergonovine, cisapride, St. John's wort, lovastatin, simvastatin, pimozide, nevirapine, sildenafil (when given as Revatio® for treatment of PAH), indinavir, triazolam, or PO midazolam will be non-preferred and require prior authorization if it is currently being used in combination with Tybost.</p> <p>DDI: Combined P-gp, UGT1A1 and strong CYP3A inhibitors may significantly increase plasma concentrations of Sunlenca®. Concomitant administration of Sunlenca® with these inhibitors is not recommended.</p> <p>Sunlenca: In combination with other antiretroviral(s) for the treatment of HIV-1 infection in heavily treatment-experienced adults with multidrug resistant HIV-1 infection failing their current antiretroviral regimen due to resistance, intolerance, or safety considerations.</p>
	MC	APRETUDE	MC/DEL	8	APTIVUS		
	MC/DEL	ATAZANAVIR	MC/DEL	8	CIMDUO		
	MC	ATRIPLA ¹	MC/DEL	8	COMBIVIR TABS		
	MC	BIKTARVY	MC/DEL	8	EDURANT		
	MC	CABENUVA	MC/DEL	8	EPZICOM ¹		
	MC	COMPLERA ¹	MC/DEL	8	FUZEON		
	MC/DEL	DELSTRIGO	MC/DEL	8	INTELENCE		
	MC	DESCOVY ¹	MC/DEL	8	ISENTRESS ³		
	MC	DIDANOSINE	MC/DEL	8	ISENTRESS HD		
	MC/DEL	DOVATO	MC	8	JULUCA		
	MC	EFAVIRENZ TAB	MC	8	KALETRA		
	MC/DEL	EFAVIRENZ CAP	MC/DEL	8	LEXIVA		
	MC	EFAVIRENZ-EMTRICITABINE-TENOFOVIR DF TAB	MC/DEL	8	NEVIRAPINE		
	MC	EMTRIVA ¹	MC	8	NORVIR		
	MC	EPIVIR SOL	MC/DEL	8	PIFELTRO		
	MC/DEL	EVOTAZ ¹	MC	8	RETROVIR		
	MC	GENVOYA ^{1,5}	MC	8	REYATAZ		
	MC/DEL	ISENTRESS 400MG ⁶	MC/DEL	8	SELZENTRY		
	MC/DEL	ISENTRESS CHEW ³	MC	8	STAVUDINE		
	MC/DEL	ISENTRESS POWDER	MC	8	STRIBILD ¹		
	MC/DEL	LAMIVUDINE TABS	MC	8	SUNLENCA ⁵		
	MC/DEL	LAMIVUDINE/ZIDOVUDINE	MC/DEL	8	SYMFI ⁹		
	MC/DEL	LAMIVUDINE SOLN	MC/DEL	8	SYMFI LO ⁵		
	MC/DEL	LOPINAVIR-RITONAVIR SOL	MC/DEL	8	SYM TUZA		
	MC	LOPINAVIR-RITONAVIR TAB	MC	8	TRIUMEQ ^{1,4}		
	MC	ODEFSEY ¹	MC/DEL	8	TRIZIVIR TABS		
	MC/DEL	PREZCOBIX	MC	8	TRUVADA ¹		
	MC	PREZISTA ²	MC/DEL	8	VIRACEPT TABS		
	MC/DEL	RITONAVIR TAB 100MG	MC	8	VITEKTA		
	MC	RUKOBIA ⁹	MC	8	ZERIT		
	MC	SUSTIVA ¹	MC	8	VIDEX EC		
	MC	TIVICAY	MC	8	VIREAD TABS ¹		
	MC	TIVICAY PD	MC/DEL	8	ZIAGEN TABS		
	MC	TROGARZO ⁵	MC/DEL	8	ZIAGEN SOL		
	MC	TYBOST	MC/DEL	9	VIRAMUNE XR		
	MC	VIREAD POW					
	MC/DEL	ZIDOVUDINE					
CYTO-MEGALOVIRUS AGENTS	MC	CIDOFOVIR	MC		VALCYTE TABS	<p>Use PA Form# 20420</p> <p>1. Must show failure or</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>
	MC	FOSCARNET SODIUM	MC/DEL		FOSCAVIR		
	MC/DEL	GANCICLOVIR	MC/DEL		LIVTENCITY ¹		

	MC/DEL		VALGANCICLOVIR	MC/DEL		PREVYMIS	contraindication to all the following ganciclovir, valganciclovir, cidofovir and foscarnet before Livtency will be approved.	Prevymis: Documentation that member is high-risk for CMV reactivation as defined by transplant guidelines or that there has been significant myelosuppression by one of the preferred agents. DDI: Livtency is a substrate of CYP3A4. Coadministration of Livtency® with strong inducers of CYP3A4 is not recommended, except for selected anticonvulsants.
HERPES AGENTS	MC/DEL MC/DEL		ACYCLOVIR VALACYCLOVIR HCL	MC/DEL MC MC/DEL MC MC/DEL	8 8 8 8 9	FAMCICLOVIR ¹ SITAVIG ZOVIRAX ¹ VALTRES TABS ¹ FAMVIR TABS ¹	1. Must fail Acyclovir and Valacyclovir before non-preferred products in step order. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
INFLUENZA AGENTS	MC MC MC/DEL		AMANTADINE CAPS RELENZA DISKHALER AEPB OSELTAMIVIR ¹	MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL		AMANTADINE TABS FLUMADINE TABS FLUMIST RIMANTADINE HCL TABS TAMIFLU ¹ TAMIFLU SUS XOFLUZA	1. Tamiflu and Osetamivir 10 caps or 60cc's per month. Will be audited for presence of positive influenza tests in patient or family member. Use PA Form# 20420 for all others	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
IMMUNE SERUMS								
IMMUNE SERUMS	MC		HYPERRHO INJ					
HEPATITIS AGENTS								
HEPATITIS C AGENTS	MC MC MC/DEL MC/DEL MC/DEL MC MC/DEL		SOFOSBUVIR/VELPATASVIR ² (Authorized generic labeler 72626 Asegua Therapeutics) MAVYRET ² PEGASYS KIT ¹ PEGASYS SOLN PEG-INTRON KIT ¹ RIBAVIRIN RIBASPHERE	MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC MC/DEL		COPEGUS TABS DAKLINZA EPCLUSA ² HARVONI ² REBETOL CAPS RIBAPAK SOVALDI ² VIEKIRA PAK ² VIEKIRA XR ² VOSEVI ZEPATIER ²	1. Dosing limits apply, please see dosage consolidation list. 2. Approvals will require clinical PA. Please see the Hepatitis PA form for criteria Use PA Form #10700	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Olysio will require a prior authorization if it is currently being used in combination with drugs known to be significant CYP3A4 inhibitors (ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, nelfinavir, ritonavir, atazanavir, saquinavir and telithromycin).
HEPATITIS AGENTS - MISC.				MC		ACTIMMUNE	Use PA Form# 20420	Approved for chronic granulomatous disease, osteopetrosis and idiopathic pulmonary fibrosis.
HEPATITIS B ONLY	MC/DEL MC		ENTECAVIR TENOFVIR	MC MC MC MC		BARACLUDE HEPSERA TABS TYZEKA VEMLIDY	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Baraclude is indicated for treatment of chronic Hep B virus (HBV) in adults with: evidence of active viral replication AND either evidence of persistent elevation in serum aminotransferases (ALT or AST) or histologically active disease, Patient is 16 years of age or older. Boxed warning: Use not recommended for those co-infected with HIV and HBV who are not also receiving highly active antiretroviral therapy (HAART). Vemlidy® remain non-preferred and require prior authorization and be available to those who have evidence of bone loss or renal insufficiency or who are unable to tolerate or who have failed on preferred medications.
RSV PROPHYLAXIS								
RSV PROPHYLAXIS				MC		SYNAGIS ¹	Use PA Form# 30120 1. MaineCare will approve Synagis PA's for start date of November 29, 2021 for infants who meet the	Please see the criteria listed on the Synagis PA form.

patients who meet the guidelines. PA will be approved for max of 5 doses. Maximum 1 dose/30 days. MaineCare will start accepting PAs November 1, 2021."

MS TREATMENTS

MULTIPLE SCLEROSIS - INTERFERONS	MC MC/DEL MC		AVONEX KIT ¹ BETASERON SOLR ¹ REBIF SOLN ¹	MC MC/DEL		PLEGRIDY ¹ EXTAVIA	1. Clinical PA is required to establish diagnosis and medical necessity. Use PA Form# 20430	Non-Preferred drugs must be tried in step-order and failed due to lack of efficacy or intolerable side effects before lower ranked non-preferred drugs will be approved , unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
----------------------------------	--------------------	--	---	--------------	--	----------------------------------	--	--

MULTIPLE SCLEROSIS - NON-INTERFERONS	MC MC/DEL MC/DEL MC/DEL MC MC MC		COPAXONE DALFAMPRIDINE ER DIMETHYL FUMARATE CAP FINGOLIMOD CAP ² KESIMPTA ² TERIFLUNOMIDE TAB ² TYSABRI ^{1,2}	MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC MC MC	8 8 8 8 8 8 8 8 8 8 8 8 8	AMPYRA AUBAGIO BAFIERTAM BRIUMVI GILENYA GLATOPIA MAVENCLAD ³ MAYZENT OCREVUS ² PONVORY ² TASCENSO ODT ^{2,4} TECFIDERA VUMERITY ZEPOSIA	1. Providers must be enrolled in the TOUCH Prescribing program, a restricted distribution program. Clinical PA is required to establish diagnosis and medical necessity. 2. Clinical PA is required to establish diagnosis and medical necessity. 3. Due to safety profile, use of Mavenclad® is generally recommended for patients who have had an inadequate response to, or are unable to tolerate, an alternate drug indicated for the treatment of MS 4. For the treatment of patients 10 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Mavenclad will require multiple trials of preferred agents including Mayzent for secondary progressive disease. DDI: Due to significant increases in exposure to siponimod, concomitant use of Mayzent® and drugs that cause moderate CYP2C9 and moderate or strong CYP3A4 inhibition is not recommended. Ponvory: Before initiation of Ponvory® treatment, assess the following: •Complete Blood Count (CBC)- Obtain a recent (i.e. within the last 6 months) CBC, including lymphocyte count. •Cardiac Evaluation- oObtain an electrocardiogram (ECG) to determine whether pre-existing conduction abnormalities are present. In patients with certain pre-existing conditions, advice from a cardiologist should be sought and first-dose monitoring is recommended. oDetermine whether patients are taking drugs that could slow heart rate of atrioventricular (AV) conduction. •Liver Function Tests- Obtain recent (i.e. within the last 6 months) transaminase and bilirubin levels. •Ophthalmic Evaluation- Obtain an evaluation of the fundus, including the macula. •Current or prior medications with immune system effects- If patients are taking anti-neoplastic, immunosuppressive, or immune-modulating therapies, or if there is a history of prior use of these drugs, consider possible unintended additive immunosuppressive effects before starting treatment with Ponvory®. •Vaccinations- Test for antibodies to varicella zoster virus (VZV) before starting Ponvory®; VZV vaccination of antibody-negative patients is recommended prior to commencing treatment with Ponvory®. If live attenuated vaccine immunizations are required, administer at least 1 month prior to initiation of Ponvory®. Mayzent for Relapsing forms of MS: multiple trials of preferred agents, including an intravenous MS product. Mayzent for Active secondary progressive disease: prior trials of two preferred agents are required. Use PA Form# 20430
--------------------------------------	--	--	---	--	---	--	---	---

MULTIPLE SCLEROSIS - MISC				MC		ZINBRYTA ¹	1. The safety and efficacy of use in children under the age of 17 years have not been established. Use PA Form #20430	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists
---------------------------	--	--	--	----	--	-----------------------	--	---

ASSORTED NEUROLOGICS

NEUROLOGICS - MISC.	MC MC MC MC		BOTOX ^{2,4} DYSPORT ⁴ PROSTIGMIN TABS PYRIDOSTIGMINE	MC/DEL MC MC MC/DEL		FIRDAPSE MESTINON MYOBLOC ¹ RUZURGI ³	1. Approval will be limited to Cervical dystonia. 2. Please see botulinum PA	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
---------------------	----------------------	--	---	------------------------------	--	--	---	--

				<p>MC</p> <p>MC</p> <p>MC</p> <p>MC/DEL</p>	<p>SKYSONA^{4,6}</p> <p>VYVGART⁵</p> <p>VYVGART HYTRULO⁵</p> <p>XEOMIN²</p>	<p>form for additional criteria</p> <p>3. For the treatment of patients between ages 6-16 years of age.</p> <p>4. Clinical PA required.</p> <p>5. For adult patients who are anti-acetylcholine receptor (AChR) antibody positive.</p> <p>6. For the treatment of patients between ages 4-17 years of age.</p> <p>Use PA Form# 10210</p>	<p>Failed/did not tolerate therapeutic trials fo muscle relaxants, unless contraindicated, including but not limited to baclofen, cyclobenzaprine, orphenadrine, Skelaxin, and tizanidine.</p> <p>Migraine: Consideration for Botox approvals will only be made after failures of required trials of the following preferred medications: tricyclic or venlafaxine,beta blocker,valproic acid ,topiramate.</p> <p>Firdapse is recommended for the treatment of Lambert-Eaton myasthenic syndrome (LEMS) in adults.</p> <p>Ruzurgi is recommended for the treatment of Lambert-Eaton myasthenic syndrome (LEMS) in patients 6 years to less than 17 years of age.</p>
NEUROLOGICS- hATTR AGENTS				<p>MC</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC/DEL</p> <p>MC/DEL</p>	<p>AMVUTTRA¹</p> <p>ONPATTRO¹</p> <p>TEGSEDI¹</p> <p>VYNDAMAX¹</p> <p>VYNDAQEL¹</p>	<p>1. PA required for appropriate diagnosis.</p> <p>Use PA Form# 20420</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.</p> <p>Tegsedi® should be non-preferred and approved for patients for whom other treatments, including Onpattro®, have been ineffective.</p> <p>Vyndamax will be considered for the treatment of the cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) in adults to reduce cardiovascular mortality and cardiovascular-related hospitalization</p>
NEUROLOGICS- SMA	<p>MC</p> <p>MC</p> <p>MC</p>		<p>GENE</p> <p>ZOLGENSMA¹</p> <p>NON-GENE</p> <p>EVRYSDI^{1,2}</p> <p>SPINRAZA¹</p>		<p>GENE</p> <p>NON-GENE</p>	<p>1. Clinical PA is required to establish diagnosis and medical necessity</p> <p>2. For patients 2 months of age and older.</p> <p>Use PA Form# 20420</p>	<p>Zolgensma: The patient is less than 2 years of age AND The diagnosis is spinal muscular atrophy (SMA) AND The patient has bi-allelic mutations of the SMN1 gene AND The patient does not have advanced SMA (e.g. complete paralysis of limbs or permanent ventilator dependence) AND Medication is prescribed per the dosing</p> <p>Spinraza: The diagnosis is spinal muscular atrophy (SMA) type 1, 2, or 3 (results of genetic testing must be submitted) AND The patient has at least 2 copies of the SMN2 gene AND The prescriber is a neurologist, pulmonologist, or other physician with expertise in treating SMA AND Baseline motor ability has been established using one of the following exams: Hammersmith Infant Neurological Exam (HINE) Hammersmith Functional Motor Scale Expanded (HFMSE) Upper Limb Module Test (non-ambulatory) Children’s Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP INTEND) AND Prior to starting therapy, and prior to each dose, the following laboratory tests will be conducted: Treating provider attests the member has a platelet count > 50,000/ml or greater Treating provider agrees to do platelet count and coagulation test before each dose Treating provider agrees to do a quantitative spot urine protein test before each dose Concomitant use of Spinraza and Zolgensma is investigational and will not be approved AND Use of Spinraza after gene replacement therapy, including Zolgensma is investigational and will not be approved Note: Initial approval will be granted for 4 loading doses (the first 3 loading doses should be administered at 14-day intervals; the 4th loading dose should be administered 30 days after the 3rd dose). Renewal may be granted for up to 12 months with a maximum of 3 doses approved per year (12mg (5ml) every 4 months). For therapy continuation, clinical documentation must be submitted documenting improvement or maintenance of motor ability OR slower progression of disease than would otherwise be expected.</p> <p>Use PA Form# 20420</p>
NEUROLOGICS- RETT SUNDROME				<p>MC</p>	<p>DAYBUE^{1,2}</p>	<p>1.Clinical PA required for appropriate diagnosis</p> <p>2. For the treatment of patients 2 years of age and older.</p> <p>Use PA Form# 20420</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>
ALS DRUGS	<p>MC/DEL</p>		<p>RILUZOLE</p>	<p>MC</p>	<p>EXSERVAN</p>		<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical</p>

	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		METHYLPREDNISOLONE TABS PREDNISOLONE PREDNISONONE SOLU-CORTEF SOLR SOLU-MEDROL SOLR	MC MC		STERAPRED TABS ZILRETTA		DDI: All preferred steroids will require clinical PA for patients over 60 that are currently on fluoroquinolone therapy.
HORMONE REPLACEMENT THERAPIES								
ANDROGENS / ANABOLICS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ANDRODERM PT24 ANDROGEL 1% ANDROGEL PUMP 1.62% DANAZOL CAPS TESTOSTERONE CYP	MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL		ANADROL-50 ANDRO LA 200 OIL ANDROGEL PACKETS 1.62% ANDROID CAPS AXIRON DELATESTRYL OIL DEPO-TESTOSTERONE OIL FORTESTA HALOTESTIN TABS JATENZO METHITEST TAB METHYLTESTOSTERONE CAP OXANDROLONE STRIANT MUC ER TESTIM TESTOSTERONE GEL PACKETS TESTOSTERONE SOL TESTRED CAPS TLANDO VOGELXO XYOSTED	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Additionally, laboratory evidence of a testosterone deficiency must be supplied. One of each dosage form should be tried (tablet, injection, and topical) Oxandrolone: Weight gain (adjunctive therapy): Adjunctive therapy to promote weight gain after weight loss following extensive surgery, chronic infections, or severe trauma, and in some patients who, without definite pathophysiologic reasons, fail to gain or to maintain normal weight. Other indications included in manufacturer labeling: Adjunctive therapy to offset protein catabolism with prolonged corticosteroid administration. Requirement for documentation of weight loss over two readings- Patient has involuntary weight loss of more than 10% of total body weight in less than four months) and, BMI < 18.5 (Normal BMI = 18.5 to 24.9)
ESTROGENS - PATCHES / TOPICAL	MC MC/DEL		EVAMIST MINIVELLE PATCH	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	5 8 8 8 8 8	ESTRADIOL PTWK DIVIGEL ¹ CLIMARA PTWK ELESTRIN ¹ MENOSTAR PATCH VIVELLE-DOT PTTW	1. Step order drugs must be used in specified step order.	Approved for failures on multiple oral estrogen agents after 90 day trials or if unable to swallow any oral medication.
ESTROGENS - TABS	MC/DEL MC/DEL		ESTRADIOL PREMARIN TABS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC		ENJUWIA ESTRADIOL-NORETHINDRONE ESTRACE TABS ESTRATAB TABS MENEST TABS NORETHINDRON-ETHINYL ORTHO-EST TABS	Must fail preferred products before non-preferred products.	Preferred drugs must be tried for at least 90 days and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ESTROGEN COMBO'S	MC/DEL MC/DEL MC/DEL MC/DEL		ANGELIQ COMBIPATCH PTTW PREMPHASE TABS PREMPRO TABS	MC/DEL MC/DEL MC MC/DEL MC/DEL		FEMHRT 1/5 TABS ¹ FYAVOLV LOPREEZA TAB ORTHO-PREFEST TABS ¹ SYNTEST H.S. TABS ¹	1. Must fail Premphase and Prempro products before non preferred products.	Preferred drugs must be tried for at least 90 days and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PROGESTINS	MC/DEL MC/DEL MC MC		MEDROXYPROGESTERONE ACETA ¹ NORETHINDRONE ACETATE TABS ¹ 17-ALPHA HYDROXYPROGESTERONE PWDR PROGESTERONE CAPS	MC/DEL MC MC MC/DEL MC/DEL		AYGESTIN TABS CYCRIN TABS PROGESTERONE POWD PROMETRIUM CAPS PROVERA TABS	1. Must fail Medroxyprogesterone and Norethindrone products before non-preferred	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

[Use PA Form# 20420](#)

ENDOMETRIOSIS

CENTRAL PRECOCIOUS PUBERTY AGENTS	MC	FENSOLVI ¹				1. For pediatric patients 2 years of age and older with central precocious puberty (CPP).	
ENDOMETRIOSIS- NASAL	MC/DEL	SYNAREL (NASAL) SPRAY					Synarel is also indicated for central precocious puberty Use PA Form# 20420
ENDOMETRIOSIS/ UTERINE FIBROIDS- ORAL	MC/DEL MC	ORILISSA ¹ MYFEMBREE ^{1,2}	MC		ORIAHNN ¹	1. Prior treatment of NSAID and hormonal contraceptives required 2. Limited to 24 months due to the risk of continued bone loss, which may not be reversible.	Use PA Form# 20420
ENDOMETRIOSIS- INJECTABLE	MC/DEL	DEPO-SUBQ PROVERA 104					Use PA Form# 20420

CONTRACEPTIVES

CONTRACEPTIVES - PROGESTIN ONLY	MC/DEL MC/DEL MC MC MC/DEL	CAMILA TABS ERRIN INCASSIA TAB HEATHER TAB NORETHINDRONE ACETATE 0.35MG TABS	MC/DEL MC/DEL MC MC/DEL		JOLIVETTE NORA-BE TABS ORTHO MICRONOR TABS SLYND		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. If member experienced adverse reactions, consider using Oral Contraceptives from other groups. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with Tracleer. Use PA Form# 20420
CONTRACEPTIVES - INJECTABLE	MC/DEL	MEDROXYPROGESTERONE ACETATE 150mg IM	MC/DEL		DEPO-PROVERA 150 mg SUSP		The preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Use PA Form# 20420
CONTRACEPTIVE - EMERGENCY	MC/DEL MC MC MC MC MC/DEL MC MC/DEL MC MC/DEL	ELLA ENCONTRA ONE STEP ECONTRA EZ NEW DAY OPCION OPTION 2 MY CHOICE MY WAY LEVONORGESTREL NEXT CHOICE ¹				1. Allowed 2 tablets per 30 days without PA	Due to the extensive list of products, any covered emergency contraceptive product preferred is and available without a PA. Use PA Form# 20420
CONTRACEPTIVES - PATCHES/ VAGINAL PRODUCTS	MC MC MC MC/DEL	ELURYNG ¹ NUVARING RING ¹ TWIRLA XULANE ²	MC MC MC		ANNOVERA PHEXXI ZAFEMY	1. Quantity limit allowing 1 every 28 days with out PA. 2. Dose limits apply allowing 3 patches per 28 days supply.	Approved if adequate clinical reason given why patient unable to comply with other preferred agents including long acting injectable. Use PA Form# 20420

CONTRACEPTIVES- LONG ACTING REVERSIBLE	MC/DEL		MIRENA	MC/DEL MC MC MC/DEL MC/DEL	KYLEENA LILETTA NEXPLANON PARAGARD SKYLA		
CONTRACEPTIVES - MONOPHASIC COMBINATION O/C'S	MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		APRI TABS AVIANE TABS BALZIVA CRYSSELLE-28 TABS DESOGEN TABS ESTARYLLA TAB HAILEY FE TAB ISIBLOOM TAB JUNEL FE TAB LARIN FE TAB LESSINA TAB LEVORA-28 TAB MILI TAB NORGESTIMATE-ETHINYL ESTRADIOL TAB MIBELAS 24 FE TAB MICROGESTIN FE TAB RECLIPSEN SAFYRAL TAB SPRINTEC 28 TABS YASMIN 28 TABS YAZ	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	BEYAZ BREVICON-28 TABS LESSINA-28 TABS LEVORA LOESTRIN FE 1/20 TABS LOESTRIN 1.5/30-21 TABS MICROGESTIN FE TABS LOESTRIN 1/20-21 TABS LO/OVRAL 21 TABS LO/OVRAL 28 TABS NEXTSTELLIS NORDETTE-28 TABS NORTREL OCELLA OVRAL PORTIA-28 TABS SAFYRAL ZOVIA	Use PA Form# 20420 If member experienced adverse reactions, consider using Oral Contraceptives from other groups.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. If member experienced adverse reactions, consider using Oral Contraceptives from other groups. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with Tracleer.
CONTRACEPTIVES - BI-PHASIC COMBINATIONS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC/DEL		AZURETTE TAB CAMRESE CAMRESE LO DESOGESTREL/ ETH/ ESTRAD 0.15/30mcg KARIVA TABS LO LOESTRIN FE PIMTREA TAB NORETHINDRONE-ETH ESTRADIOL TAB 0.5-35/1-35 SIMPESSE TBDSPK 3MO VIORELE TAB	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	LOSEASONIQUE	If member experienced adverse reactions, consider using Oral Contraceptives from other groups.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. If member experienced adverse reactions, consider using Oral Contraceptives from other groups. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with Tracleer.
CONTRACEPTIVES - TRI-PHASIC COMBINATIONS	MC/DEL MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC		ENPRESSE NORGESTIMATE-ETHINYL ESTRADIOL TAB TRIPHASIL 28 TABS TRI-LO-MILI TAB TRI-LO-ESTARYLLA TAB TRI-ESTARYLLA TRI-SPRINTEC TAB TRI-LO-SPRINTEC TRINESSA	MC/DEL MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC	NORTREL 7/7/7 ORTHO TRI-CYCLEN LO TABS	If member experienced adverse reactions, consider using Oral Contraceptives from other groups.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. If member experienced adverse reactions, consider using Oral Contraceptives from other groups. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with Tracleer.
CONTRACEPTIVES - MULTI-PHASIC COMBINATIONS				MC	NATAZIA	Use PA Form# 20420	

VASOMOTOR SYMPTOMS AGENTS

VASOMOTOR SYMPTOMS AGENTS				MC/DEL		VEOZAH		<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>DDI: Avoid concomitant use of Veozah with drugs that are weak, moderate or strong CYP1A2 inhibitors.</p>
Use PA Form# 20420								

DIABETES SUPPLIES

DIABETIC- SUPPLIES			CONTINUOUS GLUCOSE MONITORING ^{1,2} DIABETIC- LANCETS DIABETIC- LANCING DEVICES DIABETIC- LANCING DEVICES DIABETIC- PEN NEEDLES DIABETIC- SYRINGES DIABETIC- TEST STRIPS DIABETIC- METERS				<p>1. Clinical PA is required to establish diagnosis and medical necessity.</p> <p>2. Dosing limits apply. Please refer to Dose consolidation list.</p>	<p>Please refer to the MaineCare Preferred Diabetic Supply List available at www.mainearepd.org</p> <p>Continuous Glucose Monitoring Criteria: Patient has a diagnosis of Diabetes Mellitus AND Practitioner feels patient has sufficient training to use CGM</p> <ul style="list-style-type: none"> • 2 years of age or older for Dexcom G6, ≥ 14 years for Medtronic Guardian, or ≥ 4 years for Freestyle Libre 2. • At least one of the following are documented: <ul style="list-style-type: none"> o Hypoglycemic unawareness o Treated with insulin (at least 1X day) o Has history of problematic hypoglycemia with documentation of at least one recurrent level 2 hypoglycemic events, or 1 level 3 hypoglycemic event • Approval of non-preferred products will be limited to cases where the CGM is directly integrated with the patient's insulin pump. The make and model of pump must be documented on the prior authorization.
Use PA Form#20420								

DIABETES THERAPIES

DIABETIC - INSULIN	MC/DEL MC MC MC MC MC MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		<p>APIDRA</p> <p>HUMALOG KWIKPEN INJ 100/ML</p> <p>HUMALOG JUNIOR KWIKPEN 100/ML</p> <p>HUMALOG MIX 75/25</p> <p>HUMALOG 50/50 VIAL</p> <p>HUMULIN INJ 70/30 KWIKPEN</p> <p>HUMULIN INJ 70/30</p> <p>HUMULIN R INJ U-500</p> <p>INSULIN ASPART PROT MIX 70-30</p> <p>INSULIN ASPART</p> <p>INSULIN LISPRO</p> <p>LANTUS SOLN</p> <p>LEVEMIR</p> <p>NOVOLOG</p> <p>NOVOLOG MIX</p> <p>NOVOLOG MIX 70/30 FLEXPEN</p>	MC/DEL MC/DEL MC MC/DEL MC MC MC MC MC/DEL MC		<p>ADMELOG</p> <p>AFREZZA¹</p> <p>BASAGLAR</p> <p>FIASP</p> <p>HUMALOG KWIKPEN U-200</p> <p>HUMULIN INJ 50/50</p> <p>HUMULIN N INJ U-100</p> <p>HUMULIN R U-100</p> <p>LYUMJEV</p> <p>NOVOLIN</p> <p>RELION</p>	<p>Use PA Form# 20420</p> <p>1. Not to be as a monotherapy. Obtain lab values of pulmonary function and recent smoking history</p> <p>2. For the treatment of patients ≥3 years of age</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>
DIABETIC - PENFILLS	MC MC MC MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		<p>HUMALOG MIX KWIK 50/50</p> <p>HUMALOG MIX INJ 75/25 KWP</p> <p>HUMALOG KWIK INJ 100/ML</p> <p>HUMALOG KWIK INJ 200/ML</p> <p>HUMULIN R U-500 KWP</p> <p>INSULIN ASPART PROT MIX 70-30 PEN</p> <p>INSULIN ASPART PEN</p> <p>INSULIN LISPRO KWIKPEN U-100</p> <p>LANTUS SOLOSTAR</p> <p>LEVEMIR FLEXTOUCH</p> <p>LEVEMIR FLEXPEN</p> <p>NOVOLOG MIX PENFILL</p> <p>NOVOLOG PENFILL SOLN</p> <p>NOVOLOG FLEXPEN</p> <p>NOVOLOG MIX 70/30 VIAL</p>	MC MC/DEL MC MC/DEL		<p>APIDRA OPTICLIK PEN</p> <p>NOVOLIN 70/30 PEN</p> <p>REZVOGLAR KWIKPEN</p> <p>TRESIBA</p>	<p>Use PA Form# 20420</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>

	MC/DEL MC/DEL		TOUJEO MAX SOLOSTAR TOUJEO SOLOSTAR					
DIABETIC - DPP- 4 ENZYME INHIBITOR	MC/DEL MC/DEL		JANUVIA ^{1,2} TRADJENTA ²	MC/DEL MC/DEL MC/DEL		NESINA ONGLYZA ² QTERN	1. Preferred if therapeutic doses of metformin are seen in members drug profile for at least 60 days within the past 18 months or if phosphate binder is currently seen in the members drug profile. 2. Dosing limits apply. Please refer to Dose consolidation list. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Onglyza 5mg will require a prior authorization if it is currently being used in combination with drugs known to be significant CYP3A4 inhibitors (ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, nelfinavir, ritonavir, atazanavir, saquinavir and telithromycin).
DIABETIC - DPP- 4 ENZYME INHIBITOR-COMBO	MC/DEL MC/DEL MC/DEL		JANUMET ^{1,2} JANUMET XR ^{1,2} JENTADUETO ¹	MC/DEL MC/DEL MC MC/DEL		JENTADUETO XR KAZANO KOMBIGLYZE XR OSENI	1. Preferred if therapeutic doses of metformin are seen in members drug profile for at least 60 days within the past 18 months or if phosphate binder is currently seen in the members drug profile. 2. Dosing limits apply. Please refer to Dose consolidation list. Use PA Form# 20420	
DIABETIC - LANCET-LANCET DEVICE							Use PA Form# 20420	Please refer to the MaineCare Preferred Diabetic Supply List available at www.mainearepdl.org
DIABETIC - SYRINGES-NEEDLES							Use PA Form# 20420	Please refer to the MaineCare Preferred Diabetic Supply List available at www.mainearepdl.org
DIABETIC - OTHER				MC/DEL MC		CYCLOSET SYMLIN	Use PA Form #20420 for all others	
SGLT 2 INHIBITORS	MC/DEL MC/DEL MC/DEL		FARXIGA INVOKANA ¹ JARDIANCE	MC/DEL		STEGLATRO	1. Dosing limits apply please refer to Dose Consolidation List Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
SGLT 2 INHIBITOR COMBINATIONS	MC/DEL MC/DEL MC/DEL		INVOKAMET SYNJARDY XIGDOU XR	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		GLYXAMBI INVOKAMET XR SEGLUROMET STEGLUJAN SYNJARDY XR TRIJARDY XR	Use PA Form# 20420	Preferred drugs must be tried for at least 3 months at full therapeutic doses and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Glyxambi /Xigduo XR- Verify prior trials and failures or intolerance of preferred treatments from other diabetic categories Synjardy® XR is not recommended for patients with type 1 DM or for the treatment of diabetic ketoacidosis.
DIABETIC MONITOR	MC MC MC MC		ONE TOUCH ULTRA 2 KIT ONE TOUCH ULTRA MINI KIT TRUE METRIX TRUETRACK	MC MC MC MC MC		ACCUCHECK ASCENSIA ASSURE CONTOUR BREEZE Z EXACTECH	Use PA Form# 20420	Effective October 25th 2007, approvals for all non preferred meters/ test strips will require medical necessity documenting clinically significant features that are not available on any of the preferred meters.

				MC		FREESTYLE INSULINX		
				MC		FREESTYLE LITE SYSTEM KIT		
				MC		ONE TOUCH ULTRA SMART KIT		
				MC		PRECISION XTRA METER		
				MC		PRODIGY		
DIABETIC TEST STRIPS	MC MC MC		ONE TOUCH ULTRA ¹ TRUE METRIX TRUETRACK	MC MC MC MC MC MC MC MC MC MC MC		ACCUCHECK ASCENSIA ASSURE CONTOUR BREEZE Z EXACTECH FREESTYLE FREESTYLE LITE FREESTYLE INSULINX ONE TOUCH DELICA PRECISION XTRA PRODIGY	1. Only 50 ct & 100 ct package size. Use PA Form# 20420	Effective October 25th 2007, approvals for all non preferred meters/ test strips will require medical necessity documenting clinically significant features that are not available on any of the preferred meters.
INCRETIN MIMETIC	MC MC MC/DEL		BYETTA TRULICITY VICTOZA	MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL	5 5 8 8 8 8 8	OZEMPIC RYBELSUS ADLYXIN BYDUREON BCISE MOUNJARO SOLIQUA XULTOPHY		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Soliqua must try both insulin and a preferred incretin mimetic and have a medical necessity for use that is not based on convenience or simply due to the fact that one injection is needed instead of two.
DIABETIC - ORAL SULFONYLUREAS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		CHLORPROPAMIDE TABS GLIMEPIRIDE GLIPIZIDE TABS GLIPIZIDE ER TABS GLYBURIDE MICRONIZED TABS GLYBURIDE TABS ¹ TOLAZAMIDE TABS TOLBUTAMIDE TABS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL		AMARYL TABS DIABETA TABS GLUCOTROL TABS GLUCOTROL XL TBCR GLYNASE TABS MICRONASE TABS	Use PA Form# 20420 1. Pa required for members ≥65. Glyburide has a greater risk of severe prolonged hypoglycemia in older adults.	Preferred drugs must be tried for at least 3 months at full therapeutic doses and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: All sulfonylureas (except glyburide) will now be non-preferred and require prior authorization if it is currently being used with either ranitidine or cimetidine. DDI: Glimperide will now be non-preferred and require prior authorization if it is currently being used with either fluconazole (except 150mg strength) or fluvoxamine. Amaryl is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with either fluconazole or fluvoxamine.
DIABETIC -ORAL BIGUANIDES	MC/DEL MC/DEL		METFORMIN HCL TABS METFORMIN ER	MC MC MC MC/DEL		GLUCOPHAGE TABS GLUCOPHAGE XR TB24 FORTAMET METFORMIN ER OSMOTIC	Use PA Form# 20420	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
DIABETIC - THIAZOL / BIGUANIDE COMBO				MC/DEL MC/DEL MC MC		ACTOPLUS MET ¹ ACTOPLUS MET XR AVANDARYL ¹ AVANDAMET TABS ¹	Use PA Form# 20420 1. Requires use of Actos, Metformin, or other preferred anti-diabetics.	DDI: Actos, Avandia, or any combination product with Actos or Avandia will now be non-preferred and require prior authorization if it is currently being used with gemfibrozil.
DIABETIC - / THIAZOL	MC/DEL		PIOGLITAZONE HCL ¹	MC/DEL MC		ACTOS TABS ³ AVANDIA TABS ²	1. Pioglitazone HCL is non-preferred as monotherapy. Pioglitazone HCL is preferred if therapeutic	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

						doses of metformin, sulfonylurea or insulin are seen in members drug profile for at least 60 days within the past 18 months. 2. Current users of Avandia who have tried Actos will be able to continue use of Avandia. 3. Dosing limits apply please refer to Dose Consolidation List Use PA Form# 20420	DDI: Actos, Avandia, or any combination product with Actos or Avandia will now be non-preferred and require prior authorization if it is currently being used with gemfibrozil.
DIABETIC - ALPHAGLUCOSIDASE	MC/DEL			MC		PRECOSE TABS Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
DIABETIC - SULFONYLUREA / BIGUANIDE	MC/DEL		GLYBURIDE/METFORMIN	MC MC MC/DEL		GLUCOVANCE TABS ¹ METAGLIP TABS ¹ DUETACT ² Use PA Form# 20420	1. Use individual ingredients. 2. Use Actos with generic glimepiride. Approved for patients failing to achieve good diabetic control with maximal doses of individual components.
DIABETIC - MEGLITINIDES	MC		NATEGLINIDE	MC/DEL MC/DEL		PRANDIN TABS STARLIX TABS Use PA Form# 20420	Preferred drugs from other diabetic sub-categories must be tried and failed due to lack of inadequate diabetic control or intolerable side effects before non-preferred drug will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Prandin is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for current use with both Sporanox and gemfibrozil, due to a significant drug-drug interaction.
GLUCOSE ELEVATING AGENTS							
GLUCOSE ELEVATING AGENTS	MC/DEL MC/DEL	1 2	GLUCAGEN INJ. HYPOKIT ¹ BAQSIMI ^{2,4}	MC MC MC/DEL MC		GLUCAGON DIAGNOSTIC KIT GLUCAGEN DIAGNOSTIC KIT GVOKE ³ ZEGALOGUE ⁵ Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. 1. Dosing limits apply, please see dose consolidation list. 2. For the treatment of patients ≥ 4 years of age. 3. For the treatment of patients ≥ 2 years of age. 4. Baqsimi will require a step through Glucagen. 5. For the treatment of patients ≥ 6 years of age.
THYROID							
THYROID EYE DISEASE				MC		TEPEZZA Use PA Form# 20420	
THYROID HORMONES	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ARMOUR THYROID TABS CYTOMEL TABS ERMEZA ¹ LEVOTHROID TABS LEVOTHYROXINE SODIUM TABS LEVOXYL TABS UNITHROID TABS	MC MC/DEL MC MC/DEL		LEVOTHYROXINE SODIUM SOLR LIOTHYRONINE SYNTHROID TABS THYQUIDITY Use PA Form# 20420	1. Clinical PA is required to confirm diagnosis of dysphagia. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

ANTITHYROID THERAPIES	MC/DEL MC/DEL		METHIMAZOLE TABS PROPYLTHIOURACIL TABS	MC/DEL		TAPAZOLE TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
CUSHING DISEASE AGENTS								
CUSHING DISEASE AGENTS				MC MC		ISTURISA ¹ RECORLEV	1. For the treatment of adult patients with Cushing's disease for whom pituitary surgery is not an option or has not been curative. Use PA Form #20420	Recorlev® is associated with dose-related QT interval prolongation. QT interval prolongation may lead to life-threatening ventricular dysrhythmias such as Torsades de pointes.
OSTEOPOROSIS / BONE AGENTS								
OSTEOPOROSIS	MC/DEL		ALENDRONATE	MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC MC MC/DEL MC MC MC/DEL		ACTONEL TABS AREDIA SOLR BINOSTO BONIVA INJECTION KIT BONIVA TABS ^{2,4} CALCITONIN NS DUAVEE DIDRONEL TABS EVISTA TABS ¹ EVENTITY ² FORTEO FORTICAL FOSAMAX TABS AND PLUS D ³ PROLIA SOHONOS ⁶ STRENSIQ ⁵ TYMLOS XGEVA ZOMETA	Use PA Form# 20420 1. Approval only requires failure of Alendronate. 2. Quantity limits apply, please see dosage consolidation list. 3. Please use Alendronate and Vitamin D. 4. Please use other preferred agents. 5. Obtain baseline ophthalmology exams and renal ultrasounds and then periodically during treatment 6. Clinical PA ffor indication required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Binosto use preferred generic alendronate tablets Evenity® should be limited to 12 monthly doses Sohonos: For the reduction in volume of new heterotopic ossification in adults and pediatric patients aged 8 years and older for females and 10 years and older for males with fibrodysplasia ossificans progressiva (FOP).
FIBROBLAST GROWTH FACTOR 23 INHIBITORS	MC		CRYSVITA ¹				1.Preferred for patients <21 years for the treatment of X-linked hypophosphatemia. Use PA Form #20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
CALCIMIMETIC AGENTS								
CALCIMIMETIC AGENTS				MC MC		PARSABIV SENSIPAR	Use PA Form# 30115	For Sensipar baseline PTH, Ca, and phosphorous levels are required and initial approvals will be limited to 3 months. Subsequent approvals will require additional levels being done to assess changes. Will not approve if baseline Ca is less than 8.4. Parsabiv is for the treatment of secondary hyperparathyroidism (HPT) in adults with chronic kidney disease (CKD) on hemodialysis. Parsabiv® has not been studied in adults with parathyroid carcinoma, primary hyperparathyroidism, or with chronic kidney disease who are not on hemodialysis and is not recommended for use in these populations.
GROWTH HORMONE								
GROWTH HORMONE	MC/DEL MC/DEL MC/DEL		GENOTROPIN ¹ NORDITROPIN SOLN ¹ NUTROPIN AQ ¹	MC MC MC/DEL MC/DEL MC MC MC	8 8 8 8 8 8 8	HUMATROPE SOLR INCRELEX NUTROPIN NGENLA OMNITROPE SAIZEN SOLR SKYTROFA	Use PA Form# 10710 1.Clinical PA is required to establish diagnosis and medical necessity.	See Growth Hormone PA form for criteria. Step-order will still apply unless clinical contraindication supplied. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

				MC/DEL	8	SOGROYA		
				MC/DEL	8	TEV-TROPIN		
ACHONDROPLASIA TREATMENT				MC		VOXZOGO ¹	1. Pediatric patients with achondroplasia who are 5 years of age and older with open epiphyses. Use PA Form# 20420	Voxzogo: To increase linear growth in pediatric patients with achondroplasia who are 5 years of age and older with open epiphyses. This indication is approved under accelerated approval based on an improvement in annualized growth velocity. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s).
SOMATOSTATIC AGENTS				MC/DEL	7	OCTREOTIDE INJ ¹	Use PA Form# 10710	
				MC	8	BYNFEZIA ¹		
				MC	8	MYCAPSSA ¹		
				MC/DEL	8	SANDOSTATIN ¹	1. Non-preferred products must be used in specified step order.	
				MC	8	SOMATULINE ¹		
GROWTH HORMONE ANTAGONISTS								
GH ANTAGONISTS				MC		SOMAVERT	Use PA Form# 10710	Approved for acromegaly patients failing surgery/radiation/drug therapy including bromocriptine and sandostatin.
VASOPRESSIN RECEPTOR ANTAGONIST								
VASOPRESSIN RECEPTOR ANTAGONIST				MC		JYNARQUE ¹	Use PA Form# 20420	Samsca Drug Warning- Avoid use in patients with underlying liver disease, including cirrhosis, because the ability to recover from liver injury may be impaired. Limit duration of therapy to 30 days to minimize the risk of liver injury.
				MC/DEL		SAMSCA	1. Clinical PA required for appropriate diagnosis	DDI: Jynarque- Concomitant use with strong CYP3A inhibitors is contraindicated. Avoid concomitant use of Jynarque® with OATP1B1/B3 and OAT3 substrates (e.g. statins, bosentan, glyburide, nateglinide, repaglinide, methotrexate, furosemide).
URINARY INCONTINENCE								
VASOPRESSINS	MC/DEL		DESMOPRESSIN TABS	MC/DEL	5	DDAVP TABS	1. Products must be used in specified step order.	Approved for central diabetes insipidus and for nocturnal enuresis. For nocturnal enuresis- must be over 6 years old, must fail an adequate trial of alarm training (higher success rate, lower relapse rate) and must periodically attempt weaning (at 6 month intervals).
	MC/DEL		DDAVP SOLN	MC/DEL	6	DESMOPRESSIN SPRAY ¹	Nocturnal enuresis patients will be encouraged to periodically attempt stopping DDAVP.	
				MC	8	DESMOPRESSIN ACETATE SOLN ¹		
				MC/DEL	8	NOCDURNA ¹		
				MC	8	NOCTIVA ¹	2. Patients with a diagnosis of hemophilia or Von Willebrands disease will be exempt from prior authorization.	
				MC/DEL	8	STIMATE SOLN ^{1,2}	Use PA Form# 20420	
ANTISPASMODICS	MC/DEL		DETROL TABS	MC/DEL	8	DARIFENACIN ER TAB	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		DETROL LA CAPS	MC/DEL	8	DITROPAN		
	MC/DEL		OXYBUTYNIN	MC/DEL	8	FLAVOXATE HCL TAB		
				MC/DEL	8	TOLTERODINE		
ANTISPASMODICS - LONG ACTING	MC/DEL		GELNIQUE GEL PACKET	MC	8	DITROPAN XL TBCR	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		MYRBETRIQ	MC/DEL	8	ENABLEX ^{1,2}	1. See Criteria Section.	
	MC/DEL		OXYBUTYNIN ER TABS	MC	8	GEMTESA ²	2. Use a preferred long acting antispasmodic.	1. Vesicare 5mg and Enablex 7.5mg maximum doses if given with drugs known to be significant CYP3A4 inhibitors.(Ketoconazole, Sporanox, Erythromycin, Fluconazole, Nefazodone, Nelfinavir, and Ritonavir)
	MC/DEL		OXYTROL	MC/DEL	8	TOLTERODINE TAB	3. For the treatment of patients ≥ 2 years of age.	DDI: Enablex 15mg and Vesicare 10mg will now be non-preferred and require prior authorization if they are currently being used in combination with any of the following medications: clarithromycin, erythromycin, Ketek, Crixivan, Norvir, ketoconazole, fluconazole (except 150mg strength), Sporanox, nefazodone, or diltiazem.
	MC/DEL		SOLIFENACIN SUCCINATE TAB	MC	8	VESICARE ¹		
	MC/DEL		TOVIAZ	MC	8	VESICARE ³ LS		
	MC/DEL		TROSPIUM					
CHOLINERGIC	MC/DEL		BETHANECHOL	MC/DEL		URECHOLINE	Use PA Form# 20420	
HYPERAMMONIA TREATMENTS	MC		CARGLUMIC ACID TABS	MC		CARBAGLU TABS		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

						Use PA Form# 20420	
UREA CYCLE DISORDER	MC MC		BUPHENYL TABLET PHEBURANE GRANULES	MC MC MC MC/DEL MC/DEL	BUPHENYL POWDER RAVICTI LIQUID OLPRUVA SODIUM PHENYL BUTYRATE POWDER SODIUM PHENYL BUTYRATE TAB	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Olpruva: As adjunctive therapy to standard of care, which includes dietary management, for the chronic management of adult and pediatric patients weighing 20kg or greater and with a body surface area (BSA) of 1.2m2 or greater, with urea cycle disorders (UCDs) involving deficiencies of carbamylphosphate synthetase (CPS), ornithine transcarbamylase (OTC), or argininosuccinic acid synthetase (AS).
METABOLIC MODIFIER							
HERED. TYROSINEMIA				MC	ORFADIN	Use PA Form# 20420	Approved for Type 1 hereditary tyrosinemia patients. Must include laboratory evidence of dx at first PA.
FABRY DISEASE AGENTS				MC MC MC/DEL	ELFABRIO ¹ FABRAZYME ² GALAFOLD ¹	1.Clinical PA to verify appropriate diagnosis. 2.For the treatment of patients 2 years of age and older. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Elfabrio and Galfold: For the treatment of adults with confirmed Fabry disease.
ANTIHYPERTENSIVES / CARDIAC							
CARDIAC GLYCOSIDES	MC/DEL MC/DEL MC/DEL		DIGITEK TABS DIGOXIN LANOXIN			Use PA Form# 20420	
CARDIAC MYOSIN INHIBITORS				MC	CAMZYOS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Camzyos: For the treatment of adults with symptomatic New York Heart Association (NYHA) class II-III obstructive hypertrophic cardiomyopathy (HCM) to improve functional capacity and symptoms. DDI: Concomitant use of Camzyos® with a moderate to strong CYP2C19 inhibitor or a strong CYP3A4 inhibitor is contraindicated.
CARDIAC - SINUS NODE INHIBITORS				MC	CORLANOR	Use PA Form#20420	In patients with stable, symptomatic chronic heart failure with left ventricular ejection fraction ≤35%, who are in sinus rhythm with resting heart rate ≥70 beats per minute (bpm) and
CARDIAC- SOLUBLE GUANYLATE CYCLASE STIMULATORS				MC/DEL	VERQUVO	Use PA Form# 20420	
CARDIAC- SODIUM- GLUCOSE COTRANSPORTER 2 (SGLT2) INHIBITOR				MC	INPEFA ¹	1. To reduce the risk of cardiovascular death, hospitalization for heart failure, and urgent heart failure visit in adults with: Heart failure or Type 2 diabetes mellitus, chronic kidney disease, and other cardiovascular risk factors.	Other Preferred SGLT inhibitors must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

ANTIANGINALS--Isosorbide Di-nitrate/ Mono-Nitrates	MC/DEL MC/DEL		ISOSORBIDE MONONITRATE TABS ISOSORBIDE MONONITRATE ER	MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC	DILATRATE SR CPR ISORDIL TABS ISORDIL TITRADOSE TABS ISOSORBIDE DINITRATE SUBL ISOSORBIDE DINITRATE TABS ISOSORBIDE DINITRATE CR TBCR ISOSORBIDE DINITRATE ER TBCR ISOSORBIDE DINITRATE TD TBCR IMDUR TB24 ISMO TABS MONOKET TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
NITRO - OINTMENT/CAP/CR	MC/DEL MC/DEL MC MC		NITROBID OINT NITROGLYCERIN CPR NITROL OINT NITRO-TIME CPR			Use PA Form# 20420	
NITRO - PATCHES	MC/DEL MC/DEL	1 1	NITROGLYCERIN PT24 ¹ NITRO-DUR PT 24 0.8MG ¹	MC MC/DEL	NITRODISC PT24 NITRO-DUR PT24	1. At least 2 step 1's and step 3 of the preferred products must be used in specified order or PA will be required. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
NITRO - SUBLINGUAL/ SPRAY	MC/DEL		NITROSTAT SUBL	MC/DEL MC MC	NITROQUICK SUBL NITROLINGUAL SOLN NITROLINGUAL TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BETA BLOCKERS - NON SELECTIVE	MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		CARVEDILOL LEVATOL TABS NADOLOL TABS PINDOLOL TABS PROPRANOLOL HCL SOLN ¹ PROPRANOLOL HCL TABS ¹ PROPRANOLOL HCL 60MG TABS PROPRANOLOL LA CAPS RANOLAZINE ER TABS SOTALOL AF SOTALOL HCL TABS TIMOLOL MALEATE TABS	MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC MC MC MC	ASPRUZYO BETAPACE TABS BETAPACE AF TABS COREG CR ³ COREG TABS CORGARD TABS INDERAL TABS HEMANGEOL SOL INDERAL XL CAP INDERAL LA CPR INNOPRAN XL RANEXA	1. Recommend using BID since its effects do not last 24 hours. 2. Please use other strengths in combination to obtain this dose. 3. Dosing limits still apply. Please see dose consolidation list Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Concomitant use of Ranolazine products with strong CYP3A inhibitors, including ketoconazole, itraconazole, clarithromycin, nefazodone, nelfinavir, ritonavir, indinavir, and saquinavir, is contraindicated.
BETA BLOCKERS - CARDIO SELECTIVE	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ACEBUTOLOL HCL CAPS ATENOLOL TABS ¹ BETAXOLOL HCL TABS BISOPROLOL FUMARATE TABS BYSTOLIC METOPROLOL TARTRATE TABS ¹ METOPROLOL ER NEBIVOLOL HCL TAB	MC MC/DEL MC MC/DEL MC/DEL MC/DEL	KERLONE TABS LOPRESSOR TABS SECTRAL CAPS TENORMIN TABS TOPROL XL TB24 ZEBETA TABS	1. Recommend using Atenolol (and metoprolol) BID since its effects do not last 24 hours. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BETA BLOCKERS - ALPHA / BETA	MC/DEL		LABETALOL HCL TABS	MC	TRANDATE TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BETA BLOCKERS & DURECTIC COMBOS	MC/DEL		METOPROLOL-HYDROCHLOROTHIAZIDE TAB	MC/DEL	DUTOPROL	Use PA Form# 20420	
CALCIUM CHANNEL BLOCKERS-- Amlodipines, Bepridil, Diltiazems, Felodipines, Isradipines, Nifedipines, Nisoldipine, and Verapamils	MC/DEL		AMLODIPINE ¹	MC/DEL MC	KATERZIA NORLIQVA	1. Dosing limits apply, please see dose consolidation list.	

			MC/DEL		NORVASC TABS ¹	Use PA Form# 20420	
MC		DILTIA XT CP24	MC/DEL	5	DILACOR XR CP24 ¹	1. Products must be used in specified order or PA will be required. Just write "Diltiazem 24-hour" and the pharmacy will use a preferred long acting diltiazem that does not require PA.	Preferred drugs must be tried and failed (in step-order) due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: All preferred diltiazems will now be non-preferred and require prior authorization if they are currently being used in combination with either Enblex 15mg or Vesicare 10mg. All non-preferred diltiazems require prior authorization, but with any prior authorization request, the member's drug profile will also be monitored for current use with Enblex 15mg or Vesicare 10mg.
MC/DEL		DILTIAZEM HCL ER CP24	MC/DEL	6	TAZTIA ¹		
MC/DEL		DILTIAZEM HCL XR CP24	MC	8	CARDIZEM TABS ¹		
MC/DEL		DILTIAZEM CD 300MG CP24	MC	8	CARDIZEM CD CP24 ¹		
MC/DEL		DILTIAZEM CD 360MG CP24	MC	8	CARDIZEM LA TB24 ¹		
MC		CARTIA XT CP24 ¹	MC	8	CARDIZEM SR CP12 ¹		
MC/DEL		DILTIAZEM CD CP24 ¹	MC/DEL	8	DILTIAZEM HCL TABS ¹		
MC/DEL		DILTIAZEM HCL ER CP24 ¹	MC/DEL	8	DILTIAZEM HCL ER CP12 ¹		
MC/DEL		DILTIAZEM XR CP24 ¹	MC/DEL	8	DILTIAZEM HCL ER CP12 ¹		
MC/DEL		TIAZAC CP24 ¹					
			MC/DEL		PLENDIL TB24	Use PA Form# 20420	Other Preferred calcium channel blockers must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
			MC/DEL		FELODIPINE		
			MC		DYNACIRC CAPS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
			MC		DYNACIRC CR TBCR ¹	1. Established users will be grandfathered	
			MC		CARDENE SR CPCP	Use PA Form# 20420	Other Preferred calcium channel blockers must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
			MC		NICARDIPINE HCL CAPS		
MC/DEL		AFEDITAB CR	MC/DEL		ADALAT CC TBCR ¹	1. Established users of Adalat CC are grandfathered. Use PA Form# 20420	Preferred drug must be tried and failed in step order due to lack of efficacy or intolerable side effects before non-preferred drugs in step order will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MC/DEL		NIFEDIAC CC	MC/DEL		NIFEDIPINE CAPS		
MC/DEL		NIFEDICAL XL TBCR	MC/DEL		PROCARDIA CAPS		
MC/DEL		NIFEDIPINE TBCR	MC/DEL		PROCARDIA XL TBCR		
MC/DEL		NIFEDIPINE ER TBCR					
			MC		SULAR TB24	1. Established users of 10MG and 20MG strengths are grandfathered. Use PA Form# 20420	
			MC		SULAR CR ¹		
MC/DEL	1	VERAPAMIL HCL CR TBCR	MC/DEL		CALAN TABS	Products must be used in specified order or PA will be required. Just write "Verapamil 24-hour" and the pharmacy will use a preferred long acting generic that does not require PA. Use PA Form# 20420	Preferred drugs must be tried and failed (in step-order) due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MC/DEL	1	VERAPAMIL HCL ER TBCR	MC/DEL		CALAN SR TBCR		
MC/DEL	1	VERAPAMIL HCL SR TBCR	MC/DEL		COVERA-HS TBCR		
			MC		ISOPTIN-SR		
			MC/DEL		VERAPAMIL HCL ER CP24		
			MC/DEL		VERAPAMIL HCL SR CP24		
			MC/DEL		VERAPAMIL HCL TABS		
			MC/DEL		VERELAN CP24		
			MC/DEL		VERELAN PM CP24		
ANTIARRHYTHMICS			MC/DEL		CORDARONE	1. Prescription must be written by Cardiologist. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Amiodarone will now be non-preferred and require prior authorization if it is currently being used in combination with either Lovastatin (doses greater than 40mg/day) or Lipitor (doses greater than 20mg/day) or Levofloxacin or Gemifloxacin, or Moxifloxacin, or Ofloxacin. DDI: Multaq will be preferred unless the following medications are seen in the member's drug profile within the last 35 days for brand name medications or 90 days for generic medications: Erythromycin, Amiodarone and other antiarrhythmics, TCA's, Phenothiazine, Ketoconazole, Itraconazole, Voriconazole, Cyclosporine, Telithromycin, Clarithromycin, Nefazodone, Ritonavir.
			MC/DEL		DISOPYRAMIDE		
			MC/DEL		MULTAQ		
			MC/DEL		NORPACE		
			MC/DEL		PACERONE		
			MC/DEL		QUINIDEX		
			MC		TAMBOCOR		
			MC/DEL		TIKOSYN ¹		
			MC/DEL		RYTHMOL SR		
			MC/DEL		RYTHMOL		
ACE INHIBITORS			MC	5	MAVIK TABS	1. Non-preferred products must be used in specified order. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Non-preferred products are subject to step-order requirements unless clinical circumstances warrant exception.
			MC/DEL	5	ACCUPRIL TABS		
			MC/DEL	8	ACEON TABS ¹		
			MC/DEL	8	ALTACE CAPS ¹		
			MC	8	EPANED		
			MC/DEL	8	LOTENSIN TABS ¹		
			MC/DEL	8	MOEXIPRIL HCL ¹		
			MC	8	MONOPRIL HCT TABS ¹		

				MC/DEL	8	PRINIVIL TABS ¹		
				MC	8	QBRELIS		
				MC/DEL	8	UNIVASC ¹		
				MC	8	VASOTEC TABS ¹		
				MC/DEL	8	ZESTRIL TABS ¹		
ANGIOTENSIN RECEPTOR BLOCKER	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		AMLODIPINE-OLMESARTAN TAB ³ IRBESARTAN ¹ LOSARTAN ¹ MICARDIS TABS ³ OLMESARTAN ¹ TELMISARTAN ¹	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC	8 8 8 8 8 8	ATACAND TABS AVAPRO BENICAR TABS COZAAR DIOVAN EDARBI TEVETEN TABS	Use PA Form# 20420 1. Dosing limits apply, please see dose consolidation list. 2. Use preferred active ingredients which are available without PA. 3. Preferred without a PA only if patient on a diabetic therapy or prior ACE therapy.	Per best practices patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy
DIRECT RENIN INHIBITOR				MC/DEL MC/DEL MC/DEL		AMTURNIDE TEKTURNA ¹ TEKAMLO	1. Must show failure of single and combination therapy from all preferred antihypertensive categories. Use PA Form# 20420	
ANTIHYPERTENSIVES - CENTRAL	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL		CLONIDINE HCL TABS GUANFACINE HCL TABS HYDRALAZINE HCL TABS HYLOREL TABS METHYLDOPA TABS MINOXIDIL TABS PRAZOSIN HCL CAPS RESERPINE TABS	MC/DEL MC/DEL MC MC MC/DEL MC MC/DEL		CLONIDINE PATCH CLONIDINE TTS GUANABENZ ACETATE TABS ISMELIN TABS MINIPRESS CAPS NEXICLON TENEX TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ACE INHIBITORS AND CA CHANNEL BLOCKERS				MC/DEL MC MC MC/DEL	8 8 8 9	AMLODIPINE/BENAZEPRIL PRESTALIA ¹ TARKA TBCR LOTREL CAPS	1. Prestalia will only be approved for patients ≥ 18 years of age. Use individual preferred generic medications. Use PA Form# 20420	
ACE AND THIAZIDE COMBO'S	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		BENAZEPRIL HCL/HYDROCHLOR CAPTOPRIL/HYDROCHLOROTHIA ENALAPRIL MALEATE/HCTZ TABS LISINAPRIL-HCTZ TABS LOTENSIN HCT TABS	MC/DEL MC MC/DEL MC/DEL MC MC/DEL		ACCURETIC TABS MONOPRIL HCT TABS PRINZIDE TABS UNIRETIC TABS VASERETIC TABS ZESTORETIC TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BETA BLOCKERS AND DIURETIC COMBO'S	MC/DEL MC/DEL MC/DEL		ATENOLOL/CHLORTHALIDONE BISOPROLOL FUMARATE/HCTZ PROPRANOLOL/HCTZ	MC/DEL MC/DEL MC MC MC/DEL		CORZIDE TABS LOPRESSOR HCT TABS TENORETIC TIMOLIDE 10/25 TABS ZIAC TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ARB'S AND CA CHANNEL BLOCKERS	MC/DEL MC/DEL MC/DEL		AMLODIPINE/VALSARTAN AMLODIPINE/VALSARTAN HCT TRIBENZOR	MC/DEL MC MC/DEL MC/DEL		AZOR BYVALSON EXFORGE EXFORGE HCT	 Use PA Form# 20420	DDI: Byvalson will be non-preferred and require a prior authorization if it is currently being used in combination with drugs known to be significant CYP2D6 inhibitors (e.g. quinidine, propafenone, fluoxetine, paroxetine). Per best practices patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy
ARB'S AND DIURETICS	MC/DEL MC/DEL MC/DEL MC/DEL		BENICAR HCT ¹ LOSARTAN HCT ¹ MICARDIS HCT TABS ¹ VALSARTAN-HCT ¹	MC/DEL MC/DEL MC MC/DEL	7 8 8 8	IRBESARTAN HYDROCHLOROTHIAZIDE ATACAND HCT TABS AVALIDE TABS ¹ DIOVAN HCT TABS ¹	1. Dosing limits apply, please see dose consolidation list.	Per best practices patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy

				MC/DEL MC	8 8	HYZAAR TABS TEVETEN HCT TABS				
ANGIOTENSIN MODULATORS-ARB COMBINATION	MC		ENTRESTO	MC/DEL		EDARBYCLOR			Use PA Form# 20420	
ARB'S AND DIRECT RENIN INHIBITOR COMBINATION				MC/DEL		VALTURNA			Use PA Form# 20420	
DIURETICS	MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		ACETAZOLAMIDE TABS BUMETANIDE CHLOROTHIAZIDE TABS CHLORThALIDONE TABS EDECIN TABS EDECIN TABS HYDROCHLORThIAZIDE INDAPAMIDE TABS METHAZOLAMIDE TABS METHYLOThIAZIDE TABS SPIRONOLACTONE 25MG TABS SPIRONOLACTONE/HYDRO TORSEMIDE TABS TRIAMTERENE/HCTZ ZAROXOLYN TABS	MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL		ALDACTAZIDE TABS ALDACTONE TABS AMILORIDE HCL BUMEX TABS DEMADEX TABS DIAMOX DIURIL DYAZIDE CAPS CAROSPIR ENDURON TABS INSPRA KERENDIA KEVEYIS LASIX TABS MAXZIDE MICROZIDE CAPS MIDAMOR TABS NAQUA TABS SPIRONOLACTONE 50MG ¹		1. Multiples of Spironolactone 25 mg are cheaper than 50 mg strength. Inspira will be approved for severe breast tenderness and male gynecomastia. DDI: The concomitant use of Keveysis® with high dose aspirin is contraindicated.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
CCB / LIPID				MC/DEL		CADUET			Use PA Form# 20420	
NEUROGENIC ORTHOSTATIC HYPOTENSION										
NEUROGENIC ORTHOSTATIC HYPOTENSION				MC		NORTHERA			Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
LIPID DRUGS										
CHOLESTEROL - BILE SEQUESTRANTS	MC/DEL MC/DEL		CHOLESTYRAMINE COLESTIPOL HCl	MC/DEL MC/DEL MC MC/DEL		COLESTID PREVALITE QUESTRAN WELCHOL TABS			Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
CHOLESTEROL - FIBRIC ACID DERIVATIVES	MC/DEL MC/DEL MC/DEL		FENOFIBRATE TAB GEMFIBROZIL TABS NIACIN ER	MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC		ANTARA LOPID FENOFIBRATE 120mg TAB FENOFIBRATE CAP FIBRICOR LIPOFEN LOFIBRA NIASpan ER TRICOR TRIGLIDE TRILIPIX		Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Fenofibrate is preferred but will require a prior authorization requests if used concurrent with Warfarin. DDI: Gemfibrozil will now be non-preferred and require prior authorization if it is currently being used with any of the following medications: Prandin, Actos, Avandia, any Avandia/Actos combination product, any HMG-COA Reductase Inhibitors (statins), or Warfarin.	
CHOLESTEROL - HMG COA + ABSORB INHIBITORS MORE POTENT DRUGS/COMBINATIONS	MC/DEL MC/DEL MC MC/DEL		ATORVASTATIN EZETIM/SIMVA TAB ROSUVASTATIN SIMVASTATIN ¹	MC MC/DEL MC/DEL MC/DEL MC		ATORVALIQ CRESTOR EZALLOR SPRINKLES ³ LIPITOR LIPTRUZET		1. Dosing limits apply, please see dosage consolidation list. 2. Current users grandfathered.	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Lipitor (doses greater than 20mg/day) will now be non-preferred and require prior authorization if they are currently being used in combination cyclosporine.

				MC/DEL		ZOCOR	3. For the treatment of patients ≥ 18 years of age.	DDI: Lipitor (doses greater than 20mg/day) will now be non-preferred and require prior authorization if it is currently being used in combination with Amiodarone.
				MC/DEL MC		SIMVASTATIN 80MG ^{1,2} VYTORIN	Use PA Form# 20420	DDI: All preferred statins will now be non-preferred and require prior authorization if it is currently being used in combination with Gemfibrozil.
CHOLESTEROL - HMG COA + ABSORB INHIBITORS LESS POTENT DRUGS/COMBINATIONS	MC/DEL MC/DEL MC/DEL		EZETIMIBE TABS LOVASTATIN TABS ² PRAVASTATIN ²	MC MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL	8 8 8 8 8 8 8 8 8 8	ALTOPREV TB24 FLUVASTATIN TAB ER LESCOL XL TB24 LIVALO MEVACOR TABS NEXLETOL NEXLIZET PRAVACHOL TABS PRAVIGARD	2. Dosing limits apply, please see dosage consolidation list	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Zetia will be approved for patients unable to tolerate all other therapies or unable to achieve cholesterol goal with maximally tolerated dose of most potent statins. DDI: Lescol will now be non-preferred and require prior authorization if it is currently being used in combination with diclofenac. DDI: Lovastatin (doses greater than 40mg/day) will now be non-preferred and require prior authorization if it is currently being used in combination with Amiodarone. DDI: Lovastatin (doses greater than 20mg per day) will now be non-preferred and require prior authorization if it is currently being used in combination cyclosporine. DDI: All preferred statins will now be non-preferred and require prior authorization if it is currently being used in combination with Gemfibrozil.
CHOLESTEROL - HMG COA + ABSORB INHIBITORS STATIN/ NIACIN COMBO	MC		SIMCOR	MC		ADVICOR TBCR	Use PA Form# 20420	
FAMILIAL HYPERCHOLESTEROLEMIA	MC MC		PRALUENT (LABLER 72733) PEN ^{1,2,3,5,6} REPATHA ^{1,2,3}	MC MC MC MC		EVKEEZA ^{1,4} JUXTAPID KYNAMRO ¹ LEQVIO	1. Clinical PA required for appropriate diagnosis 2. Quantity limits apply 3. Documented adherence to lipid lowering medications and abstinence from tobacco for previous 90 days 4. For the treatment of patients ≥ 12 years of age. 5. Approval of Praluent NDC's with labeler code 00024 will be considered only if labeler code 72733 NDC's are on a long-term backorder and unavailable from the manufacturer.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists Juxtapid is contraindicated with strong CYP3A4 inhibitors. Juxtapid dosage should not exceed 30mg daily when it is used concomitantly with weak CYP3A4 inhibitors. Kynamro requires an appropriate lab testing prior to starting (ALT<AST), Alkaline phosphatase and total billrubin, monthly liver-related tests for the first year, then every three months. Repatha and Praluent Criteria for approval: The patients's age is FDA approved for the given indication AND • Concurrent use with statin therapy AND • Documented adherence to prescribed lipid lowering medications for the previous 90 days AND • Recommended or prescribed by a lipidologist or cardiologist AND • Inability to reach goal LDL-C despite a trial of 2 or more maximum tolerated dose of statins (one of which must be atorvastatin or rosuvastatin) and ezetimibe 10mg daily Additional criteria for the diagnosis of heterozygous familial hypercholesterolemia (HeFH): (both are required): Total cholesterol > 290 mg/dL OR LDL-C > 190 mg/dL AND one of the following • Presence of tendon xanthomas OR • In 1st or 2nd degree relative-documented tendon xanthomas, MI at age ≤ 60 years or TC > 290 mg/dL . Additional criteria for the diagnosis of clinical atherosclerotic cardiovascular disease: History of MI, angina, coronary or other arterial revascularization, stroke, TIA, or PVD of atherosclerotic origin. Additional criteria for the diagnosis of homozygous familial hypercholesterolemia (Repatha only): Total cholesterol levels > 290mg/dL or LDL-C > 190mg/dL (adults) OR Total cholesterol levels > 260mg/dL or LDL-C > 155mg/dL (children < 16 years) and TG within reference range OR Confirmation of diagnosis by gene testing.
PULMONARY ANTI-HYPERTENSIVES								
PULMONARY ANTI-HYPERTENSIVES	MC MC/DEL MC/DEL MC		EPOPROSTENOL INJ ^{3,6} SILDENAFIL TADALAFIL VENTAVIS ³	MC/DEL MC MC/DEL MC MC MC MC MC/DEL MC MC MC MC		ADEMPAS ^{1,3} ADCIRCA ⁴ ALYQ TAB FLOLAN ³ LIQREV OPSUMIT ^{1,2} ORENITRAM REMODULIN ³ REVATIO ⁴ TADLIQ ⁴ TYVASO UPTRAVI VELVETRI ³	1. Requires previous trials/failure of multiple preferred medications. 2. Dosing limits apply, please see the dose consolidation list. 3. Require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and NYHA functional class 3 or 4. 4. Require WHO Group 1 diagnosis of primary PAH	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Sildenafil will be preferred with clinical PA for treatment of pulmonary arterial hypotension (WHO Group 1) in adults to improve exercise ability and delay clinical worsening. Avoid concomitant use of Sildenafil with moderate or strong Cyp3A inhibitors DDI: Uptravi will require a prior authorization if it is currently being used in combination with strong inhibitors of CYP2C8 (gemfibrozil) DDI: Opsumit will require a prior authorization if it is currently being used in combination with drugs known to be significant CYP3A inhibitors (ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, nelfinavir, ritonavir, atazanavir, saquinavir and telithromycin). DDI: Adempas will require a prior authorization if it is currently being used in combination with drugs known to be PDE inhibitors should be avoided (including dipyridamole, addcira and tadalafil) with adempas

							Primary Pulmonary Hypertension) and NYHA functional class 2 or 3.	Liqrev: treatment of pulmonary arterial hypertension (WHO Group 1) in adults to improve exercise ability and delay clinical worsening. Avoid concomitant use of Liqrev with moderate or strong CYP3A inhibitors.
							Use PA Form# 20420	
ERA / ENDOTHELIN RECEPTOR ANTAGONIST	MC MC		LETAIRIS ^{1,2} TRACLEER				1. Providers must be registered with LEAP Prescribing program, a restricted distribution program. 2. Clinical PA is required to establish diagnosis and medical necessity.	Tracleer approvals will require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and NYHA functional class 2 thru 4. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with Tracleer. Letairis approvals will require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and functional class 2 or 3 symptoms.
							Use PA Form# 20420	
IMPOTENCE AGENTS								
IMPOTENCE AGENTS							As of January 1, 2006, per CMS (federal govt.), impotence agents are no longer covered.	As of January 1, 2006, per CMS (federal govt.), impotence agents are no longer covered.
ANTI-EMETOGENICS								
ANTIEMETIC - ANTICHOLINERGIC / DOPAMINERGIC	MC MC/DEL MC MC/DEL MC		BONJESTA MECLIZINE HCL TABS PROMETHAZINE SUPP PROMETHAZINE TRANSDERM-SCOP PT72	MC MC MC MC MC		ANTIVERT TABS BARHEMSYS PHENERGAN SOLN PROMETHAZINE 50MG SUPP PROMETHEGAN SUPP TORECAN TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Concomitant use of MAOIs and Bonjesta® is contraindicated.
ANTIEMETIC - 5-HT3 RECEPTOR ANTAGONISTS/ SUBSTANCE P NEUROKININ	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		DICLEGIS DRONABINOL CAPS GRANISETRON TAB ONDANSETRON TAB ONDANSETRON ODT TBDP ONDANSETRON SOL	MC MC MC MC MC MC MC/DEL MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL MC	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	AKYNZEO ¹ APREPITANT ALOXI ANZEMET TABS APONVIE ⁴ CESAMET ¹ CINVANTI ⁴ EMEND ² KYTRIL MARINOL CAPS SANCUSO SUSTOL SYNDROS TRIMETHOBENZAMIDE CAP VARUBI ZOFTRAN ODT TBDP ³ ZOFTRAN TABS ³ ZOFTRAN INJ ³ ZUPLLENZ	1. Approvals will require diagnosis of chemo-induced nausea/vomiting and failed trials of all preferred anti-emetics, including 5-HT3 class (Ondansetron) and Marinol. 2. Clinical PA is required for members on highly emetic anti-neoplastic agents. 3. Dosing limits apply, please see Dosage Consolidation List 4. Clinical PA required for appropriate diagnosis	Preferred drugs and step therapy must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. * Ondansetron limits still apply as listed on the Ondansetron PA form for covered indications including chemotherapy, radiotherapy, post operative nausea & vomiting and hyperemesis gravidarum. Other medical indications will be approved or denied on a case by case basis. Hyperemesis and other medical indications approved are still subject to failure of multiple preferred antiemesis drugs. Akynzeo- Concomitant use should be avoided in patients who are chronically using a strong CYP3A inducer such as rifampin. Varubi – Available to the few who are unable to tolerate or who have failed on preferred medications Aponvie is for the prevention of postoperative nausea and vomiting (PONV) in adults.
							Use PA Form# 20420	
NON-SEDATING ANTIHISTAMINES / DECONGESTANTS								
ANTIHISTIMINES - NON-SEDATING	MC MC/DEL MC/DEL MC		ALAVERT TABS CETIRIZINE TABS LORATADINE TAVIST ND (OTC)	MC MC MC/DEL MC/DEL MC/DEL	5 5 5 5 5	CLARINEX TABS ^{1,5} CLARINEX SYR ^{1,2} FEXOFENADINE ¹ ZYRTEC ¹ ZYRTEC SYR ^{1,2}	1. Must fail preferred drugs, OTC loratidine and cetirizine before moving to non-preferred step order drugs.	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. No combination product with decongestant will be approved since pseudoephedrine available without PA.

				MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL	8 8 8 8 8 9	ALLEGRA ³ CLARITIN ³ DES LorATADIN LORATADINE ODT ⁴ LEVOCETIRIZINE ⁴ XYZAL ³	2. Clarinex and Zyrtec syrps <6 yr w/o PA. 3. Must fail all step 5 drugs (Clarinex, Fexofenadine and Zyrtec) before moving to next step product. 4. All OTC versions of loratadine ODT are now non-preferred. 5. Pa's for Clarinex RediTabs will only be approved if between the ages of 6-11 years old. Use PA Form# 20530	Pseudoephedrine is available with prescription.	
ANTIHISTIMINES - OTHER	MC/DEL MC/DEL MC/DEL		CLEMASTINE CHLORPHENIRAMINE DIPHENHYDRAMINE				Use PA Form# 20530		
ALLERGY / ASTHMA THERAPIES									
ANAPHYLACTIC DEVICES	MC/DEL MC/DEL MC/DEL		EPINEPHRINE EPIPEN EPIPEN JR	MC MC/DEL		TWINJECT SYMJEPI		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Use PA Form# 20420	
ALLERGEN IMMUNOTHERAPY				MC MC MC MC MC		ODACTRA ORALAIR ¹ PALFORZIA RAGWITEK GRASTEK	Use PA Form# 20420 1. See criteria section	Prescriber must provide the testing to show that the patient is allergic to the components in the prescribed therapy and must provide a clinically valid rationale why single agent sublingual therapy is being chosen over subcutaneous therapy Palforzia® is approved for use in patients with a confirmed diagnosis of peanut allergy. Initial dose escalation may be administered to patients aged 4 through 17 years. Up-dosing and maintenance may be continued in patients 4 years of age and older. Odactra® is approved for use in persons 12 through 65 years of age. Note that Odactra® is not indicated for the immediate relief of allergic symptoms. Treatment must start 12 weeks before expected onset of pollen season and only after confirmed by positive skin test or in vitro testing for pollen-specific IgE antibodies for any of the 5 grass species contained in Oralair Oralair: Patient age ≥10 years and ≤65 years Have an auto-injectable epinephrine on-hand	
ANTI-ASTHMATIC - ANTICHOLINERGICS - INHALER	MC MC/DEL MC/DEL		INCRUSE ELLIPTA ³ SPIRIVA HANDIHALER ^{1,2} SPIRIVA RESPIMAT	MC/DEL MC MC/DEL		FLUTICASONE-SALMETEROL LONHALA MAGNAIR TUDORZA	Use PA Form# 20420 1. Quantity limit of 1 inhalation daily (1 capsule for combination inhaler) 2. We ask physicians to write "asthma" on the prescription whenever Spiriva is primarily being used for that condition. 3. Quantity limit of 1 inhalation daily	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	

ANTIASTHMATIC - PHOSPHODIESTERASE 4 INHIBITORS				MC/DEL		DALIRESP	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - ANTICHOLINERGICS - NEBULIZER	MC/DEL		IPRATROPIUM BROMIDE SOLN	MC MC/DEL		ATROVENT SOLN YUPELRI	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - ANTIINFLAMMATORY AGENTS	MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL		CROMOLYN SODIUM NEBU DUPIXENT ^{2,4} FASENRA ² FASENRA ² AUTO INJCT NUCALA ² SYRINGE 40MG XOLAIR ¹	MC MC		CINQAIR ³ TEZSPIRE ⁵	1. Need max inhaled steroids and written by pulmonary or allergy specialist. Must have elevated IgE and ≥ to age 6. 2. For patients with severe asthma aged 12 years or older and eosinophilia. 3. For patients ≥ 18 years of age with eosinophilia. 4. Clinical PA required. 5. For adult and pediatric patients aged 12 years and older with severe asthma. Use PA Form# 20420	All will require suboptimal response to maximal doses of inhaled steroid as evidenced by asthmatic ER/Hospital admissions and Allergy/Pulmonary specialist management. Dupixent limited to patient with asthma not controlled on high dose ICS-LABA who have eosinophil greater than or equal to 150 cells or the patient is depend on an oral corticosteroid Fasenra, Nucala and Cinqair are not indicated for treatment of other eosinophilic conditions and are not indicated for the relief of acute bronchospasm or status asthmaticus.
ANTIASTHMATIC - NASAL STEROIDS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC		BUDESONIDE SPRAY FLUTICASON SPR ³ OLOPATADINE SPRAY OMNARIS SPR ³ TRIAMCINOLONE NS QNASL	MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC MC MC/DEL MC MC/DEL	5 8 8 8 8 8 8 8 8 8 8 8	BECONASE AQ INHA ^{1,3} DYMISTA FLONASE SUSP ^{2,3} FLUNISOLIDE SOLN ^{1,3} NASONEX SUSP RHINOCORT AERO ^{2,3} RHINOCORT AQUA SUSP ^{2,3} RYALTRIS ⁴ TRI-NASAL SOLN ^{2,3} VANCENASE POCKETHALER AERS ^{2,3} VERAMYST ^{2,3} XHANCE ² ZETONNA ³	Use PA Form# 20420 1. All preferred drugs must be tried before moving to non preferred steps. 2. All step 5 medications need to be tried before moving to step 8's. 3. Dosing limits apply to whole category, please see dosage consolidation list. 4. Use of individual ingredients or other preferred agents.	Preferred drugs and step therapy must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Xhance will be considered for the treatment of nasal polyps in patients 18 years of age or older. The patient has had a documented side effect, allergy, or treatment failure of two preferred nasal glucocorticoids, one of which must be fluticasone.
ANTIASTHMATIC - NASAL MISC.	MC/DEL MC/DEL MC		AZELASTINE CROMOLYN NASAL 4% IPRATROPIUM NASAL SOL ¹	MC/DEL MC/DEL	8 8	ASTEPRO ² PATANASE	Use PA Form# 20420 1. Ipratropium will be approved if submitted with documentation supporting use of CPAP machine. 2. Utilize Multiple preferred, as well as step therapy Azelastine.	Approved if patient fails on non-sedating antihistamines and steroid nasal sprays. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - BETA - ADRENERGICS	MC/DEL MC/DEL MC MC/DEL MC MC/DEL		ALBUTEROL NEB METAPROTERENOL PROAIR RESPICLICK PROVENTIL HFA SEREVENT TERBUTALINE SULFATE TABS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL		ACCUNEB NEBU ALBUTEROL HFA BRETHINE LEVALBUTEROL TARTRATE PROAIR DIGHALER ⁴ STRIVERDI	1. Xopenex users w/ prior asthma hospitalization due to albuterol nebulizer failure will be grandfathered. 2. Quantity Limit: 12	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

	MC/DEL MC	ALBUTEROL 0.63mg/3ml VENTOLIN HFA AERS	MC MC MC MC	VOLMAX TBCR VOSPIRE ER TB12 XOPENEX HFA ³ XOPENEX NEBU ^{1,2}	cc/day. 3. Dosing limits apply, please see dosage consolidation list. 4. For the treatment of patients ≥ 4 years of age. Use PA Form# 20420	
ANTIASTHMATIC - ADRENERGIC COMBINATIONS	MC MC MC MC MC/DEL MC/DEL	ADVAIR DISKUS ¹ ADVAIR HFA ¹ AIRDUO RESPICLICK ² BREQ ELLIPTA ¹ DULERA SYMBICORT	MC MC/DEL MC/DEL MC	AIRDUO DIGIHALER ² AIRSUPRA BREZTRI AEROSPHERE TRELLEGY ELLIPTA ¹	1. Dosing limits apply, please see dosage consolidation list. 2. For patients ≥ 12 years and older. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. AirDuo® Respiclick be non-preferred and require prior authorization and be available to those who are unable to tolerate or who have failed on preferred medications DDI: Avoid concomitant use of strong CYP3A4 inhibitors (e.g. ritonavir, atazanavir, clarithromycin, indinavir, itraconazole, nefazodone, nelfinavir, saquinavir, ketoconazole, telithromycin) with AirDuo® Respiclick is not recommended due to increased systemic corticosteroid and increased cardiovascular adverse effects
ANTIASTHMATIC - ADRENERGIC ANTICHOLINERGIC	MC/DEL MC MC/DEL MC/DEL	ALBUTEROL/IPRATROPIUM NEB. SOLN ANORO ELLIPTA COMBIVENT RESPIMAT STIOLTO	MC/DEL MC/DEL MC/DEL	BEVESPI AEROSPHERE ^{2,3} DUAKLIR PRESSAIR DUONEB SOLN ¹	1. Please use preferred individual ingredients Albuterol and Ipratropium. 2. Dosing limits apply, please see dosing consolidation list. 3. The safety and efficacy of use in children under the age of 18 years have not been established. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Duoneb components are available separately without PA. DDI: Avoid concomitant use of Bevespi with other anticholinergic-containing drugs, due to an increased risk of anticholinergic adverse events. Bevespi® should be used with extreme caution in patients being treated with MAO inhibitors, TCAs, or other drugs known to prolong the QTc interval. Bevespi should be used with extreme caution in patients being treated with MAO inhibitors, TCAs, or other drugs known to prolong the QTc interval.
ANTIASTHMATIC - XANTHINES	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL	AMINOPHYLLINE TABS THEOCHRON TB12 THEOLAIR-SR TB12 THEOPHYLLINE CR TB12 THEOPHYLLINE ELIX THEOPHYLLINE SOLN THEOPHYLLINE ER CP12 THEOPHYLLINE ER TB12	MC/DEL MC MC/DEL	THEO-24 CP24 THEOLAIR TABS UNIPHYL TBCR	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - STEROID INHALANTS	MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC	ARNUITY ELLIPTA ASMANEX TWISTHALER ^{3,4} ASMANEX HFA ⁵ BUDESONIDE NEB 0.25MG & 0.5MG ¹ FLOVENT DISKUS ³ PULMICORT FLEXHALER ³ QVAR AERS ³	MC MC/DEL MC MC/DEL MC/DEL MC	8 AEROSPAN 8 ALVESCO ³ 8 ARMONAIR DIGIHALER 8 BUDESONIDE NEB 1MG 8 PULMICORT SUSP 8 FLOVENT HFA ³	1. Budesonide Neb 0.25mg & 0.5mg will be preferred for members under the age of 8 years old. PA will be required for members 8 years of age and older, please consider other preferred options. 2. All preferreds must be	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

							<p>tried before moving to non preferred steps.</p> <p>3. Dosing limits apply, please see dosage consolidation list.</p> <p>4. Asmanex 110mcg will be limited to member between the ages of 4-11years old.</p> <p>5. Asmanex HFA will be preferred for members under the age of 6 years old. PA will be required for members 6 years of age and older, please consider other preferred options.</p> <p>Use PA Form# 20420</p>	
ANTIASTHMATIC - 5-Lipoxygenase Inhibitors				MC		ZYFLO CR TABS		Other Preferred asthma controller drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - LEUKOTRIENE RECEPTOR ANTAGONISTS	MC/DEL MC/DEL MC/DEL		MONTELUKAST GRANULE ¹ MONTELUKAST SODIUM TAB MONTELUKAST SODIUM CHEW TAB	MC/DEL MC/DEL MC/DEL	8 8 8	ACCOLATE TABS SINGULAIR ² SINGULAIR GRANULES	Use PA Form# 20420 1.Montelukast Granules will only be approved if between ages of 6months-24 months. 2.Singulair Chewables 4mg from 2years-5years and Singulair Chewables 5mgs from 6years-14years old.	
ANTIASTHMATIC - ALPHA-PROTEINASE INHIBITOR				MC MC/DEL MC MC	8 8 8 8	ARALAST ZEMAIRA GLASSIA PROLASTIN SUSR	Use PA Form# 20420	Prolastin and Azemaira will be approved for members with A1AT deficiency and clinically demonstrable panacinar emphysema.
ANTIASTHMATIC - HYDRO-LYTIC ENZYMES				MC/DEL		PULMOZYME SOLN	Use PA Form# 20420	Will be approved for cystic fibrosis patients.
ANTIASTHMATIC - MUCOLYTICS	MC/DEL		ACETYLCYSTEINE ¹	MC		MUCOMYST	1. Acetylcysteine is covered with diagnosis of CF. Use PA Form# 20420	
ANTIASTHMATIC-CFTR POTENTIATOR AND COMBINATIONS				MC MC MC MC MC/DEL		BRONCHITOL ¹ ORKAMBI KALYDECO SYMDEKO TRIKAFTA	1. For the treatment of patients ≥18 years of age with CF. Use PA Form# 20420	<p>Kalydeco will be considered for patients with cystic fibrosis (CF) aged 1 month and older who have at least one mutation in the CFTR gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use.</p> <p>Symdeko will be considered for patients with cystic fibrosis (CF) aged 6 years and older who are homozygous for the <i>F508del</i> mutation or who have at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor based on in vitro data and/or clinical evidence. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use.</p> <p>Bronchitol will be considered as add-on maintenance therapy to improve pulmonary function in adult patients 18 years and older with cystic fibrosis (CF). Use Bronchitol® only for adults</p>

who have passed the Bronchitol® Tolerance Test (BTT). (see Recommended Dosage section for further information)

Trikafta will be considered for the treatment of cystic fibrosis (CF) in patients aged 2 years and older who have at least one F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene or mutation in the CFTE gene that is responsive based on in vitro data. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to confirm the presence of at least one F508del mutation or a mutation that is responsive based on in vitro data.

Orkambi will be considered for patients with cystic fibrosis (CF) aged 1 year and older who are homozygous for the F508del mutation in the CFTR gene. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of the F508del mutation on both alleles of the CFTR gene. The efficacy and safety of Orkambi have not been established in patients with CF other than those homozygous for the F508del mutation.

[Use PA Form# 20420](#)

IDIOPATHIC PULMONARY FIBROSIS	MC/DEL		OFEV ¹	MC MC	ESBRIET ¹ PIRFENIDONE	1. Diagnosis required Use PA Form# 20420	Ofev- Avoid concomitant use with P-gp and CYP4A inducers (e.g. carbamazepine, phenytoin, and St. John's wort) Esbriet- The concomitant use with strong CYP1A2 inhibitors (e.g. fluvoxamine, enoxacin) is not recommended
COUGH/COLD							
COUGH/COLD	MC/DEL MC/DEL MC/DEL MC MC		DEXTROMETHORPHAN CAPS ¹ DEXTRO-GUAIF SYRP ¹ GUAIFENESIN SYRP ¹ PSEUDOEPHEDRINE ¹ ROBITUSSIN DM SYRP ¹ ROBITUSSIN SUGAR FREE SYRP ¹			1. All of cough cold preparations are not covered except these preferred products. Use PA Form# 20420	All non-preferred products are not covered as permitted by Federal Medicaid regulations and MaineCare Policy.
DIGESTIVE AIDS / ASSORTED GI							
GI - ANTIPERISTALTIC AGENTS	MC/DEL MC/DEL MC/DEL MC		DIPHENOXYLATE DIPHENOXYLATE/ATROPINE LOPERAMIDE HCL CAPS/LIQ OPIUM TINCTURE TINC PAREGORIC TINC	MC/DEL MC MC	LOFENE TABS LONOX TABS MOTOFEN TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.
GI - ANTI-DIARRHEAL/ ANTACID - MISC.	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ATROPINE SULFATE SOLN BISMATROL BISMUTH SUBSALICYLATE CALCIUM CARBONATE (ANTACID) CHEW DICYCLOMINE HCL GLYCOPYRROLATE TABS HYOSCYAMINE CAPS & TABS HYOSCYAMINE SULFATE KAOPECTATE MAGNESIUM OXIDE TABS MAG-OX 400 TABS PAMINE TABS PROPANTHELINE BROMIDE TABS SODIUM BICARBONATE TABS TUMS	MC/DEL MC/DEL MC MC MC MC MC/DEL MC MC MC MC MC MC MC MC MC MC MC MC MC	BELLADONNA ALKALOIDS & OP BENTYL TABS BENTYL SYRP CUVPOSA DARTISLA ODT ² ED-SPAZ MYTESI ¹ GLYCOPYRROLATE INJ LEVSIN TABS LEVSIN/SL SUBL NULEV TBP OSCIMIN ROBINUL INJ ROBINUL TABS	Use PA Form# 20420 1. Dosing limits apply please refer to Dose Consolidation List 2. It is not indicated as monotherapy for treatment of peptic ulcer because effectiveness in peptic ulcer healing has not been established.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. Preferred products that used to require diag codes still require diag codes unless indicated otherwise. Mytesi requires a diagnosis of non-infectious diarrhea in patients with HIV/AIDS on anti-retroviral therapy, prior trials of preferred, more cost effective anti-diarrheals.
GI- BILE ACID				MC	CHOLBAM	Use PA Form# 20420	Indication of bile acid synthesis disorders due to single enzyme defects (SEDs) AND for adjunctive treatment of peroxisomal disorders (PDs)
GI - H2-ANTAGONISTS	MC MC/DEL MC/DEL		ACID REDUCER TABS CIMETIDINE FAMOTIDINE	MC MC MC/DEL MC/DEL MC	AXID CAPS AXID AR TABS NIZATIDINE CAPS PEPCID PEPCID AC	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Cimetidine will now be non-preferred and require prior authorization if it is currently being used with any sulfonylurea (except for glyburide). DDI: Cimetidine will require prior authorization if being used in combination with Plavix.

GI- IBAT INHIBITORS				MC MC		BYLVAY ^{1,2} LIVMARLI ^{1,2}	Use PA Form# 20420 1. For the treatment of patients ≥ 3months of age 2. Clinical PA required for appropriate diagnosis	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.
GI - PROTON PUMP INHIBITOR	MC/DEL MC/DEL MC/DEL		OMEPRAZOLE CAPS ² PANTOPRAZOLE ² LANSOPRAZOLE CAPS ²	MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	6 6 7 7 8 8 8 8 8 8 8 8 8 8	NEXIUM CPDR ³ NEXIUM SUS ⁵ PRILOSEC OTC ³ ACIPHEX TBEC ³ DEXILANT (KAPIDEX) ² KONVOME ² OMEPRAZOLE-SODIUM BICARBONATE CAPS OMEPRAZOLE MAGNESIUM PREVACID CPDR ³ PREVACID SOLUTABS ^{1,4} PRILOSEC CPDR PROTONIX INJ PROTONIX ²	1. Prevacid Solutabs available without PA for children less than 9 years old. 2. Dosing limits apply, please see dosage consolidation list. 3. All preferreds and step therapy must be tried and failed 4. Payment for Prevacid SoluTabs for patients 9 and older will be considered for those patients who cannot tolerate a preferred solid oral dosage form. 5. Nexium sus available without PA if member is < 12 yrs of age and ≤ 1 pack per day Use PA Form# 20720	All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please refer to the PPI PA form for additional criteria on Non-Preferred PPIs DDI: Omeprazole will require prior authorization if being used in combination with Plavix. DDI: Lansoprazole will require prior authorization if being used in combination with Plavix. DDI: Prevacid, Omeprazole and pantoprazole will now be non-preferred and require prior authorization if they are currently being used in combination with any of the following medications: Ampicillin, B-12, Fe salts, Griseofulvin, Sporanox, Ketoconazole, Reyataz, or Vantin. DDI: All non-preferred PPIs require prior authorization, but with any prior authorization request, the member's drug profile will also be monitored for current use with ampicillin, B-12, Fe salts, griseofulvin, itraconazole, ketoconazole, Reyataz or Vantin due to a significant drug-drug interaction.
GI - ULCER ANTI-INFECTIVE	MC MC		PYLERA TALICIA				Use PA Form# 20420	
GI - PROSTAGLANDINS	MC		MISOPROSTOL TABS	MC/DEL		CYTOTEC TABS	Use PA Form# 20420	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drug will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
GI - DIGESTIVE ENZYMES	MC/DEL MC		CREON ¹ ZENPEP ¹	MC/DEL MC/DEL MC/DEL		PERTZYE ULTRESA VIOKACE	Use PA Form# 20420 1. Clinical PA is required to establish CF diagnosis and medical necessity. In all cases except cystic fibrosis patients, objective evidence of pancreatic insufficiency (fat malabsorption test etc...) must be supplied.	Non -Preferred drugs must be tried and failed in step-order due to lack of efficacy or intolerable side effects before other non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
GI - ANTI - FLATULENTS / GI STIMULANTS	MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		AMITIZA CALULOSE SYRP CONSTULOSE SYRP ENULOSE SYRP GASTROCROM CONC GENERLAC SYRP LACTULOSE SYRP METOCLOPRAMIDE HCL	MC MC/DEL MC MC/DEL		CEPHULAC SYRP INFANTS GAS RELIEF SUSP GIMOTI SPRAY REGLAN TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.

GI - INFLAMMATORY BOWEL AGENTS	<p>MC MC/DEL MC MC MC/DEL MC/DEL</p>		<p>APRISO BALSALAZIDE MESALAMINE ENMA KIT PENTASA SULFAZINE EC TBEC SULFASALAZINE TABS</p>	<p>MC/DEL MC/DEL MC MC MC/DEL MC MC/DEL MC/DEL MC MC MC/DEL MC MC MC</p>	<p>ASACOL 800MG HD AZULFIDINE EN-TABS TBEC AZULFIDINE TABS COLAZAL CAPS DELZICOL DIPENTUM CAPS GIAZO LIALDA TABS¹ MESALAMINE TAB ROWASA ENEM SFROWASA UCERIS RECTAL FOAM² UCERIS TABS²</p>	<p>Use PA Form# 20420 1. Current users grandfathered. 2. Diagnosis required</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Giazio is only indicated for males, as the safety.efficacy for use in females has not been established.Prior trials of preferred products. Uceris Rectal Foam or Tab- Concomitant use with CYP3A inhibitors (e.g. ketoconazole, itraconazole, ritonavir, indinavir, saquinavir, erythromycin, cyclosporine, and grapefruit juice) should be avoided. Verify prior trials and failures or intolerance of preferred treatments</p>
GI - IRRITABLE BOWEL SYNDROME AGENTS	MC/DEL		LOTRONEX TABS	MC	VIBERZI	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
GI- SHORT BOWL SYNDROME				MC	GATTEX	Use PA Form #20420	Gattex requires a diagnosis of adult SBS who are dependent on parenteral support. Appropriate colonoscopy and lab assessments 6months prior to starting
MISCELLANEOUS GI							
GI - MISC.	<p>MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC MC MC/DEL</p>		<p>BISAC-EVAC SUPP BISACODYL BISCOLAX SUPP CINOBAC CAPS CITRATE OF MAGNESIA SOLN CITRUCEL CLENPIQ SOL COLYTE DIOCTO SYRP DOCUSATE CALCIUM CAPS DOCUSATE SODIUM FIBER LAXATIVE TABS FLEET GENFIBER POWD GLYCERIN HIPREX TABS KRISTALOSE PACK LINZESS 145mcg & 290mcg MAALOX MILK OF MAGNESIA SUSP MINERAL OIL OIL MIRALAX BULK POWD (BRAND) MOVANTIK MOVIPREP POWD PACK NULYTELY SOLR PEG 3350- ELECTROLYTE SOL PEG 3350 POWDER SENNA SENOKOT GRAN SENOKOT SYRP SENOKOT CHILDRENS SYRP SENOKOT XTRA TABS STOOL SOFTENER CAPS</p>	<p>MC/DEL MC MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC MC MC/DEL</p>	<p>ACTIGALL CAPS BENEFIBER CARAFATE CLEARLAX POW COLACE CAPS DIOCTO-C SYRP DOC SOD /CAS CAP DOC-Q-LAX CAPS DOCUSATE SODIUM/CAS CAPS DOK PLUS DULCOLAX SUPP FIBER CON TABS FIBER-LAX TABS GAVILYTE-H GOLYTELY SOLR IBSRELA LINZESS 72mcg⁴ MALTSUPEX MIRALAX PACKETS MOTEGRITY OCALIVA¹ PEG-ELECTROLYTES SOLR PEG 3350 PACKETS PREPOPIK PAK RELISTOR TABS SENEXON TABS SENOKOT TABS SENOKOT S TABS SORBITOL STOOL SOFTENER PLUS CAPS SUFLAVE SUTAB SYMPROIC³</p>	<p>1. PA required to confirm FDA approved indication. 2. For the treatment of carcinoid syndrome diarrhea in combination with somatostatin analog (SSA) therapy in adults inadequately controlled by SSA therapy 3. For the treatment of Opioid Induced Constipation(OIC) 4. Established users will be grandfathered</p>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. Linzess is preferred for adults as treatment of IBS-Constipation AND treatment of chronic idiopathic constipation in adults. Trulance should be avoided in pediatric patients less than 18 years of age.</p>

	MC/DEL		SUCRALFATE TABS	MC/DEL		UNI-CENNA TABS	Use PA Form# 20420	
	MC/DEL		SUPREP SOL	MC		UNI-EASE PLUS CAPS		
	MC		TRULANCE ²	MC		V-R NATURAL SENNA LAXATIV TABS		
	MC		UNI-EASE CAPS	MC		URSO 250		
	MC		URSO FORTE	MC		XERMELO ²		
	MC/DEL		URSODIOL					
MISC. UROLOGICAL								
UROLOGICAL - MISC.	MC		ACETIC ACID 0.25% SOLN	MC		CITRIC ACID/SODIUM CITRAT SOLN	1. Elmiron requires adequate proof of Dx with supportive testing.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC		CYTRA-K SOLN	MC/DEL		CYTRA-2 SOLN		
	MC		FOSFOMYCIN (NDC 82036427401 ONLY)	MC/DEL		ELMIRON CAPS ¹	Use PA Form# 20420	
	MC		K-PHOS MF TABS	MC		FURADANTIN SUSP		
	MC/DEL		METHENAMINE MANDELATE TABS	MC/DEL		MACROBID CAPS		
	MC/DEL		NEOSPORIN GU IRRIGANT SOLN	MC/DEL		MACRODANTIN CAPS		
	MC/DEL		NITROFURANTOIN MONO CAPS	MC/DEL		NITROFURANTOIN MACR SUSP		
	MC/DEL		PHENAZOPYRIDINE HCL TABS	MC		POTASSIUM CITRATE/CITRIC SOLN		
	MC/DEL		PHENAZOPYRIDINE PLUS	MC/DEL		PYRIDIUM PLUS TABS		
	MC		POT CITRATE TAB	MC		PYRIDIUM TABS		
	MC/DEL		PROSED/DS TABS	MC/DEL		RENACIDIN SOLN		
	MC		TRICITRATES SYRP	MC		UROCIT-K		
	MC/DEL		URELIEF PLUS					
	MC		UREX TABS					
	MC/DEL		URISED TABS					
	MC/DEL		UROQID #2 TABS					
PHOSPHATE BINDERS								
PHOSPHATE BINDERS	MC/DEL		CALCIUM ACETATE CAP ¹	MC		AURYXIA ¹	Use PA Form# 20420	Preferred drugs must be tried and failed in step-order due to lack of efficacy or intolerable side effects before less preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		FOSRENOL CHEW ¹	MC/DEL		CALCIUM ACETATE TAB ¹	1. Diag required.	
	MC/DEL		MAGNEBIND - 400 ¹	MC/DEL		ELIPHOS ¹		
	MC		PHOSLYRA ¹	MC/DEL		FOSRENOL PWDR ¹		
	MC/DEL		REVELA ¹	MC		VELPHORO ¹		
INTRA-VAGINALS								
VAGINAL - ANTIBACTERIALS	MC/DEL		CLEOCIN CREA	MC/DEL		METROGEL VAGINAL GEL ¹	1. Dosing limits apply, please see Dosage Consolidation List.	Preferred drugs must be tried and failed in step-order due to lack of efficacy or intolerable side effects before less preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		CLEOCIN SUPP	MC/DEL		VANDAZOLE		
	MC		CLINDESSE CREA	MC		XACIATO		
	MC/DEL		METRONIDAZOLE VAGINAL GEL ¹				Use PA Form# 20420	
	MC/DEL		NUVESSA					
VAGINAL - ANTI FUNGALS	MC/DEL		CLOTRIMAZOLE CREA	MC		AVC CREA	1. Quantity limit: 1/script/2 weeks	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		CLOTRIMAZOLE-3 CREA	MC		CLOTRIMAZOLE 3 DAY CREA		
	MC/DEL		GYNE-LOTRIMIN CREA	MC		GYNAZOLE-1 CREA	Use PA Form# 20420	
	MC		MICONAZOLE CREA	MC		GYNE-LOTRIMIN 3 TABS		
	MC		MICONAZOLE 3 KIT CREA OTC	MC/DEL		MICONAZOLE 3 COMBO PACK KIT ¹		
	MC/DEL		MICONAZOLE 7 CREA	MC/DEL		MICONAZOLE 3 SUPP		DDI: Miconazole will require prior authorization if being used in combination with Warfarin.
	MC/DEL		MICONAZOLE NITRATE CREA	MC		TERAZOL 3 CREA		
	MC		NYSTATIN TABS	MC		TERAZOL 7 CREA		
	MC/DEL		TERCONAZOLE CREAM	MC/DEL		TERCONAZOLE SUPP		
	MC		VAGITROL					
	MC		V-R MICONAZOLE-7 CREA					
VAGINAL - CONTRACEPTIVES							Use PA Form# 20420	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drug will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

VAGINAL - ESTROGENS	MC/DEL MC/DEL		ESTRING RING PREMARIN CREA	MC/DEL MC/DEL		ESTRACE CREA ¹ VAGIFEM TABS ¹	1. Must fail all preferred products before non-preferred. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
VAGINAL - OTHER	MC/DEL MC MC		ACID JELLY GEL ACI-JEL GEL CERVICAL AMINO ACID CREA	MC		AMINO ACID CERVICAL CREA	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BENIGN PROSTATIC HYPERPLASIA (BPH)								
BPH	MC/DEL MC/DEL MC/DEL MC/DEL		DOXAZOSIN MESYLATE TABS FINASTERIDE ¹ 5mg TERAZOSIN HCL CAPS TAMSULOSIN HCL	MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL	5 8 8 8 8 8 8	FLOMAX CP24 ALFUZOSIN AVODART ^{2,4} CARDURA TABS ⁴ ENTADFI ^{5,6} JALYN ^{3,4} PROSCAR TABS ⁴ RAPAFLO ⁴ UROXATRAL ⁴	1. There will be dosing limits of 1 tab per day with out PA. 2. Prior use of preferred agent prior to any approvals. 3. Use of preferred (tamsulosin and finasteride) and (tamsulosin and non-preferred Avodart). 4. Non-preferred products must be used in specified order. 5. Use of individual ingredients preferred (Finasteride and tadalafil). 6. Entadfi® is not recommended for more than 26 weeks Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approval of a non-preferred 5-alpha reductase inhibitor requires objective clinical evidence of a very enlarged prostate rather than just the presence of obstructive urinary outflow symptoms along with adequate trial of preferred Proscar.
ANXIOLYTICS								
ANXIOLYTICS - BENZODIAZEPINES	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ALPRAZOLAM TABS CHLORDIAZEPOXIDE HCL CAPS CLORAZEPATE DIPOTASSIUM TABS DIAZEPAM LORAZEPAM OXAZEPAM CAPS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL	8 8 8 8 8 8 9	ALPRAZOLAM ER ATIVAN LOREEV XR NIRAVAM SERAX TRANXENE XANAX TABS XANAX XR	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANXIOLYTICS - MISC.	MC/DEL MC MC MC/DEL MC/DEL MC/DEL		BUSPIRONE HCL TABS HYDROXYZINE HCL SOLN HYDROXYZINE HCL SYRP HYDROXYZINE HCL TABS ¹ HYDROXYZINE PAMOATE CAPS MEPROBAMATE TABS	MC MC MC/DEL MC/DEL MC/DEL		BUSPAR TABS DROPERIDOL SOLN DROPERIDOL SOLN DROPERIDOL SOLN	Use PA Form# 20420 1. Dosing limits apply, please refer to Dose consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTI-DEPRESSANTS								
ANTIDEPRESSANTS - MAO INHIBITORS	MC/DEL		NARDIL TABS	MC/DEL		TRANLYCYPROMIINE	Use PA Form# 20420	
ANTIDEPRESSANTS - MAO INHIBITORS TOPICAL				MC/DEL		EMSAM ¹	1. Dosing limits apply, please refer to Dose consolidation list. Use PA Form# 20420	Preferred drugs (including a preferred SSRI, a non-SSRI, and Venlafaxine ER) must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIDEPRESSANTS - SELECTED SSRI's AND OTHERS	MC/DEL MC/DEL MC/DEL MC/DEL		BUPROPION HCL TABS BUPROPION SR BUPROPION XL 150mg and 300mg CITALOPRAM	MC/DEL MC MC/DEL MC/DEL	8 8 8 8	APLENZIN ⁴ AUVELITY ¹¹ BUPROPION XL 450mg CELEXA	1. Strong caution with pediatric population. 2. Max daily dose allowed is 120mg, Combination of	Preferred drugs (including failure of at least one preferred SSRI, one SNRI and one non-SSRI/SNRI) must be tried for at least 4 weeks each and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

	MC/DEL	DULOXETINE ^{2,9}	MC	8	CYMBALTA ²	multiple strengths require PA	
	MC/DEL	ESCITALOPRAM	MC/DEL	8	DRIZALMA SPRINKLES	4. Dosing limits allowing 2 tabs/day and a max daily limit of 200mg / day applies. Please see dose consolidation list.	CYMBALTA: Fibromyalgia diagnosis- prior use and failure of preferred generics (amitriptyline or cyclobenzaprine) <u>and</u> gabapentin prior to approval.
	MC/DEL	FLUOXETINE 10mg AND 20mg AND 40mg CAPS	MC/DEL	8	EFFEXOR TABS		
	MC/DEL	FLUOXETINE HCL LIQD	MC/DEL	8	EFFEXOR XR CP24		
	MC/DEL	FLUVOXAMINE MALEATE TABS	MC/DEL	8	FETZIMA ⁷		
	MC/DEL	MIRTAZAPINE	MC/DEL	8	FLUOXETINE 10mg AND 20mg AND 60mg TABS	5. Dosing limits apply, please refer to Dose consolidation list and max daily dose applies. Max daily dose allowed is 375mg.	DDI: Fluvoxamine will now be non-preferred and require prior authorization if it is currently being used with glimepiride (Amaryl).
	MC/DEL	NEFAZODONE	MC	8	FORFIVO XL		DDI: Preferred nefazodone will now be non-preferred and require prior authorization if it is currently being used in combination with either Onglyza 5mg, Enablex 15mg or Vesicare 10mg.
	MC/DEL	PAROXETINE ¹	MC/DEL	8	IRENKA		
	MC/DEL	SERTRALINE HCL	MC/DEL	8	KHEDEZLA		
	MC/DEL	TRAZODONE HCL TABS	MC/DEL	8	LEXAPRO TABS		
	MC/DEL	VENLAFAXINE ER CAPS ⁵	MC	8	LUVOX TABS	6. Non-preferred products must be used in specified step order.	DDI: Fluoxetine will require prior authorization if being used in combination with Plavix.
	MC/DEL	VENLAFAXINE TABS ⁵	MC	8	MAPROTILINE HCL TABS		DDI: Fluvoxamine will require prior authorization if being used in combination with Plavix.
			MC/DEL	8	MIRTAZAPINE ODT		
			MC	8	OLEPTRO	7. Requires previous trials/failure of multiple preferred medications. Dosing limits apply, please see the dose consolidation list. Max daily dose of 80mg if used concomitantly with strong CYP3A4 inhibitor.	SAVELLA: Fibromyalgia diagnosis and trial of a preferred generic amitriptyline, cyclobenzaprine, duloxetine and gabapentin prior to approval.
			MC/DEL	8	PAROXETINE CR ¹		DDI: Drizalma Sprinkle avoid the concomitant use of duloxetine with potent CYP1A2 inhibitors (e.g. fluvoxamine, cimetidine, ciprofloxacin, enoxacin).
			MC/DEL	8	PAXIL ¹		
			MC/DEL	8	PAXIL CR ¹		
			MC/DEL	8	PRISTIQ		Zulresso® is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called the Zulresso® REMS.
			MC	8	PROZAC		
			MC	8	PROZAC CAPS		
			MC	8	PROZAC WEEKLY CPDR		
			MC/DEL	8	REMERON TABS	8. Psychiatry recommended. Please see criteria section.	Spravato: Treatment Resistant Depression
			MC/DEL	8	SARAFEM CAPS		• Must be 18 years of age or older; and medication must be administered under the direct, on site, supervision of a licensed healthcare provider with post-administration observation of a minimum of least 2-hours. The medication must be prescribed by or in consultation with a psychiatrist and prescriber must be enrolled in the REMS program.
			MC/DEL	8	SPRAVATO ⁸	9. Please use multiples of the 20mg, the 40mg is still non-preferred.	• Approval is based upon failure of at least two antidepressants and failure of an antidepressant used adjunctively with one recognized augmentation strategy such as lithium, an atypical antipsychotic, thyroid hormone, etc
			MC/DEL	8	TRAZODONE HCL 300MG TABS		• Ongoing use of Spravato beyond 3 months is based upon a positive response as evidenced by at least a 30 % reduction from baseline as measured by a standardized rating scale
			MC/DEL	8	TRINTELLIX		
			MC	8	WELLBUTRIN TABS	10. For the treatment of patients ≥ 18 years of age.	
			MC	8	WELLBUTRIN SR TBCR		
			MC	8	WELLBUTRIN XL		
			MC/DEL	8	REMERON SOLTAB TBDP	11. Use individual ingredients separately.	Spravato: MDD with Suicidal Ideation
			MC/DEL	8	SAVELLA ⁴		Approval for this indication only if it is started in an inpatient unit, given adjunctively with an optimized antidepressant regimen, and with an 8-12 week initial approval with ongoing use dependent upon documentation of ongoing benefit.
			MC/DEL	8	ZOLOFT		
			MC/DEL	8	ZULRESSO ¹⁰		
			MC/DEL	8	VENLAFAXINE ER TABS ⁵		
			MC/DEL	9	VIIBRYD ⁶		
			MC/DEL	9	FLUOXETINE 90mg TABS ⁵		
						Use PA Form# 20420	
ANTIDEPRESSANTS - TRI-CYCLICS	MC/DEL	AMITRIPTYLINE HCL TABS ¹	MC/DEL		AMOXAPINE TABS	1. Users over the age of 65 require a pa.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	CLOMIPRAMINE HCL CAPS ¹	MC/DEL		ANAFRANIL CAPS		
	MC/DEL	DESIPRAMINE HCL TABS ¹	MC/DEL		DOXEPIH HCL 150 MG ²		
	MC/DEL	DOXEPIH HCL ¹ (not generic Silenor)	MC/DEL		DOXEPIH (generic Silenor)	2. Use multiples of 50mg.	
	MC/DEL	IMIPRAMINE HCL TABS ¹	MC/DEL		NORPRAMIN TABS		
	MC/DEL	NORTRIPTYLINE HCL ¹	MC/DEL		PAMELOR		
	MC	PROTRIPTYLINE HCL TABS ¹	MC		TOFRANIL		
	MC	SURMONTIL CAPS ¹	MC		VIVACTIL TABS		
						Use PA Form# 20420	
						Use PA Form# 10220 for Brand Name requests	
SEDATIVE / HYPNOTICS							
SEDATIVE/HYPNOTICS - BARBITURATE	MC	BUTISOL SODIUM TABS ¹	MC		LUMINAL SOLN	1. PA required for new users of preferred products if over 65 years.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	CHLORAL HYDRATE SYRP ¹	MC/DEL		SOMNOTE CAPS		
	MC	MEBARAL TABS ¹					
	MC/DEL	PHENOBARBITAL ¹					
						Use PA Form# 20420	
SEDATIVE/HYPNOTICS - BENZODIAZEPINES	MC/DEL	DORAL TABS ¹	MC		HALCION TABS ¹	1. Dosing limits apply, please see dosing consolidation list	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	ESTAZOLAM TABS ¹	MC		MIDAZOLAM HCL SYRP		

	MC/DEL		FLURAZEPAM HCL CAPS ¹	MC/DEL		RESTORIL CAPS ¹	consolidation list.	preferred drug(s) exists. Benzodiazepines do cause dependence with continued use and usage should be limited to 7-10 days at a time. Chronic intermittent use (2-3 days per week max) is the standard of care	
	MC/DEL		TEMAZEPAM CAPS 15 & 30MG ¹	MC/DEL		TEMAZEPAM 7.5MG ¹	Use PA Form# 30110		
	MC/DEL		TRIAZOLAM TABS ¹	MC/DEL					
SEDATIVE/HYPNOTICS - Non-Benzodiazepines	MC/DEL	1	MIRTAZAPINE	MC/DEL	7	AMBIEN ¹	1. Quantity Limit of 12 per 34 days. 2. Quantity limits will be allowed up to 30/30, but intermittent therapy is recommended. 3. Only zolpidem trial/failure will be required to obtain Zaleplon. 4. Must fail all preferred products before non-preferred Use PA Form# 30110	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Ambien, Ambien CR, Lunesta, Sonata, Zaleplon and Zolpidem may cause dependence with continued use and as with benzodiazepines, usage should be limited to 7-10 days at a time. Chronic intermittent use (2-3 days per week max) is the standard of care. Please refer to Sedative/Hypnotic PA form. DDI: Belsomra® with strong CYP3A inhibitors (e.g. ketoconazole, itraconazole, posaconazole, clarithromycin, nefazodone, ritonavir, saquinavir, nelfinavir, indinavir, boceprevir, telaprevir, telithromycin, and conivaptan) is not recommended	
	MC	1	TRAZODONE	MC/DEL	7	ESZOPICLONE			
	MC/DEL	1	ZOLPIDEM ²	MC/DEL	7	ZOLPIDEM ER			
	MC/DEL	2	ZALEPLON ^{2,3}	MC/DEL	8	AMBIEN CR ¹			
				MC/DEL	8	BELSOMRA ¹			
				MC	8	DAYVIGO ¹			
				MCDEL	8	EDLUAR			
				MC	8	HETLIOZ			
				MC/DEL	8	INTERMEZZO			
				MC/DEL	8	LUNESTA ¹			
				MC/DEL	8	SONATA CAPS ¹			
				MC/DEL	8	ROZEREM			
				MC	8	QUVIVIQ			
			MC/DEL	8	ZOLPIMIST				
ANTI-PSYCHOTICS									
ANTIPSYCHOTICS - ATYPICALS	MC		ABILIFY MAINTENA	MC	8	ABILIFY ASIMTUFI	If prescribing 2 or more antipsychotics, PA will be required for both drugs, except if one is Clozapine. This also includes combination of Seroquel with Seroquel XR. Use PA form# 20440 for Multiple Antipsychotic requests Use PA form# 10130 for non preferred single therapy atypical requests	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Non preferred atypicals will be approved for patients with FDA-approved indications, and for specific conditions supported by at least two published peer-reviewed double-blinded, placebo-controlled randomized trials that are not contradicted by other studies of similar quality and as long as all first line preferred therapies have been tried and failed at full therapeutic doses for adequate durations (at least two weeks). Prescriptions for quetiapine are limited to a maximum daily dose of 800mg. Uzedy: Establish tolerability with oral risperidone prior to initiating Uzedy Atypicals: Prior Authorization will be required for preferred medication to assure indication is in accordance with FDA approved or literature supported evidence-based best practices. The approved indications are: • schizophrenia • bipolar disorder • agitation related to autism • adjunct in major depressive disorder	
	MC/DEL		ARIPIRAZOLE TAB ³	MC/DEL	8	ABILIFY DISC TAB, INJ and SOL ¹			
	MC		ARISTADA	MC	8	ABILIFY TABS ²			
	MC		ARISTADA INITIO	MC/DEL	8	ARIPIRAZOLE SOL			
	MC/DEL		OLANZAPINE ^{2,3}	MC/DEL	8	ARIPIRAZOLE ODT			
	MC/DEL		OLANZAPINE ^{2,3} ODT	MC	8	CAPLYTA			
	MC/DEL		INVEGA HAFYERA	MC	8	FANAPT			
	MC		INVEGA SUSTENNA	MC/DEL	8	GEODON			
	MC/DEL		INVEGA TRINZA INJ	MC	8	INVEGA			
	MC/DEL		LURASIDONE TAB	MC	8	IGALMI			
	MC/DEL		PALIPERIDONE ER	MC	8	LATUDA			
	MC/DEL		PERSERIS	MC	8	LYBALVI			
	MC		RISPERDAL CONSTA	MC	8	NUPLAZID			
	MC/DEL		RISPERIDONE ODT	MC	8	REXULTI			
	MC/DEL		RISPERIDONE TAB ^{2,3}	MC	8	RISPERDAL TAB			
	MC/DEL		RISPERIDONE SOLN ²	MC	8	RISPERDAL M TAB ¹			
	MC/DEL		QUETIAPINE ^{2,3}	MC	8	RISPERDAL SOLN			
	MC/DEL		QUETIAPINE XR	MC	8	RYKINDO			
	MC		VRAYLAR ⁴	MC/DEL	8	SAPHRIS ¹			1. Established users of single therapy atypicals were grandfathered.
	MC/DEL		ZIPRASIDONE ^{2,3}	MC	8	SECUADO			2. Prior Authorization will be required for preferred medications for members under the age of 5. 3. Dosing limits apply please refer to the dose consolidation list.
				MC/DEL	8	SEROQUEL TABS			
				MC	8	UZEDY			
				MC	8	ZYPREXA TABS			
			MC	8	ZYPREXA RELPREVV				
			MC	8	ZYPREXA ZYDIS TBDP ¹				
			MC/DEL	9	SEROQUEL XR	DDI: The concomitant use of Nuplazid with other drugs known to prolong the QT interval (e.g. Class IA antiarrhythmics, Class 3 antiarrhythmics, antipsychotics, and antibiotics such as gatifloxacin and moxifloxacin).			

						4.Requires step through 1 preferred drug for all indications except AMDD. AMDD requires insufficient response from two antidepressants	Lybalvi: Step through aripiprazole and Latuda. If criteria is met then initial approval for 3 months. Subsequent approvals will be based on evidence of not gaining >= 10 % baseline body weight for ongoing approval. If weight gain >= 10 % of initial body weight, then criteria for ongoing use not met. Invega Hafyera: The patient is started and stabilized on the medication OR The patient has been adequately treated with Invega Sustenna (paliperidone palmitate 1-month) for at least four months or Invega Trinza (paliperidone palmitate 3- month) following at least one 3-month injection cycle.
ANTIPSYCHOTICS - SPECIAL ATYPICALS	MC/DEL		CLOZAPINE TABS	MC/DEL MC/DEL MC/DEL		CLOZAPINE ODT CLOZARIL TABS VERSACLOZ SUSP	Use PA Form# 20420 Preferred generic drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred brand will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Patients previously stabilized on brand name drug will be approved.
ANTIPSYCHOTICS - TYPICAL	MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		CHLORPROMAZINE HCL FLUPHENAZINE DECANOATE FLUPHENAZINE HCL HALDOL HALOPERIDOL HALOPERIDOL DECANOATE SOLN HALOPERIDOL LACTATE SOLN LOXAPINE SUCCINATE CAPS LOXITANE-C CONC MOBAN TABS PERPHENAZINE PROCHLORPERAZINE SERENTIL THIORIDAZINE HCL THIOTHIXENE TRIFLUOPERAZINE HCL TABS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		COMPAZINE COMPRO SUPP FLUPHENAZINE HCL CONC HALDOL DECANOATE LOXITANE CAPS MELLARIL NAVANE CAPS PROLIXIN STELAZINE TABS	Use PA Form# 20420 If prescribing 2 or more antipsychotics, PA will be required for both drugs, except if one is Clozapine. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. If prescribing 2 or more antipsychotics, PA will be required for both drugs, except if one is Clozapine.
LITHIUM							
LITHIUM	MC/DEL MC/DEL		LITHIUM CARBONATE LITHIUM CITRATE SYRP	MC/DEL MC/DEL		ESKALITH CAPS ESKALITH CR TBCR	Use PA Form# 20420
COMBINATION - PSYCHOTHERAPEUTIC							
PSYCHOTHERPEUTIC COMBINATION	MC/DEL MC/DEL		CHLORDIAZEPOXIDE/AMITRIPT PERPHENAZINE/AMITRIPTYLIN				Use PA Form# 20420
STIMULANTS							
STIMULANT - AMPHETAMINES -SHORT ACTING	MC/DEL MC/DEL MC		AMPHETAMINE SALT COMBO ^{1,4} DEXTROAMPHET SULF TABS PROCENTRA	MC/DEL MC MC/DEL MC		ADDERALL TABS EVEKEO METHAMPHETAMINE HCL ZENZEDI	1. Preferred stimulants will be available without PA if diagnosis of ADHD or Narcolepsy. 2. As per recent FDA alert, Adderal & Dexedrinel should not be used in patients with underlying heart defects since they may be at increased risk for sudden death.

						<p>3. Dosing limits apply, please see dosing consolidation list.</p> <p>4. Max daily dose of 50mg.</p> <p>Use PA Form# 20420</p>
STIMULANT - LONG ACTING AMPHETAMINES SALT	<p>MC/DEL</p> <p>MC</p> <p>MC</p>		<p>AMPHETAMINE/DEXTROAMPHET ER^{3,4,7}</p> <p>ADDERALL XR CP24^{1,3,4,7}</p> <p>VYVANSE^{2,3,4}</p>	<p>MC</p> <p>MC</p> <p>MC</p>	<p>MYDAYIS⁵</p> <p>VYVANSE CHEW^{2,3,4,6}</p> <p>FOCALIN IR TABS</p> <p>XELSTRYM⁸</p>	<p>Use PA Form# 20420</p> <p>1. As per recent FDA alert, Adderall should not be used in patients with underlying heart defects since they may be at increased risk for sudden death.</p> <p>2. FDA approval is currently for adults and children 6 or older. Will be available without PA for this age group if within dosing limits. Limit of one capsule daily. Max dose of 70MG daily.</p> <p>3. Preferred stimulants will be available without PA if diagnosis of ADHD.</p> <p>4. Dosing limits apply, please see dosing consolidation list.</p> <p>5. For the treatment of Attention Deficit Hyperactivity Disorder (ADHD) in patients 13 years and older</p> <p>6. Vyvanse chew grace period for current user through June 2022.</p> <p>7. FDA approval is currently for adults and children 6 or older. Will be available without PA for this age group if within dosing limits. Max dose of 50MG daily without a PA.</p> <p>8. For the treatment of patients 6 years of age and older.</p> <p>DDI: The concomitant use of Mydayis® is contraindicated with monoamine oxidase inhibitors (MAOIs) or within 14 days after discontinuing MAOI treatment, as concomitant use can increase hypertensive crisis.</p>
LONG ACTING AMPHETAMINES	<p>MC</p> <p>MC/DEL</p>		<p>DEXTROAMPHET SULF CPSR^{1,3}</p> <p>DEXTROAMPHETAMINE ER</p>	<p>MC/DEL</p> <p>MC</p>	<p>ADZENYS ER³</p> <p>ADZENYS XR- ODT</p>	<p>1. Preferred stimulants will be available without PA if diagnosis of ADHD.</p>

	MC	DYANAVEL XR SUS	MC MC MC	ADZENYS XR ³ DEXEDRINE CAP SR ^{2,3} DYANAVEL XR TAB	2. As per recent FDA alert, Adderall & Dexedrine should not be used in patients with underlying heart defects since they may be at increased risk for sudden death. 3. Dosing limits apply, please see dosing consolidation list. Use PA Form# 20420	DDI: : The concomitant use of Adzenys® XR is contraindicated with monoamine oxidase inhibitors (MAOIs) or within 14 days after discontinuing MAOI treatment.
STIMULANT - METHYLPHENIDATE	MC/DEL MC/DEL MC/DEL MC/DEL	DEXMETHYLPHENIDATE IR TABS METHYLPHENIDATE SOL METHYLPHENIDATE TAB METHYLIN TABS ^{1,2}	MC/DEL MC/DEL MC MC MC/DEL MC/DEL	FOCALIN IR TABS METADATE ER METHYLPHENIDATE HCL CHEW METHYLIN CHEWABLES METHYLIN SOL RITALIN	1. Preferred stimulants will be available without PA if diagnosis of ADHD. Use PA Form# 20420 2. Dosing limits apply, please see dosing consolidation list. Maximum daily doses are as follows: 72mg daily for methylphenidate and 36mg daily for dexmethylphenidate.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please refer to General Criteria category E.
STIMULANT - METHYLPHENIDATE - LONG ACTING	MC MC/DEL MC MC MC/DEL	CONCERTA TBCR DEXMETHYLPHENIDATE CAP ER 50/50 QUILLICHEW ER ^{5,1} QUILLIVANT XR SUS ^{1,5} RITALIN LA ⁴	MC MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	5 METADATE CD CPR 8 ADHANSIA XR ^{2,6} 8 APTENSIO XR ² 8 AZSTARYS ⁶ 8 COTEMPLA XR ² 8 COTEMPLA XR ODT ² 8 DAYTRANA ^{2,3} 8 FOCALIN XR ² 8 JORNAY PM ^{2,6} 8 METHYLPHENIDATE ER CAPS ^{2,4} 8 METHYLPHENIDATE LA CAPS ² 8 METHYLPHENIDATE ER ^{2,4} CAPS 50/50 8 METHYLPHENIDATE ER ² CAPS 40/60 8 METHYLPHENIDATE CD CAPS ² 30-70	1. Preferred stimulants will be available without PA if diagnosis of ADHD. 2. Non-preferred products must be used in specified step order. 3. FDA approval currently only for ages 6-16. Limit of one patch daily. Max dose of 30MG daily. 4. Dosing limits apply, please see dosing consolidation list 5. Quillivant XR and Quillichew ER are only indicated for use in patients 6 years of age and older. 6. For the treatment of patients ≥ 6 years of age. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
STIMULANT - STIMULANT LIKE	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC	ATOMOXETINE HCL ARMODAFINIL CLONIDINE ER GUANFACINE ER MODAFINIL TABS QELBREE ^{6,7}	MC/DEL MC MC MC/DEL MC MC/DEL MC	7 PROVIGIL TABS ³ 7 STRATTERA ^{1,2} 8 CAFKIT SOLN ³ 8 INTUNIV 8 KAPVAY 8 SUNOSI 8 WAKIX	1. Failure of both an amphetamine and methylphenidate is required for consideration for approval of Strattera, unless history of substance abuse without current use of abusable medication(s). Additionally, for patients <17 years of age, a trial of	Provigil requests require diagnosis of Narcolepsy, ADHD, or Obstructive Sleep Apnea. Previous failures of methylphenidate and amphetamine is required for Narcolepsy and ADHD diagnosis, with additional Strattera trial needed with ADHD diagnosis. Please refer to detailed criteria on Provigil PA form Sunsosi is non-preferred and is indicated for to improve wakefulness in adult patients with excessive daytime sleepiness associated with narcolepsy or obstructive sleep apnea (OSA). Wakix is non-preferred and is indicated for the treatment of excessive daytime sleepiness (EDS) in adults with narcolepsy DDI: Sunosi® is contraindicated with MAO inhibitors or within 14 days after discontinuing the MAO inhibitor.

				MC	8	XYREM SOL	<p>guanfacine in required before approval of Strattera.</p> <p>2. Strattera currently has dosing limitations allowing one tablet per day for all strengths if obtain approval. Max daily dose of Strattera is 100mg. Please see dosing consolidation list.</p>	
				MC	8	XYWAV ⁵		<p>Xywav: Diagnosis of cataplexy associated with narcolepsy OR excessive daytime sleepiness associated with narcolepsy. Diagnosis must be confirmed by submission of supporting documentation to include the specialist's interpretation of the Polysomnography (PSG) and Multiple Sleep Latency Test (MSLT) results</p> <p>FDA reminded healthcare professionals and patients that the combined use of Xyrem (sodium oxybate) with alcohol or central nervous system (CNS) depressant drugs can markedly impair consciousness and may lead to severe breathing problems (respiratory depression)</p> <p>DDI: Concomitant use of Qelbree® with an MAO inhibitor or within 2 weeks after discontinuing an MAO inhibitor is contraindicated</p> <p>DDI: Concomitant use of Qelbree® significantly increases the total exposure, but not peak exposure, of sensitive CYP1A2 substrates, which may increase the risk of adverse reactions associated with these CYP1A2 substrates. Coadministration of Qelbree® with sensitive CYP1A2 substrates or CYP1A2 substrates with a narrow therapeutic range (e.g. alosetron, duloxetine, ramelteon, tasimelteon, tizanidine, theophylline), is contraindicated.</p> <p>Use PA Form# 20710 for Provigil, Nuvigil and Xyrem</p> <p>Use PA Form# 20420 for all others</p>
				MC/DEL	9	NUVIGIL ³	3. Non-preferred products must be used in specified	
				MC	9	DESOXYN TABS ³	4. Please use generic Guanfacine.	
				MC	9	DESOXYN CR ³	5. For patients 7 years of age and older with 6. For pediatric patients 6 years of age or older 7. Preferred with a trial and fail either Atomoxetine OR any 2 preferred ADHD agents.	

ANTI-CATAPLECTIC AGENTS

PSYCHOTHERAPEUTIC AGENTS - MISC.				MC		NUDEXTA		
				MC		XENAZINE		Use PA Form# 20710 for Xenazine

WEIGHT LOSS

WEIGHT LOSS							No longer covered: PHENTERMINE, XENICAL, DIDREX, and MERIDIA	Weight loss drugs are not covered as permitted by Federal Medicaid regulations and Maine Medicaid (MaineCare) Policy.
-------------	--	--	--	--	--	--	--	---

ALZHEIMER DISEASE

ALZHEIMER - Cholinomimetics/Others	MC/DEL		DONEPEZIL HYDROCHLORIDE TABS ¹	MC	6	ARICEPT TABS ²	1. PA is required to establish dementia diagnosis and baseline mental status score.	<p>Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>Aduhelm and Leqembi: Testing to rule out reversible causes of dementia (CBC, CMP, TSH, B12, urine drug screen, RPR/VDRL, (folate (if alcohol abuse is present), HIV (if risk present) and an assessment including a review of current medications as a cause of intellectual decline</p> <ul style="list-style-type: none"> - Prescribed by or in consultation with a neurologist or geriatrician or geriatric psychiatrist. Diagnosis of Alzheimer's disease defined as: <ul style="list-style-type: none"> •Confirmed presence of amyloid pathology and mild cognitive impairment or mild dementia stage of disease, consistent with Stage 3 and Stage 4 Alzheimer's disease OR •Confirmed presence of amyloid pathology and prodromal or mild dementia stage of disease, consistent with Stage 3 and Stage 4 Alzheimer's disease -Testing: <ul style="list-style-type: none"> •Clinical Dementia Rating (CDR) global score of 0.5 or 1.0 OR •Repeatable Battery for Assessment of Neuropsychological Status (RBANS) delayed memory index score ≤ 85 OR •Mini-Mental State Examination (MMSE) score of 20-30 OR •Montreal Cognitive Assessment (MoCA) score ≤ 22 - Member is age 50 or older - Obtain recent (within one year) brain magnetic resonance imaging (MRI) prior to initiating treatment - Provider attestation to obtain MRIs prior to the 7th infusion (first dose of 10 mg/kg) and 12th infusion (sixth dose of 10 mg/kg)
	MC/DEL		DONEPEZIL HYDROCHLORIDE ODT ¹	MC	6	ARICEPT ODT ²		
	MC/DEL		EXELON DIS ¹	MC/DEL	7	DONEPEZIL HYDROCHLORIDE TABS 23MG		
	MC/DEL		GALANTAMINE CAPS ¹	MC	8	ADLARITY ³		
	MC/DEL		GALANTAMINE TAB ¹	MC	8	ADUHELM	2. Must fail all preferred products before moving to non-preferred.	
	MC/DEL		MEMANTINE ¹	MC/DEL	8	EXELON CAP		
	MC/DEL		RIVASTIGMINE TARTRATE CAPS ¹	MC/DEL	8	GALANTAMINE HYDROBROMIDE SOL	3. Approvals will require trials and failure or clinical rationale why preferred patches cant be used.	
				MC	8	LEQEMBI ^{1,2}		
				MC/DEL	8	MEMANTINE HCL SOL		
				MC/DEL	8	NAMENDA		
				MC/DEL	8	NAMENDA XR CAPS		
				MC/DEL	8	NAMZARIC		
				MC	8	RAZADYNE ²		

				MC	9	COGNEX CAPS ²		<ul style="list-style-type: none"> - Member does NOT have history or increased risk of amyloid related imaging abnormalities-edema (ARIA-E), which includes brain edema or sulcal effusions and amyloid related imaging abnormalities hemosiderin deposition (ARIA-H), which includes microhemorrhage and superficial siderosis - Member does NOT have hypersensitivity to any components of Aduhelm - Failure of or inability to tolerate at least two other preferred Alzheimer therapies for at least four months each, one of which should include a combination of a cholinesterase inhibitor with memantine - If the initial drug utilized is the combination of a cholinesterase inhibitor and memantine, then only that single trial of two drugs is required
SMOKING CESSATION								
NICOTINE PATCHES / TABLETS	MC/DEL MC/DEL MC/DEL MC/DEL		CHANTIX TAB ¹ CHANTIX STARTER PACK NICOTINE DIS PT24 ¹ VARENICLINE TAB	MC/DEL		NICODERM CQ PT24 ¹	Use PA Form# 20420 1. See criteria section for exemptions	<p>As of July 1, 2014 per MaineCare policy, smoking cessation products will be covered without a copay(including MEDEL). No annual or lifetime limits, must follow FDA approved indications and therapy guidelines.</p> <p>Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>Note: MaineCare policy, smoking cessation product were "not covered" except for during pregnancy between 9/1/12 and 1/1/14, between 1/1/2014 and 7/1/14 smoking cessation products were covered with limitations</p> <p>Patients may qualify for the medication through The Maine Tobacco Helpline if they do not have MaineCare or MEDEL. Patients are encouraged to call The Maine Tobacco helpline at 1-800-207-1230.</p>
NICOTINE REPLACEMENT - OTHER	MC/DEL MC/DEL MC/DEL		NICOTINE POLACRILEX GUM ¹ NICOTINE LOZENGE MINI NICOTINE LOZENGE	MC/DEL MC/DEL MC/DEL MC	8 8 8 8	NICOTROL INHALER ^{1,2} NICOTROL NASAL SPRAY ^{1,2} NICORETTE GUM ^{1,2} NICORETTE LOZENGES	Use PA Form# 20420 1. See criteria section for exemptions 2. Must use non-preferred products in specified step order.	<p>As of July 1, 2014 per MaineCare policy, smoking cessation products will be covered without a copay(including MEDEL). No annual or lifetime limits, must follow FDA approved indications and therapy guidelines.</p> <p>Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>Note: MaineCare policy, smoking cessation product were "not covered" except for during pregnancy between 9/1/12 and 1/1/14, between 1/1/2014 and 7/1/14 smoking cessation products were covered with limitations</p> <p>Patients may qualify for the medication through The Maine Tobacco Helpline if they do not have MaineCare or MEDEL. Patients are encouraged to call The Maine Tobacco helpline at 1-800-207-1230.</p>
ALCOHOL DETERRENTS								
ALCOHOL DETERRENTS	MC/DEL MC MC MC/DEL		ACAMPROSATE ANTABUSE TABS DISULFIRAM TABS NALTREXONE HCL TABS	MC/DEL		ACAMPRO ¹	1. Should only be used in conjunction with formal structured outpatient detoxification program. Use PA Form# 20420	Preferred generic drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MISCELLANEOUS ANALGESICS								
ANALGESICS - MISC.	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL		ACETAMINOPHEN ASPIRIN ASPRIN/ APAP/ CAFF TAB BUTAL/ASA/CAFF BUTALBITAL COMPOUND BUTALBITAL/ACET TABS BUTALBITAL/APAP CAPS BUTALBITAL/APAP/CAFFEINE TABS CHOLINE MAGNESIUM TRISALI DIFLUNISAL TABS EXCEDRIN SALSALATE TABS	MC MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC MC MC		AXOCET CAPS ESGIC-PLUS FIORICET TABS FIORINAL CAPS FIORTAL CAPS FORTABS TABS PHRENILIN TABS PHRENILIN FORTE CAPS TRILISATE LIQD TRILISATE TABS ZEBUTAL CAPS ZORPRIN TBCR	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
LONG ACTING NARCOTICS								
NARCOTICS - LONG ACTING	MC/DEL MC/DEL		FENTANYL PATCH ⁴ BUTRANS ⁴	MC MC	8 8	ARYMO ER AVINZA	Use PA Form# 20510 Use PA form #10300 for	Preferred drugs (Fentanyl Patch, Morphine Sulfate ER tab, Butrans and Embeda) must be tried for at least 2 weeks each & failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Adequate clinical data on initiation/treatment of non-preferred drugs is required.

	MC/DEL MC/DEL MC	MORPHINE SULFATE ER TB12 NUCYNTA ER XTAMPZA ER	MC MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC MC MC MC/DEL	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 9	BELBUCA EXALGO HYSINGLA ER KADIAN METHADONE METHADOSE MORPHABOND ER MORPHINE SULFATE ER CAP MORPHINE SULFATE SUPP MS CONTIN TB12 OPANA ER ORAMORPH SR TB12 OXYCONTIN TB12 ¹ XARTEMIS ER ZOHYDRO ER OXYCODONECONC OXYCODONE ER ^{3,5}	PAs over the opiate limit 1. Oxycotin will be available without PA for patients treated for or dying from cancer or hospice patients. CA (cancer) or HO (hospice) diag code may be used but store must verify since all scripts will be audited and stores will be liable. 2. Established users are grandfathered. 3. Oxycodone ER allowed only 2 per day for all strengths except 80 mg, please see attached to 4. Dosing limits apply. Please see dose consolidation list. 5. Non-preferred products must be used in specific order. 6. Methadone will be available without PA for patients treated for or dying from cancer or hospice patients or similar conditions as supported by clinical documentation. CA (cancer) or HO (hospice) diag code may be used but store must verify since all scripts will be audited and stores will be liable.	or the preferred drug or a significant potential drug interaction between another drug & the preferred drug(s) exists. Adequate trials include prevention/treatment or common adverse effects associated w/ narcotics (antinausea, antipruritics, etc.) as well as adequate equianalgesic dosing when converting from one narcotic to another. Also, adequate documentation of attempts to titrate dose of preferred agents to achieve adequate pain relief & desired clinical response must be provided. Member's drug regimen for additions &/or discontinuations of medications that may affect absorption &/or metabolism of preferred agents must be monitored. Approvals will not be granted if patient had access to either non-preferred products or high doses of short acting narcotics during the trial period. Non-preferred drugs will not be approved for patients showing evidence of usage patterns consistent w/ controlled substance abuse such as: 1.Frequent or persistent early refills of controlled drugs; 2.Multiple instances of early refill overrides due to reports of misplacement, stolen, dropped in toilet or sink, distant travel, etc.; 3.Breaches of narcotic contracts with any provider; 4.Failure to comply with patient responsibilities in attached opioid documentation (see PA form) including but not limited to failing to submit to and pass pill counts; 5.Failing to take or pass random drug testing; 6.Failing to provide old records regarding prior use of narcotics; 7.Receiving controlled substances from other prescribers that the provider submitting the PA is unaware of 8.Documented history of substance abuse. Substance abuse evaluations may be required for patients with medical records displaying documented substance abuse or potential signs of narcotic misuse and abuse such as chronic early refills, short dosing intervals, frequent dose increases, multiple lost/stolen etc scripts and intolerance or "allergy" to all products but Oxycotin. 9.Circumventing MaineCare prior authorization requirements for narcotics by paying cash for affected narcotics (prescribers failed to submit prior authorization prior to cash narcotic scripts being filled by member). 10.Requests for any Brand name controlled substance, considered by authorities to be highly abused and diverted (Oxycontin, Percocet, Tylox, Vicodin, Dilaudid, Ultracet...) with an available AB rated generic equivalent will be denied unless it will be provided in a setting that virtually eliminates the risk of diversion. 11.Allergic reactions to any product within a specific narcotic class will justify and preclude use of any other product in the same class due to the risk of cross-hypersensitivity. Hysingla ER- Concomitant use should be avoided with mixed agonist/antagonist analgesics, partial agonist analgesics, and MAOIs. Verify prior trials and failures or intolerance of preferred treatments Methadone – Established users must have a trial and failure of at least 2 preferred drugs for least 2 weeks. Otherwise they will be allowed 180 days to transition to a preferred product.
NARCOTICS - SELECTED	MC/DEL MC/DEL	TRAMADOL HCL TABS TRAMADOL/APAP TABS	MC/DEL MC MC/DEL MC MC MC MC MC MC MC MC	7 8 8 8 8 8 8 8 8 8 9	RYZOLT BUPRENEX SOLN BUTORPHANOL NALBUPHINE HCL SOLN QDOLO SOLN SEGLENTIS ¹ STADOL NS SOLN TRAMADOL ER ULTRACET TABS ¹ ULTRAM ER	Use PA Form# 20420 Use PA form #10300 for PAs over the opiate limit 1. Only available if component ingredients are unavailable.	Preferred drugs from this and other narcotic classes must be tried for at least 2 weeks each and failed due to lack of efficacy or intolerable side effects before non-preferred drugs from this class will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approvals will not be granted if patient had access to either non-preferred products or high doses of short acting narcotics during the trial period. Substance abuse evaluations may be required for patients with medical records displaying potential signs of narcotic misuse and abuse such as chronic early refills, short dosing intervals, frequent dose increases, multiple lost/stolen etc scripts and intolerance or "allergy" to all products but desired product. Allergic reactions to any product within a specific narcotic class will justify and preclude use of any other product in the same class due to the risk of cross-hypersensitivity. Non-preferred drugs will not be approved for patients showing evidence of usage patterns consistent with controlled substance abuse such as: 1.frequent or persistant early refills of controlled drugs; 2.multiple instances of early refill overrides due to reports of misplacement, stolen, dropped in toilet or sink, distant travel; 3.breaches of narcotic contracts with any provider; 4.failure to comply with patient responsibilities in attached opioid documentaion (see PA form) including but not limited to failing to submit to and pass pill counts; 5.failing to take or pass random drug testing; 6.failing to provide old records regarding prior use of narcotics; 7.receiving controlled substances from other prescribers that the provider submitting the PA is unaware of. in Substance abuse evaluations may be required for patients with medical records displaying potential signs of narcotic misuse and abuse such as chronic early refills, short dosing intervals, frequent dose increases, multiple lost/stolen etc scripts and intolerance or "allergy" to all products but Oxycotin. Allergic reactions to any product within a specific narcotic class will justify and preclude use of any other product in the same class due to the risk of cross-hypersensitivity.

Beginning January 2017, all current opiate users who are above the maximum combined daily dose of 100 MME must titrate their total daily dose of opioid medications below 300 MME. Also, the maximum daily supply of an opiate prescription for acute pain will be limited to 7-day supplies. The maximum day supply of an opiate prescription for chronic pain will be limited to 30-day supplies. As of July 1, 2017 all users of opioid medications must comply with the maximum combined daily dose of 100 MME.

However, for MaineCare members, effective January 1, 2017, opioid prescription(s) for more than a 7-day supply and/or more than 30 MME/ day will require a prior authorization. Please note that MaineCare implemented a 30 MME limit January 1, 2013 that is still effective.

Post-surgical members may receive prior authorizations for opiates up to a 60 days in length if medical necessity is provided by the surgical provider.

An MME conversion chart is available at www.mainearepd.org. Click on "General Pharmacy Info."

Please see the Pain Management Policy tab for the complete criteria

MISCELLANEOUS NARCOTICS

MISCELLANEOUS NARCOTICS						
NARCOTICS - MISC.	MC/DEL	ACETAMINOPHEN/CODEINE	MC/DEL	8	ABSTRAL	1. Fentanyl OT loz (Barr) and Capital and codeine suspension products require PA for users over 18 years of age. PA is not required if under 18 years of age.
	MC/DEL	ASPIRIN/CODEINE TABS	MC/DEL	8	APADAZ	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please refer to General Criteria category E.
	MC/DEL	BUTAL/ASA/CAFF/COD CAPS	MC/DEL	8	ASCOMP/CODEINE CAPS	
	MC	BUTALBITAL/ASPIRIN/CAFFEI CAPS	MC/DEL	8	BUTALBITAL/APAP/CAFFEINE/ CAPS	
	MC	CAPITAL AND CODEINE SUSP ¹	MC/DEL	8	BUTALBITAL COMPOUND- CODEINE CAP	
	MC	CAPITAL/CODEINE SUSP ¹	MC	8	DEMEROL	
	MC/DEL	CODEINE PHOSPHATE SOLN	MC/DEL	8	DILAUDID	
	MC/DEL	CODEINE SULFATE TABS	MC	8	DILAUDID-HP SOLN	2. Oxycodone/acet 10/650 is 8 times more expensive. Use twice as many of oxycod/acet 5/325 instead.
	MC/DEL	ENDOCET TABS ³	MC	8	FENTANYL CITRATE SOLN	You can mix andmatch preferred strengths of oxycodone and acet. dose similar to certain non-preferred drugs.
	MC/DEL	ENDODAN TABS	MC/DEL	8	FENTORA	
	MC/DEL	FENTANYL OT LOZ ¹	MC/DEL	8	FIORICET/CODEINE CAPS	However, for MaineCare members, effective January 1, 2017, opioid prescription(s) for more than a 7-day supply and/or more than 30 MME/ day will require a prior authorization. Please note that MaineCare implemented a 30 MME limit January 1, 2013 that is still effective.
	MC/DEL	FENTANYL OT LOZ1	MC	8	FIORINAL/CODEINE #3 CAPS	
	MC/DEL	HYDROCODONE/ACETAMINOPHEN	MC	8	FIORTAL/CODEINE CAPS	
	MC/DEL	HYDROMORPHONE HCL ³	MC/DEL	8	HYDROCODONE/IBUPROFEN	Post-surgical members may receive prior authorizations for opiates up to a 60 days in length if medical necessity is provided by the surgical provider.
	MC	LORTAB ELX	MC/DEL	8	HYDROMORPHONE ER	
	MC/DEL	MEPERIDINE SOL	MC/DEL	8	HYDROMORPHONE RECTAL SUPP	An MME conversion chart is available at www.mainearepd.org . Click on "General Pharmacy Info."
	MC/DEL	NUCYNTA	MC	8	IBUDONE	
	MC/DEL	OXYCODONE TAB	MC/DEL	8	LEVORPHANOL TARTRATE TAB	
	MC/DEL	OXYCODONE/ACETAMINOPHEN ^{2,3}	MC/DEL	8	LORCET	3. Only preferred manufacturer's products will be available without prior authorization.
	MC/DEL	ROXICET	MC	8	LORTAB	
	MC	ROXIPRIN TABS	MC	8	MAXIDONE TABS	Please see the Pain Management Policy for the complete criteria
			MC/DEL	8	MEPERIDINE TABS	
			MC/DEL	8	NORCO TABS	
			MC/DEL	8	ONSOLIS	
			MC/DEL	8	OXECTA	
			MC/DEL	8	OXYCODONE CAP	
			MC/DEL	8	OXYCODONE/APAP 10/650	
			MC/DEL	8	OXYCODONE/APAP 7.5/500	
			MC/DEL	8	PENTAZOCINE/ACET TABS	
			MC/DEL	8	PENTAZOCINE/NALOXONE TABS	
			MC	8	PERCOCET TABS	
			MC	8	PERCOCET TABS	
			MC	8	PHRENILIN W/CAFFEINE/CODE CAPS	
			MC/DEL	8	ROXICET 5/500 TABS	
			MC	8	ROXICODONE TABS	
			MC/DEL	8	ROXYBOND	
			MC	8	SYNALGOS-DC CAPS	
			MC	8	TALACEN TABS	

			MC	8	TREZIX		
			MC	8	TYLENOL/CODEINE #3 TABS		
			MC	8	TYLOX CAPS		
			MC	8	XOLOX	Use PA Form# 20420	
			MC	8	VICODIN		
			MC	8	VICOPROFEN TABS	Use PA form #10300 for PAs over the opiate limit	
			MC	8	ZYDONE TABS		
			MC	9	ACTIQ LPOP		
			MC	9	CONZIP		
			MC	9	OPANA		
OPIOID DEPENDENCE TREATMENTS	MC				SUBOXONE FILM ²		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		MC		BUPRENORPHINE/NALOXONE TABS ²		
			MC/DEL		BRIXADI	Use PA Form #20100	Members will continue to be required to follow the criteria listed below:
			MC		BUPRENORPHINE ^{1,2}		1-Induction period for 30 days
			MC		PROBUPHINE ³	1. Buprenorphine will only be approved for use during pregnancy.	2-Members will be allowed multiple induction periods per year where they can receive max 24 mg Daily for up to 30-days, without a PA once they have been on a maintenance dose.
			MC		SUBLOCADE	2. See Criteria Section	3-Max dose of 24 mg for induction
			MC		ZUBSOLV	3. Both provider and patient need to be registered as part of the Probuphine REMS program	4-Max dose of 16 mg for maintenance
						Use PA form #20200 for Extended Release Buprenorphine	5- Suboxone will not require a PA if patient requires concomitant use of an opioid for acute pain.
							6- Should be evidence provided of monthly monitoring including random pill counts urine drug tests and prescription monitoring program reports.
							7- Buprenorphine monotherapy is preferred if member is pregnant and dose not > 16 mg day and pregnancy diagnosis is noted on the prescription.
							Brixadi and Sublocade: The prescriber can attest (and medical record should document) that: -member has a documented history of opioid use disorder (OUD), -XRB is being used for the treatment of OUD (rather than pain or any other non-FDA approved indication) and -member's total daily dose of sublingual buprenorphine is less than or equal to 24 mg daily. AND at least one of the following is true: -The member's previous use of sublingual buprenorphine has included misuse, overuse, or diversion. -The member is at high risk of overdose (e.g., individuals leaving incarceration or abstinence-based treatment programs; individuals who are unhoused; or those facing potential gaps in care due to delays in care or geographically limited treatment access). -The member has experienced significant medical complications of OUD and/or of injection drug use. Occurrence should be in the last 5 years, or it should be clearly documented that the risk indicated by this infection or complication is ongoing (Examples of medical complications of OUD include: threatened the function of organs or life or limb threatening and required medical and/or surgical therapy. Examples of medical complications of injection drug use include osteomyelitis, endocarditis, renal failure, joint infection or other serious medical complications directly related to OUD.) -The member has treatment-resistant OUD, including those with ongoing illicit substance use in the context of sublingual buprenorphine treatment as documented by positive urine drug screens or other clear objective evidence, and/or further functional decline with explicit documentation of the functional decline. -The member has a significant intolerance of, or documented allergy to, sublingual buprenorphine (either buprenorphine monotherapy or buprenorphine/naloxone combination therapy) that has resulted in the patient's inability to comply with continued treatment using the sublingual product. (A true allergy is usually accompanied by rash, respiratory symptoms, or anaphylaxis. Other complaints such as bad taste, mouth tingling, etc. do not constitute evidence of allergy or significant intolerance. Formulation preference or convenience are not, in and of themselves, indications for using XRB.) -The member is in ongoing treatment with XRB and would like to continue the medication.
OPIOID WITHDRAWAL AGENTS			MC		LUCEMYRA ¹		
						1. Clinical PA for appropriate approved use and patient has documented contraindication to clonidine Use PA Form#20420	
NARCOTIC ANTAGONISTS							
NARCOTIC - ANTAGONISTS	MC/DEL		MC		NALTREXONE HCL TABS		
	MC		MC		NALOXONE INJ	Use PA Form# 20420	
					EVZIO		
					OPVEE ²		

	MC	NARCAN NS				1. Will only be approved for side effects experienced with generic that are not described in the literature as occurring with the brand version.
	MC MC MC	NALOXONE SPRAY OTC VIVITROL INJ ZIMHI	MC MC/DEL	KLOXXADO REVIA TABS ¹		2. For the treatment of adult and pediatric patients 12 years of age and older.
COX 2 / NSAIDS						
COX 2 INHIBITORS - SELECTIVE / HIGHLY SELECTIVE	MC/DEL MC/DEL MC/DEL MC/DEL	CELECOXIB ^{4,5} KETOROLAC TROMETHAMINE ^{2,3,5} NABUMETONE TABS ⁵ MELOXICAM TABS ^{1,5}	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	CELEBREX CAPS ^{4,5} MELOXICAM CAPS ⁵ MOBIC ⁵ MOBIC SUSP ⁵ RELAFEN TABS ⁵ QMIIZ ODT VIVLODEX	Use PA Form# 20420 1. Meloxicam has dosing limits allowing one tablet daily of all strengths without PA. 2. Ketorolac Tromethamine is indicated for the short term (up to 5 days) management of moderately severe acute pain that requires analgesic at the opioid level in adults. Not indicated for minor or chronic pain conditions. 3. Ketorolac has dosing limits allowing 24 tablets for a 5 day supply every 30 days. 4. Dosing limits will be set at a maximum of 400mg daily 5. The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
NSAIDS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	CHILDRENS IBUPROFEN DICLOFENAC POTASSIUM TABS DICLOFENAC SODIUM TABS DICLOFENAC SODIUM 1% GEL ¹ ETODOLAC FENOPROFEN CALCIUM TABS FLURBIPROFEN TABS IBUPROFEN INDOMETHACIN KETOPROFEN MECLOFENAMATE SODIUM CAPS NAPROSYN SUSP NAPROXEN SUSP NAPROXEN TABS NAPROXEN SODIUM TABS	MC MC MC MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL	ADVIL TABS ANAPROX TABS ANAPROX DS TABS CAMBIA CATAFLAM TABS CHILDRENS ADVIL SUSP CHILD'S IBUPROFEN SUSP CHILDREN'S MOTRIN SUSP CLINORIL TABS DAYPRO TABS DICLIFENAC GEL EC-NAPROSYN TBEC ETODOLAC ER 600MG FELDENE CAPS FLECTOR PATCH	The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use. 1. Dosing limits apply, please see Dosage Consolidation List. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approvals will be granted for other requests based on failure of at least one generic NSAID from at least 3 different NSAID classes as described in the COX-II PA form. DDI: Diclofenac will now be non-preferred and require prior authorization if it is currently being used in combination with Ilescol. The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use.

	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		NAPROXEN SODIUM CAPS NAPROXEN DR TBEC OXAPROZIN TABS SULINDAC TABS TOLMETIN SODIUM VOLTAREN GEL	MC/DEL MC MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC MC MC MC MC MC MC		IBU-200 INDOCIN LICART LODINE LOFENA MOTRIN NALFON CAPS NAPRELAN TBCR NAPROSYN TABS NAPROXEN SODIUM TBCR PENNSAID PIROXICAM CAPS PONSTEL CAPS RELAFEN DS SB IBUPROFEN TABS SPRIX TIVORBEX TOLECTIN V-R IBUPROFEN TABS ZORVOLEX			
NSAID - PPI				MC MC/DEL		PREVACID NAPRA-PAC VIMOVO ¹		1. Use a preferred NSAID and PPI separately. Use PA Form# 20420	
RHEUMATOID ARTHRITIS									
RHEUMATOID ARTHRITIS	MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL		ACTEMRA VIALS ACTEMRA SYRINGES AVSOLA AZATHIOPRINE ENBREL ² ENBREL SURECLICK ² KINERET SOLN LEFLUNOMIDE METHOTREXATE ORENCIA SULFASALAZINE TABS SIMPONI PEN SIMPONI AUTOINJECTOR HUMIRA ^{1,2} XELJANZ ^{3,6} XELJANZ XR XELJANZ XR SOL	MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC/DEL MC MC MC/DEL MC/DEL MC MC MC MC MC		AMJEVITA ARAVA CIMZIA CYLTEZO ENTYVIO HADLIMA HULIO HYDROXYCHLOROQUINE ² HYRIMOZ IDACIO ILARIS ^{1,3,4} INFLECTRA INFLIXIMAB VIAL KEVZARA OLUMIANT OTREXUP RASUVO ⁷ REDITREX REMICADE RENFLEXIS RINVOQ YUFLYMA YUSIMRY XATMEP ⁹		Use PA Form# 20900 1. Dosing limits apply. Please see dose consolidation list. 2. Established users will be grandfathered. 3. Clinical PA is required to establish diagnosis and medical necessity. 4. Verification of age for appropriate indication. 5. Treatment failure or intolerance to other forms of preferred methotrexate 6. See criteria section	See criteria as listed on Rheumatoid Arthritis PA form. Preferred injectable products allowed without PA if trial of a preferred oral agents (azathioprine, hydroxychloroquine, leflunomide, methotrexate, sulfasalazine tabs) are seen in the members drug profile. Dosing limits apply. Xeljanz is limited to adults with moderate to severe RA and UC who have had an inadequate response or intolerance to methotrexate. Should not be used concomitantly with biologic DMARDs or potent Immunosuppressants. DDI: The concomitant use of Xeljanz® XR with biologic DMARDs or potent immunosuppressants such as azathioprine and cyclosporine are not recommended. The concomitant use of Xeljanz® XR with potent CYP3A4 inducers (e.g. rifampin) is not recommended
ALOPECIA AREATA AGENTS									
ALOPECIA AREATA AGENTS				MC MC/DEL	7 8	OLUMIANT LITFULO			Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

[Use PA Form# 20420](#)

MISCELLANEOUS ARTHRITIS

ARTHRITIS - MISC.	MC MC		RIDAURA CAPS MYOCHRYSLINE SOLN	MC/DEL	ARTHROTEC ¹	1. The individual components of Arthrotec are available without PA. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. The individual components of Arthrotec are available without PA.
-------------------	----------	--	-----------------------------------	--------	------------------------	---	---

LUPUS-SLE

LUPUS-SLE				MC MC MC	BENLYSTA ¹ LUPKYNIS SAPHNELO	Use PA Form# 20420 1. Approvals will require previous trial of corticosteroids, antimalarials, NSAIDs and immunosuppressives.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Lupkynis is a sensitive CYP3A4 substrate. Co-administration with strong or moderate CYP3A4 inhibitors increases voclosporin exposure, which may increase the risk of Lupkynis® adverse reactions. Co-administration of Lupkynis® with strong CYP3A4 inhibitors (e.g. ketoconazole, itraconazole, clarithromycin) is contraindicated. Reduce Lupkynis® dosage when co-administered with moderate CYP3A4 inhibitors (e.g. verapamil, fluconazole, diltiazem)
-----------	--	--	--	----------------	---	--	---

PIK3CA-Related Overgrowth Spectrum (PROS)

PIK3CA-Related Overgrowth Spectrum (PROS)				MC	VJOICE ¹	Use PA Form# 20420 1. PA required to confirm FDA approved indication.	Preferred drugs must be tried and failed, in step-order, due to lack of efficacy (failure to reach target IOP reduction) or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
---	--	--	--	----	---------------------	--	--

MIGRAINE THERAPIES

MIGRAINE - ERGOTAMINE DERIVATIVES				MC/DEL MC	D.H.E. 45 SOLN TRUDHESA	Use PA Form# 10110	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
-----------------------------------	--	--	--	--------------	----------------------------	------------------------------------	--

MIGRAINE - CARBOXYLIC ACID DERIVATIVES	MC		DIVALPROEX ER TB24	MC	DEPAKOTE ER TB24	Use PA Form# 10110	
--	----	--	--------------------	----	------------------	------------------------------------	--

MIGRAINE - SELECTIVE SEROTONIN AGONISTS (5HT)--Tabs/Nasal	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	1 1 1 1 1 1 2	MIGRANAL NASAL SPRAY RELPAK ¹ RIZATRIPTAN ODT RIZATRIPTAN TABS SUMATRIPTAN TABS ¹ ZOLMITRIPTAN TAB ¹ NARATRIPTAN HCI TABS ¹	MC MC MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	AMERGE TABS ^{1,2} AXERT TABS ^{1,2} FROVA TABS ^{1,2} IMITREX NASAL SPRAY ¹ IMITREX TABS ^{1,2} MAXALT ^{1,2,3} MAXALT MLT ^{1,2,3} ONZETRA XSAIL ² SUMATRIPTAN NASAL SPRAY ¹ ZOLMITRIPTAN ODT ZOLMITRIPTAN SPRAY ZOMIG TABS ^{1,2} ZOMIG NASAL SPARY ^{1,2} ZOMIG ZMT TBDP ^{1,2}	1. All drugs in this category have dosing limits. Please refer to dose consolidation table. 2. Must fail all preferred products before non-preferred. 3. Established users will be grandfathered Use PA Form# 10110	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Quantity limit exceptions will require ongoing therapy with therapeutic doses of highly effective prophylactic medication as listed on the Triptan PA form.
---	--	---------------------------------	---	--	---	--	--

MIGRAINE - SELECTIVE SEROTONIN AGONISTS (5HT)--Injectables	MC MC/DEL MC/DEL		IMITREX CARTRIDGE ¹ SUMATRIPTAN SYRINGE ¹ SUMATRIPTAN PEN INJCTR ¹	MC/DEL MC MC	TOSYMRA ZEMBRACE ¹ IMITREX PEN INJCTR ¹	Use PA Form# 10110 1. Dosing limits apply. Please refer to the dose consolidation table.	
--	------------------------	--	---	--------------------	---	---	--

MIGRAINE - SELECTIVE SEROTONIN AGONISTS (5HT)--Combinations				MC/DEL	TREXIMET ^{1,2}	Use PA Form# 10110 1. Dosing limits apply. Please see dose consolidation list.	
---	--	--	--	--------	-------------------------	---	--

						2. Use preferred Sumatriptan and Naproxen separately. Treximet only available if component ingredients of sumatriptan and naproxen are unavailable.		
MIGRAINE - MISC.	MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL		AIMOVIG ¹ AJOVY ¹ AJOVY AUTO INJCT ¹ EMGALITY SYRINGE ¹ 200mg/ml EMGALITY PEN ¹ NURTEC ODT ² SPASTRIN TABS	MC MC MC/DEL MC/DEL MC MC MC MC/DEL		BELCOMP-PB SUPP ELYXYB MIGRAZONE CAPS MIGERGOT SUP QULIPTA REYVOW ² UBRELVY ² VYEPTI ² ZAVZPRET ²	Use PA Form# 10110 1. See criteria section 2. Dosing limits apply, please see the dose consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Aimovig, Ajovy and Emgality: The patient is 18 years of age or older AND patient has a diagnosis of episodic migraine (4-14 headache days per month with migraine lasting 4 hours or more) or chronic migraine (≥ 15 headache days per month, of which ≥ 8 are migraine days, for at least 3 months) AND patient has failed or has a contraindication to an adequate trial (≥ 60 days) of at least 2 medications for migraine prophylaxis from at least 2 different classes. Ubrelyv is non-preferred and is indicated for the acute treatment of migraine with or without aura in adults. This is not indicated for the preventive treatment of migraine. Reyvow is non-preferred and is indicated for the acute treatment of migraine with or without aura in adults. Reyvow® is not indicated for the preventive treatment of migraine. Zavzpret: The patient must have a documented side effect, allergy, or treatment failure to preferred oral CGRP Inhibitor and two non-preferred oral CGRP Inhibitors. Nurtec ODT will be preferred after 2 adequate trials of at least two preferred triptans
GOUT								
GOUT	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ALLOPURINOL TABS COLCHICINE TAB FEBUXOSTAT TAB PROBENECID TABS PROBENECID/COLCHICINE TABS	MC/DEL MC MC MC/DEL MC MC		COLCHICINE CAP COLCRYS GLOPERBA ULORIC ¹ MITIGARE ZYLOPRIM TABS	Use PA Form# 20420 1. Failure of therapeutic (300mg) dose of Allopurinol (failure define as not being able to get uric acid levels below 6mg/dl) or severe renal disease.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: The concomitant use of Gloperba® and CYP3A4 inhibitors (e.g. clarithromycin, ketoconazole, grapefruit juice, erythromycin, verapamil, etc.) should be avoided due to the potential for serious and life-threatening toxicity.
MISC.								
ACID SPHINGOMYELINASE DEFICIENCY (ASMD)				MC		XENPOZYME ^{1,2}	1.For treatment of non-central nervous system manifestations of acid sphingomyelinase deficiency (ASMD) in adult and pediatric patients 2. Clinical PA required for appropriate diagnosis and clinical parameters.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANESTHETICS - MISC.	MC MC MC		BUPIVACAINE HCL SOLN LIDOCAINE HCL SOLN MARCAINE SOLN	MC MC/DEL MC		SENSORCAINE-MPF SOLN SYNVISC INJ XYLOCAINE SOLN	Use PA Form# 30130	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
COLD AGGLUTININ DISEASE (CAD)				MC		ENJAYMO ¹	1.Indicated to decrease the need for red blood cell transfusion due to hemolysis in adults with cold agglutinin disease (CAD).	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

PRIMARY HYPEROXALURIA TYPE 1 (PH1)						OXLUMO ¹	1. PA is required to establish diagnosis and medical necessity. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
SICKLE CELL DISEASE	MC/DEL MC		HYDROXYUREA DROXIA	MC MC MC MC/DEL		ADAKVEO ENDAR ¹ OXBRYTA ² SIKLOS	1. Evidence of other preferred L-glutamine products utilization and reason for failure. 2. For the treatment of patients ≥ 12 years of age. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: The concomitant use of Oxbryta and strong CYP3A4 inhibitors or fluconazole may increase voxelotor plasma levels and may lead to increased toxicity.
HUTCHINSON- GILFORD PROGERIA SYNDROME (HGPS)				MC		ZOKINVY ^{1,2}	1. In patients 12 months of age and older with a body surface area (BSA) of 0.39m ² and above 2. PA required to confirm FDA approved indication. Use PA Form# 20420	ZOKINVY: To reduce the risk of mortality in Hutchinson-Gilford Progeria Syndrome (HGPS). For the treatment of processing-deficient Progeroid Laminopathies with either: Heterozygous LMNA mutation with progerin-like protein accumulation OR Homozygous or compound heterozygous ZMPSTE24 mutations
VACCINES	MC/DEL MC MC/DEL MC/DEL		ABRYVVO AREXVY GARDASIL 9 SHINGRIX				Use PA Form# 20420	Gardasil 9 will be preferred by MaineCare for ages 19-45 for FDA approved indications. Under the Maine Immunization Program Gardasil 9 is covered under the Vaccine for Children Program for ages 9-18. Please contact 1-800-867-4775 or 207-287-3746 for assistance. Abryvvo will be a preferred vaccine indicated for active immunization for the prevention of lower respiratory tract disease (LRTD) caused by respiratory syncytial virus (RSV) in individuals 60 years of age and older. Active immunization of pregnant individuals at 32 through 36 weeks gestational age for the prevention of LRTD and severe LRTD caused by RSV in infants from birth through 6 months of age. Arexvy will be preferred for active immunization for the prevention of LRTD caused by respiratory syncytial virus (RSV) in individuals 60 years of age and older. SHINGRIX (>= 50yo) is preferred as of 11-20-20 with respective age edit.
APDS				MC		JOENJA ^{1,2,3}	Use PA Form# 20420 1. Clinical PA required for appropriate diagnosis 2. For the treatment of patients 2 years of age and older. 3. Avoid CYP3A drug drug interaction.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ALPHA- MANNOSIDOSIS				MC		LAMZEDE	Use PA Form# 20420 1. Clinical PA required for appropriate diagnosis	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTI-CONVULSANTS								
ANTICNVLSANTS	MC/DEL MC MC/DEL MC/DEL MC/DEL		CARBAMAZEPINE CARBAMAZEPINE ER CAP CARBATROL CP12 CELONTIN CAPS CLOBAZAM	MC MC MC/DEL MC MC	8 8 8 8 8	APTIOM BANZEL BRIVIACT ⁷ CARBAMAZEPINE SUS DEPAKOTE	Use PA Form# 20420 All non-preferred meds must be used in specified order	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

MC/DEL	CLONAZEPAM TABS	MC	8	DEPAKOTE ER		
MC	DEPAKOTE SPRINKLES CPSP	MC	8	DIACOMIT	1. Quantity limit. 5/month	
MC/DEL	DIASTAT ¹	MC/DEL	8	DIVALPROEX SODIUM SPRINKLE CAPS	2. Dosing limits apply, please see dose consolidation list.	Approvals will be for patients with a variety of drug-specific FDA-approved indications and for specific conditions supported by at least two published peer-reviewed double-blinded, placebo-controlled randomized trials that are not contradicted by other studies of similar quality after recommendation by the DUR Committee and as long as all first line therapies have been tried and failed at full therapeutic doses for adequate durations (at least two weeks).
MC/DEL	DIAZEPAM GEL ¹	MC	8	ELEPSIA XR ¹⁰		
MC/DEL	DILANTIN	MC	8	EPRONTIA SOLN ¹¹		
MC/DEL	DIVALPROEX SODIUM	MC/DEL	8	FELBATOL	3. Dosing limits apply per strength as well as a maximum daily dose of 600mg. Please see dose consolidation list.	
MC	DIVALPROEX SPRINKLE CAP	MC/DEL	8	FELBATOL SUS		
MC/DEL	EPIDIOLEX ⁸	MC/DEL	8	FELBAMATE SUS		
MC/DEL	EPITOL TABS	MC	8	FINTEPLA ⁹		
MC/DEL	ETHOSUXIMIDE SYRP	MC	8	FYCOMPA ²		
MC/DEL	EQUETRO	MC/DEL	8	HORIZANT	4. Adjunctive therapy 17 and older.	*** SEE CHART AT END OF DOCUMENT
MC/DEL	GABAPENTIN ² CAP	MC	8	GRALISE		
MC/DEL	GABAPENTIN ² TAB	MC/DEL	8	KEPPRA TABS	5. Max dose 2400mg	
MC/DEL	GABAPENTIN SOL	MC/DEL	8	KEPPRA SOLN	6. Clinical PA required for appropriate diagnosis	Topamax and Neurontin - Second line therapy for migraine prophylaxis after trial of at least three preferred preventive medications from Group 1 listed on page 2 of the Acute Migraine PA form.
MC/DEL	GABITRIL TABS	MC/DEL	8	KLONOPIN TABS		
MC/DEL	LACOSAMIDE SOL	MC	8	LAMICTAL IR		
MC/DEL	LACOSAMIDE TAB	MC	8	LAMICTAL ODT		
MC	LAMICTAL CHEW	MC/DEL	8	LEVETIRACETAM INJ		All non-preferred meds must be used in specified order.
MC	LAMICTAL XR	MC/DEL	8	LYRICA CR	7. Adjunctive therapy in the treatment of partial-onset seizures in patient's ≥16 years of age with epilepsy.	Please use Drug-Drug Interaction PA form #10400 for this combination.
MC/DEL	LAMOTRIGINE ER ODT	MC/DEL	8	LYRICA SOL ³		
MC/DEL	LAMOTRIGINE IR ²	MC	8	MOTPOLY XR		
MC/DEL	LEVETIRACETAM SOLN	MC/DEL	8	MYSOLINE TABS		
MC/DEL	LEVETIRACETAM TABS	MC	8	ONFI		
MC/DEL	LEVETIRACETAM ER TABS	MC/DEL	8	OXCARBAZEPINE SUS	8. Epidiolex is for the treatment of seizures associated with Lennox-Gastaut syndrome (LGS), Dravet syndrome (DS) or TS (Tuberous Sclerosis Complex) in patients 1 years of age and older.	Epidiolex Criteria for Lennox-Gastaut syndrome (LGS) and Dravet: a trial of two drugs (clobazam, levetiracetam, valproate derivatives, lamotrigine, topiramate, rufinamide, or felbamate).
MC/DEL	LYRICA ³	MC	8	OXTELLAR XR ⁵		Diacomit is for the treatment of seizures associated with Dravet syndrome (DS) in patients 6 months of age and older and weighing 7kg or more There are no clinical data to support the use of Diacomit® as monotherapy in DS.
MC/DEL	NAYZILAM ¹	MC/DEL	8	PHENYTEK CAPS		DDI: Concomitant use of Diacomit® with other CNS depressants, including alcohol, may increase the risk of sedation and somnolence. Concomitant use of strong inducers (CYP1A2, CYP3A4, or CYP2C19 inducers, such as rifampin, phenytoin, phenobarbital, and carbamazepine) should be avoided, or dosage adjustments should be made.
MC/DEL	OXCARBAZEPINE	MC/DEL	8	POTIGA		
MC/DEL	PREGABALIN CAPS	MC/DEL	8	PREGABALIN (ORAL) SOL		
MC/DEL	PHENYTOIN	MC	8	ROWEEPPRA TAB		
MC/DEL	PRIMIDONE TABS	MC	8	SABRIL		
MC/DEL	QUDEXY XR	MC	8	SEZABY	9. For seizures associated with Dravet syndrome in patients 2 years of age and older	DDI: Avoid concomitant use of Nayzilam® with moderate or strong CYP3A inhibitors.
MC/DEL	TEGRETOL SUS	MC	8	SPRITAM		
MC/DEL	TOPIRAMATE	MC	8	SYMPAZAN		
MC/DEL	TOPIRAMATE SPRINKLE IR CAPS	MC/DEL	8	TEGRETOL TAB		
MC/DEL	TRILEPTAL SUS	MC/DEL	8	TIAGABINE	10. Adjunctive therapy 12 and older.	Xcopri criteria: History of trials with at least 4 AEDs (2 generic, 2 branded or Uncontrolled seizures on three AEDs; or Uncontrolled on 2 AEDs given along with VNS. Uncontrolled defined as 3 or more TC seizures per year (increases risk of SUDEP); > 6 disabling seizures per year . Any patient who has gone to the ED 2 or more times in the prior 12 months (who has also tried and failed at least 3 other drugs). Ongoing use requires 50 percent reduction in seizure frequency after three months.
MC/DEL	VALPROIC ACID TABS	MC	8	TOPAMAX		
MC/DEL	VALPROIC ACID SOL	MC/DEL	8	TOPIRAMATE ER CAPS		
MC	VALTOCO ²	MC	8	TOPAMAX SPRINKLE ER CAPS ²		Motpoly XR: pediatric patient weight must be > 50kg and requires multiple preferred medication trials including generic lacosamide
MC/DEL	ZONISAMIDE	MC	8	TOPAMAX SPRINKLE IR CAPS ²		
		MC/DEL	8	TOPIRAMATE SPRINKLE ER CAPS ²		
		MC	8	TROKENDI ^{2,6}		
		MC/DEL	8	VIMPAT ⁴		
		MC/DEL	8	VIMPAT SOL ⁴	11. Initial monotherapy for the treatment of partial-onset or primary generalized tonic-clonic seizures in patients 2 years of age and older. Adjunctive therapy for the treatment of partial-onset seizures, primary generalized tonic-clonic seizures, and seizures associated with Lennox Gastaut syndrome in patients 2 years of age and older. The preventive treatment of migraine in patients 12 years and older. Will require a step though topiramate.	
		MC	8	XCOPRI		
		MC/DEL	8	ZARONTIN SYRP		
		MC/DEL	8	ZARONTIN CAP		
		MC/DEL	8	ZARONTIN SOL		
		MC	8	ZONISADE		
		MC	8	ZTALMY		
		MC/DEL	9	KEPPRA XR		
		MC/DEL	9	NEURONTIN		
		MC/DEL	9	TEGRETOL-XR TB12		

						<p>BIPOLAR DISORDER: STEP ORDER</p> <p>M ~ A</p> <p>4 ~ 4 LAMICTAL</p> <p>4 ~ 4 LITHIUM</p> <p>4 ~ 4 CARBAMAZEPINE</p> <p>4 ~ 4 VALPROATE</p> <p>4 ~ 4 ATYPICAL ANTIPSYCHOTICS EXC. CLOZAPINE</p> <p>5 ~ 5 TRILEPTAL</p> <p>9 ~ 6 TOPAMAX</p> <p>9 ~ 7 KEPPRA TABS</p> <p>9 ~ 8 GABITRIL TABS</p> <p>9 ~ 9 NEURONTIN</p>	<p>SEE ANTICONVULSANT INDICATION CHART AT THE END OF THIS DOCUMENT</p> <p>M= Monotherapy</p> <p>A= Adjunctive</p> <p>9= No Evidence</p> <p>The step orders show the relative strength of evidence for use in bi-polar and will guide prior authorization determinations.</p> <p>Step 4 drugs-no PA required.</p>
						<p>PEDIATRIC BIPOLAR1 DISORDER: STEP ORDER</p> <p>M ~ A</p> <p>(6-18 YEARS WITH OR WITHOUT PSYCHOSIS)</p> <p>4 ~ 4 LITHIUM</p> <p>4 ~ 4 CARBAMAZEPINE</p> <p>4 ~ 4 VALPROATE</p> <p>4 ~ 4</p> <p>ATYPICAL ANTIPSYCHOTICS EXC. CLOZAPINE</p> <p>4 ~ 4 LAMICTAL</p> <p>5 ~ 5 TRILEPTA</p>	<p>Two-step 1 preferred drugs must be tried before Trileptal.</p> <p>The step orders show the relative strength of evidence for use in bi-polar and will guide prior authorization determinations.</p> <p>Step 4 drugs-no PA required.</p>
ANTI-PARKINSON DRUGS							
PARKINSONS - ANTICHOLINERGICS	MC/DEL MC MC/DEL		BENZTROPINE MESYLATE TABS COGENTIN SOLN TRIHENXYPHENIDYL				Use PA Form# 20420
PARKINSONS - ADENOSINE RECEPTOR ANTAGONIST				MC/DEL		NOURIANZ	<p>Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>DDI: Avoid use of Nourianz® with strong CYP3A4 inducers (e.g. carbamazepine, rifampin, phenytoin, St. John's wort).</p>
PARKINSONS - COMT INHIBITORS				MC/DEL MC MC/DEL		COMTAN TABS ONGENTYS TASMAR TABS	Use PA Form# 20420 Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PARKINSONS - SELECTED DOPAMIN AGONISTS	MC/DEL MC/DEL		PRAMIPEXOLE ROPINIROLE	MC/DEL MC MC/DEL MC/DEL	5 8 8 8	MIRAPEX TABS ¹ REQUIP TABS MIRAPEX ER NEUPRO PATCH	Use PA Form# 20420 1. As of 12/08 users of Mirapex will be grandfathered if diagnosis is Parkinsons. Preferred drug must be tried and failed in step-order due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PARKINSONS- MAOIS				MC		XADAGO	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

PARKINSONS - DOPAMINERGICS/CARBI/ LEVO	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL		AMANTADINE HCLCAPS AMANTADINE HCL TABS BROMOCRIPTINE MESYLATE TABS BROMOCRIPTINE MESYLATE CAPS CARBIDOPA/LEVODOPA TABS ³ CARBIDOPA/LEVODOPA ER CARBIDOPA/LEVO/ENTACAPONE TAB LARODOPA TABS SELEGILINE CAPS HCL SELEGILINE TABS HCL	MC/DEL MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC/DEL MC MC MC MC		APOKYN AZILECT ² CARBIDOPA/LEVODOPA RAPDIS ELDEPRYL CAPS GOCOVRI INBRIJA KYNMOBI LODOSYN TABS OSMOLEX ER PARLODEL CAPS PARLODEL TABS RYTARY SINEMET TABS SINEMET TBCR ZELAPAR ¹	Use PA Form# 20420 1. Approvals will require concurrent therapy with Levodopa and failed trials of Selegiline, Comtan, and Stalevo. 2. Approvals will require trials of Carbidopa/Levodopa, Selegiline, Comtan, and Stalevo. 3. Only preferred manufacturer's products will be available without prior authorization.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Inbrija is recommended for the intermittent treatment of OFF episodes in patients with Parkinson's disease treated with carbidopa/levodopa.
PARKINSONS - COMBO.				MC/DEL MC		STALEVO ¹ CARBIDOPA/LEVODOPA/ENTACA ¹	Use PA Form# 20420 1. Clinical PA is required to establish diagnosis and medical necessity.	
MUSCLE RELAXANTS								
MUSCLE RELAXANTS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL		BACLOFEN TABS CHLORZOXAZONE TABS CYCLOBENZAPRINE HCL 5mg & 10mg TABS LIORESAL INTRATHECAL KIT METHOCARBAMOL TABS TIZANIDINE HCL TABS	MC/DEL MC/DEL MC/DEL MC MC MC MC MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	7 8 8 8 8 8 8 8 8 8 8 8 8 9 9 9 9 9	ORPHENADRINE CITRATE CARISOPRODOL 350MG TABS AMRIX DANTRIMUM CAPS FLEQSUVY LIORESAL TABS LORZONE LYVISPAH METAXALONE NORFLEX TBCR OZOBAX ROBAXIN-750 TABS VECUROMIUM INJ ZANAFLEX TABS CARISOPRODOL 250MG TABS CHLORZOXAZONE 250mg TABS SKELAXIN TAB SOMA TABS	Use PA Form# 20420	At least 4 preferred drugs (including tizanidine) must be tried for at least 2 weeks and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an..... acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Elderly patients, over 65, will require written notice of the increased sedative risks and impaired driving. Prior Authorization will not be given for: 1. frequent or persistent early refills of controlled drugs; 2. multiple instances of early refill overrides due to reports of misplacement, stolen, dropped in toilet or sink, distant travel, etc. Non-preferred drugs will not be approved if members circumventing MaineCare prior authorization requirements by paying (prescribers failed to submit prior authorization prior to cash narcotic scripts being filled by member). Non-preferred products must be used in specified step order. Lorzone is non preferred and requires at least 4 preferred drugs (including tizanidine) and step care therapy (orphenadrine), as well as reasons for why chlorzoxazone is not acceptable.
MUSCLE RELAXANT - COMBO.				MC/DEL MC/DEL MC MC/DEL MC/DEL MC		CARISOPRODOL/ASPIRIN TABS CARISOPRODOL/ASPIRIN/CODE NORGESIC TABS ORPHENADRINE COMPOUND ORPHENADRINE/ASA/CAFF ORPHENGESIC	Use PA Form# 20420	Individual components are available with PA described in the section above. 1. frequent or persistent early refills of non-controlled drugs; 2. multiple instances of early refill overrides due to reports of misplacement stolen, dropped in toilet or sink, distant travel, etc.
PARATHYROID HORMONE								
PARATHYROID HORMONE				MC		NATPARA ¹	1. Recommended only for those who cannot be well-controlled on calcium supplements and active forms of vitamin D alone.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

VITAMINS						
VITAMINS	MC MC MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC	CYANOCOBALAMIN SOLN FERIVA CAP FERIVAFA CAP FOLIC ACID TABS MEPHYTON TABS NIACIN NIACOR TABS NICOTINIC ACID SR CPCR PYRIDOXINE HCL TABS TANDEM CAP THIAMINE HCL SOLN VITAMIN B-1 TABS VITAMIN B-12 VITAMIN B-6 TABS VITAMIN C VITAMIN E CAPS VITAMIN E/D-ALPHA CAPS VITAMIN K1 SOLN V-R VITAMIN E CAPS	MC MC MC MC MC MC MC MC MC MC MC MC MC MC MC MC	AQUASOL E SOLN AQUAVIT-E SOLN DHT SOLN FUSION PLUS CAP HEMOCYTE PLU CAP INTEGRA CAP INTEGRA F CAP INTEGRA PLUS CAP NASCOBAL GEL TANDEM PLUS CAP	Use PA Form# 20420 Please refer to OTC list for covered products. Click here for the OTC List	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. Please refer to OTC list for covered products. DDI: B-12 will now be non-preferred and require prior authorization if it is currently being used in combination with either Prevacid, pantoprazole, Prilosec, or any currently non preferred PPI. Preferred products that used to require diag codes still require diag codes unless indicated otherwise.
VITAMIN D's	MC/DEL MC/DEL MC/DEL MC/DEL MC	CALCITRIOL CAPS ¹ ROCALTROL VITAMIN D2 ² VITAMIN D3 ² VITAMIN DROPS PARICALCITOL CAPS	MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC MC	CALCIJEX DOXERCALCIF CAP DOXERCALCIF INJ PARICALCITROL CAP PARICALCITROL INJ HECTOROL (ORAL) HECTOROL (PARENTERAL) RAYALDEE ZEMPLAR INJ ZEMPLAR CAPS	1. Diagnosis of dialysis (renal failure) required. 2. Only specific NDCs available Use PA Form# 20420	Preferred products require dialysis/renal failure diagnosis. Rayaldee requires clinical PA to verify stage 3 or 4 CKD.
EMZYMES						
POMPE DISEASE AGENTS			MC MC MC MC	NEXVIAZYME ¹ LUMIZYME OPFOLDA POMBILITI	1. For patients 1 year of age and older with late-onset Pompe disease (lysosomal acid alpha-glucosidase [GAA] deficiency). Use PA Form# 20420	All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Pombiliti and Opfolda are for the treatment of adult patients with late-onset Pompe disease (lysosomal acid alpha-glucosidase [GAA] deficiency) weighing ≥40kg and who are not improving on their current enzyme replacement therapy (ERT).
MISC MULTI-VITAMINS						
VITAMINS - MISC.	MC MC MC MC MC/DEL MC MC/DEL MC MC	CENTRUM TABS CENTRUM JR/IRON CHEW CENTRUM-LUTEIN TABS CEROVITE ADVANCED FO TABS CHEWABLE MULTIVIT/FL CHEW COD LIVER OIL CAPS COMPLETE NATAL DHA (ORAL) COMBO PKG COMPLETE SENIOR TABS DAILY MULTI VIT/IRON	MC MC/DEL MC MC MC MC MC MC MC	ADEKS ADVANCED NATALCARE TABS AQUADEKS CENTRUM JR/EXTRA C CHEW CENTRUM PERFORMANCE TABS CENTRUM SILVER TABS DALYVITE LIQD EMBREX 600 MISC FERRALET 90	1. Diag codes are no longer required on prenatal vitamins. Please refer to OTC list. Use PA Form# 20420 Click here for the OTC List	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. Please refer to OTC list. Preferred products that used to require diag codes still require diag codes unless indicated otherwise.

MC/DEL	DIALYVITE 1MG	MC	IBERET
MC/DEL	DIALYVITE 800MG	MC	MATERNA TABS
MC/DEL	FULL SPECTRUM B	MC	MAXARON
MC	M.V.I.-12 INJ	MC	MULTIRET FOLIC -500 TBCR
MC	MULTI-VIT/FLUORIDE	MC/DEL	NATAFORT TABS
MC/DEL	NATALCARE RX TABS	MC/DEL	NATALCARE CFE 60 TABS ¹
MC/DEL	NEPHRONEX	MC/DEL	NATALCARE GLOSS TABS ¹
MC/DEL	NIVA-PLUS (ORAL) TABLET	MC	NATALCARE PIC TABS ¹
MC/DEL	ONE DAILY TABS	MC	NATALCARE PIC FORTE TABS ¹
MC/DEL	ONE-DAILY MULTIVITAMINS	MC/DEL	NATALCARE PLUS TABS ¹
MC/DEL	ONE-TABLET-DAILY	MC	NATALCARE THREE TABS ¹
MC/DEL	POLY-VIT/IRON/FLUORID SOLN	MC/DEL	NATACHEW CHEW
MC/DEL	POLY-VITAMIN/FLUORIDE SOLN	MC	NATALFIRST TABS
MC/DEL	POLY-VITAMINS/IRON SOLN	MC	NATATAB RX TABS
MC	PRENATA (ORAL) TAB CHEW	MC/DEL	NEPHPLEX RX TABS
MC/DEL	PRENATAL TABS ¹	MC/DEL	NEPHROCAPS CAPS
MC/DEL	PRENATAL FORMULA 3 TABS ¹	MC/DEL	NEPHRO-VITE TABS
MC/DEL	PRENATAL PLUS TABS ¹	MC	NESTABS RX TABS
MC/DEL	PRENATAL PLUS NF TABS ¹	MC/DEL	NIFEREX
MC	PRENATAL PLUS/27MG IRON ¹	MC/DEL	OCUVITE TABS
MC	PRENATAL PLUS/IRON TABS ¹	MC	POLY-VI-FLOR SOLN
MC	PRENATAL VITAMIN PLUS LOW IRON (ORAL) TAB	MC	POLY-VI-SOL SOLN
MC/DEL	PRENATAL RX/BETA-CAROTENE ¹	MC	POLY-VI-SOL/IRON SOLN
MC/DEL	PREPLUS (ORAL) TABLET	MC	POLY-VITAMIN DROPS SOLN
MC/DEL	RENAL CAPS	MC	PRECARE
MC/DEL	RENAPHRO CAPS	MC	PREFERA OB
MC	STRESS TAB NF TABS	MC	PREMESIS RX TABS
MC	THERAPEUTIC-M TABS	MC	PRENATABS CBF TABS ¹
MC	THERAVITE LIQD	MC	PRENATAL CARE TABS ¹
MC/DEL	TRINATAL RX 1 (ORAL) TABLET	MC	PRENATAL MR 90 TBCR ¹
MC/DEL	TRIVEEN-DUO DHA (ORAL) COMBO. PKG	MC/DEL	PRENATAL MTR/SELENIUM TABS ¹
MC/DEL	TRI-VITAMIN/FLUORIDE SOLN	MC	PRENATAL OPTIMA ADVANCE TABS ¹
MC	VITA CON FORTE CAPS	MC	PRENATAL PC 40 TABS ¹
MC	VITAPLEX PLUS TABS	MC/DEL	PRENATAL RX TABS ¹
		MC	PRENATE ¹
		MC	PRENATE ELITE ¹
		MC	PRIMACARE MISC
		MC	PROTEGRA CAPS
		MC	STUARTNATAL PLUS 3 TABS ¹
		MC	TRI-VI-SOL SOLN
		MC	TRI-VI-SOL/IRON SOLN
		MC/DEL	ULTRA NATALCARE TABS
		MC	ULTRA-NATAL TABS ¹
		MC	VICON FORTE CAPS
		MC	VINATAL FORTE TABS ¹
		MC	VINATE ¹
		MC/DEL	VINATE ADVANCED TABS ¹

MISCELLANEOUS MINERALS

MINERALS	MC	CALCARB	MC	ANEMAGEN	Use PA Form# 20420 Please refer to OTC list. Click here for the OTC List	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. DDI: Fe salts will now be non-preferred and require prior authorization if it is currently being used in combination with either Prevacid, pantoprazole, Prilosec, or any currently non preferred PPI. Please refer to OTC list.
	MC	CALCI-MIX CAPSULE CAPS	MC	CALCET TABS		
	MC	CALCIQUID SYRP	MC/DEL	CALCIUM 600-D TABS		
	MC	CALCITRATE/VITAMIN D TABS	MC	CALCIUM/VITAMIN D TABS		
	MC/DEL	CALCIUM	MC	CALTRATE 600 PLUS/VIT D TABS		
	MC/DEL	CALCIUM CARBONATE	MC	CALTRATE PLUS TABS		
	MC/DEL	CALCIUM CITRATE TABS	MC	CHROMAGEN		
	MC/DEL	CALCIUM GLUCONATE TABS	MC	CITRACAL PLUS TABS		
	MC/DEL	CALCIUM LACTATE TABS	MC	CONTRIN CAPS		

MC	CALCIUM/MAGNESIUM TABS	MC	FEOGEN FORTE CAPS
MC/DEL	CALCIUM/VITAMIN D TABS	MC	FEROCON CAPS
MC	CALTRATE 600 TABS	MC/DEL	FERREX 150 CAPS
MC/DEL	CHEWABLE CALCIUM CHEW	MC	FERRO-SEQUELS TBCR
MC	CITRACAL TABS	MC	FE-TINIC CAPS
MC	CITRACAL + D TABS	MC	FE-TINIC 150 FORTE CAPS
MC	CITRUS CALCIUM TABS	MC/DEL	FLUOR-A-DAY SOLN
MC	CITRUS CALCIUM 1500 + D TABS	MC	HEMOCYTE TABS
MC	EFFERVESCENT POTASSIUM TBEF	MC/DEL	K-DUR TBCR
MC/DEL	FEOSTAT CHEW	MC	KLOR-CON PACK
MC	FERATAB TABS	MC	K-LYTE
MC/DEL	FER-GEN-SOL SOLN	MC/DEL	K-PHOS TABS NEUTRAL
MC	FER-IRON SOLN	MC	K-TABS TBCR
MC	FERRONATE TABS	MC	K-VESCENT PACK
MC/DEL	FERROUS SULFATE	MC	MICRO-K 10 MEG CPCR
MC/DEL	FLUOR-A-DAY CHEW	MC	NU-IRON 150 CAPS
MC	FLUORIDE CHEW	MC/DEL	OYSTER SHELL CALCIUM/VITA TABS
MC	FLUORIDE SODIUM CHEW	MC/DEL	POLY-IRON 150 CAPS
MC	FLUORITAB CHEW	MC/DEL	POLYSACCHARIDE IRON CAPS
MC	HM CALCIUM TABS	MC/DEL	POTASSIUM BICARB/CHLORIDE
MC	K+ POTASSIUM PACK	MC/DEL	POTASSIUM CHLORIDE 10MEQ CAPS
MC	KAON ELIX	MC/DEL	POTASSIUM CHLORIDE 8MEQ CAPS
MC	KAON-CL-10 TBCR	MC	TUMS 500 CHEW
MC	KCL 0.075%/D5W/NACL 0.2% SOLN	MC	VIACTIV CHEW
MC	K-EFFERVESCENT TBEF		
MC	KLOR-CON		
MC	KLOTRIX TBCR		
MC/DEL	K-PHOS TABS		
MC/DEL	K-VESCENT TBEF		
MC/DEL	LURIDE CHEW		
MC/DEL	MAGNESIUM GLUCONATE TABS		
MC/DEL	MAGNESIUM SULFATE SOLN		
MC	MAGTABS		
MC	MICRO-K 8 MEG		
MC/DEL	OS-CAL TABS		
MC/DEL	OS-CAL 500 + D TABS		
MC/DEL	OYSCO		
MC/DEL	OYST-CAL TABS		
MC/DEL	OYST-CAL D TABS		
MC/DEL	OYST-CAL/VITAMIN D TABS		
MC/DEL	OYSTER CALCIUM TABS		
MC/DEL	OYSTER SHELL		
MC	PHARMA FLUR		
MC/DEL	PHOSPHA 250 NEUTRAL TABS		
MC	POTASSIUM BICARBONATE TBEF		
MC/DEL	POTASSIUM CHLORIDE 8MEQ		
MC	POTASSIUM EFFERVESCENT		
MC/DEL	SELENIUM TABS		
MC	SLOW-MAG TBCR		
MC/DEL	SODIUM FLUORIDE		
MC	V-R CALCIUM		
MC	V-R OYSTER SHELL CALCIUM		
MC	ZINC SULFATE CAPS		

Preferred products that used to require diag codes still require diag codes unless indicated otherwise.

PHENYLKETONURIA (PKU) TREATMENT AGENTS- INJECTABLES				MC		PALYNZIQ ¹	1. For the treatment of patients ≥ 18 years of age. Use PA Form# 20420	Palynziq is not to be used in combination with Kuvan
PHENYLKETONURIA (PKU) TREATMENT AGENTS- ORAL				MC		KUVAN	Use PA Form# 20420	
MISC. ELECTROLYTES/NUTRITIONALS								
ELECTROLYTES/ NUTRITIONALS	MC MC MC		INTRALIPID EMUL ¹ P.T.E. -5 SOLN ¹ SEA-OMEGA CAPS ¹	MC MC		BOOST ¹ CASEC POWD ¹ CHOICE DM LIQD ¹ DELIVER 2.0 LIQD ¹ DOJOLVI ENFAMIL ¹ ENSURE ¹ GLUCERNA ¹ ISOCAL LIQD ¹ KINDERCAL TF LIQD ¹ KINDERCAL TF/FIBER LIQD ¹ L-CARNITINE CAPS ¹ LIPISORB LIQD ¹ LOVAZA ^{1,2} MODULEN IBD POWD ¹ NUTRAMIGEN POWD ¹ NUTREN ¹ NUTRITIONAL SUPPLEMENT LIQD ¹ NUTRIVENT 1.5 LIQD ¹ PEPTAMEN ¹ PHENYLADE ¹ PHENYL-FREE ¹ PKU 3 POWD ¹ PREGESTIMIL POWD ¹ PROBALANCE LIQD ¹ PROSOBEE ¹ SCANDISHAKE PACK ¹ VASCEPA	1. This list of nutritionals is incomplete. All nutritionals still require a PA except for the miscellaneous products listed as preferred. SGA form required for nutritionals unless member has a G/I tube. 2. Formerly known as Omacor. Use PA Form# 20420 & SGA Form	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. Medical foods are not to be authorized solely for the purpose of enhancing nutrient intake or managing body weight if the participant is able to eat conventional foods adequately. Medical foods may be approved if the member has a medical condition which precludes or restricts the use of conventional foods and necessitates the use of a formula. Concurrent Stimulant therapy is not an acceptable medical reason/condition for use of medical foods for enhancing nutrient intake or managing body weight. For children under the age of 5, MaineCare will not provide milk- or soy-based standard infant formulas. Regular formulas may be sought through your nearest WIC office. MaineCare will continue to cover medical food for all participants in MaineCare when medical necessity is met. Vascepa requires adjunct therapy for specific indication to reduce TG in those with severe hypertriglyceridemia (500mg per deciliter or more). Proper indication per lab values is required before approval
ERYTHROPOEITINS	MC MC MC		EPOGEN SOLN MIRCERA SYRINGE RETACRIT	MC MC	8 8	ARANESP SOLN ¹ PROCRT SOLN ¹	Use PA Form# 10520 1. Clinical PA is required to establish medical necessity and that appropriate lab monitoring is being done.	Non-Preferred drugs must be tried and failed in step-order, due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please see the EPO PA form for other approval and renewal criteria.
GRANULOCYTE CSF								
GRANULOCYTE CSF	MC MC MC/DEL MC/DEL		NEUPOGEN SYRINGE NEUPOGEN VIAL NYVEPRIA SYRINGE ZIEXTENZO	MC MC MC MC MC/DEL MC MC MC/DEL MC	8 8 8 8 8 8 8 8 8 8 9	FULPHILA FYLNETRA GRANIX SYRINGE GRANIX VIAL LEUKINE NIVESTYM ROLVEDON STIMUFEND ZARXIO NEULASTA ¹	1. Must be used in specified step order. Use PA Form# 20520	See approval criteria detailed on Granulocyte Colony Stimulating Factor PA form.

GAUCHER DISEASE

GAUCHER DISEASE				MC		CERDELGA ¹	1. Clinical PA for indication required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Exceeding days supply limits for LMWH class requires PA.
				MC		YARGESA ¹		Yargesa: As monotherapy for the treatment of adult patients with mild to moderate type 1 Gaucher disease for whom enzyme replacement therapy is not a therapeutic option (e.g., due to allergy, hypersensitivity, or poor venous access).
							Use PA Form# 20420	

ANTICOAGULANTS / PLATELET AGENTS

ANTICOAGULANTS	MC MC/DEL MC MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		COUMADIN TABS ENOXAPARIN ¹ ELIQUIS ELIQUIS STARTER PACK HEPARIN SODIUM/NACL 0.9% SOLN HEP-LOCK SOLN INNOHEP HEPARIN LOCK SOLN HEPARIN LOCK FLUSH SOLN HEPARIN SODIUM SOLN HEPARIN SODIUM LOCK FLUSH SOLN PRADAXA JANTOVEN WARFARIN SODIUM TABS XARELTO XARELTO STARTER PACK	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL		ARIXTRA SOLN FONDAPARINUX FRAGMIN INJ FRAGMIN VIAL LOVENOX SOLN LOVENOX 300 ² LOVENOX SUBQ SYRINGE PRADAXA ORAL PELLETS ⁴ IPRIVASK SAVAYSAS ³	1. Enoxaparin therapy durations greater than 7 days every 30 days require PA 2. Use other strengths available to obtain desired dose. 3. Diagnosis required 4. For the treatment of patients aged 3 months to less than 12 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Exceeding days supply limits for LMWH class requires PA. DDI: Warfarin will require prior authorization if being used in combination with fluconazole, miconazole, or voriconazole. DDI: Warfarin will require prior authorization if being used in conjunction with Gemfibrozil or Fenofibrate. DDI: Rifampin will require prior authorization if being used in combination with Savaysa
							Use PA form# 20420	

ANTHEMOPHILIC AGENTS	MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC/DEL MC MC MC/DEL MC MC/DEL MC MC/DEL		AFSTYLA ALPHANATE ALPHANINE SD ALPROLIX VIAL BEBULIN VIAL BENEFIX SOLR HELIXATE FS KIT HEMLIBRA HEMOPIL - M HUMATE-P SOLR IXINITY VIAL JIVI ³ KOATE-DVI KONYNE - 80 KOVALTRY MONARC - M MONOCLATE - P MONONINE NOVOEIGHT NOVOSEVEN SOLR NUWIQ PROFILNINE RECOMBINATE SOLR REFACTO RIXUBIS VIAL WILATE INJ XYNTHA	MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC MC MC MC MC MC MC MC MC MC/DEL MC MC/DEL		ADYNOVATE VIAL ADVATE ^{1,2,5} ALTUVIIIIO ⁴ ESPEROCT ELOCTATE HEMGENIX IDELVION KOGENATE FS ⁵ REBINYN RECOMBINATE VIAL ⁵ ROCTAVIAN ⁴ SEVENFACT	1. Only if other products unavailable. 2. Advate may be available with PA in cases of large volume dosing in patients with poor venous access. 3. Not indicated for use in children <12 years of age due to greater risk for hypersensitivity reactions and is not indicated for use in previously untreated patients. 4. Clinical PA required for appropriate diagnosis. 5. Established users will be grandfathered	Non-preferred will only be approved if other preferred products are unavailable. Hemgenix® is an adeno-associated viral vector-based gene therapy for IV infusion after dilution. For treatment of adults with Hemophilia B (congenital Factor IX deficiency) who: Currently use Factor IX prophylaxis therapy, or have current or historical life-threatening hemorrhage, or Have repeated, serious spontaneous bleeding episodes. Altuviiiio is a von Willebrand Factor (VWF) independent recombinant DNA-derived, Factor VIII concentrate indicated for use in adults and children with hemophilia A (congenital factor VIII deficiency) for: Routine prophylaxis to reduce the frequency of bleeding episodes, On-demand treatment and control of bleeding episodes, Perioperative management of bleeding. Roctavian: For the treatment of adults with severe hemophilia A (congenital factor VIII deficiency with factor VIII activity <1 IU/dL) without antibodies to adeno-associated virus serotype Inclusion: Severe factor VIII deficiency (less than 1% native factor VIII). Exclusion Criteria: Antibodies to the virus AAV5 Factor VIII inhibitors (or history of) Known significant fibrosis of cirrhosis of the liver, or unexplained elevated LFTs History of inadequate compliance with prophylaxis, or regular bleeds despite adequate prophylaxis Conditions in which high-dose steroids are contraindicated. -Inability to abstain from alcohol for one year Plan to impregnate a partner within 6 months of infusion -Hypersensitivity to mannitol -Active infections, either acute or uncontrolled chronic
----------------------	--	--	---	--	--	---	--	---

							Use PA Form# 20420	-HIV infection (limited information on use in this population)
PLATELET AGGREGATION INHIBITORS	MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL		ASPIRIN ASPIRIN-DIPYRIDAMOLE ER CPMP 12HR BRILINTA ¹ DIPYRIDAMOLE TABS CLOPIDOGREL 75MG PRASUGREL HCL TAB	MC/DEL MC MC MC/DEL MC/DEL MC/DEL	7 8 8 8 8 8	TICLOPIDINE HCL TABS DURLAZA EFFIENT PERSANTINE TABS PLAVIX TABS ZONTIVITY	Use PA Form# 20715 for Plavix, Effent & Brilinta Use PA form# 20420 for other requests 1. Dosing limits apply, please see dose consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. A special PA may be obtained at the pharmacy for members scheduled for "stent" placement or have had placement if in the last 12months. Please indicate on prescription date of stent placement. DDI: Plavix will require prior authorization if being used in combination with omeprazole, esomeprazole, cimetidine, fluconazole, ketoconazole, intelence, fluoxetine, ticlopidine, and fluvoxamine. DDI: exists for using maintenance ASA dose >100mg, as it reduces the effectiveness of Brilinta Brilinta- Concomitant use with strong CYP3A4 inhibitors should be avoided (including ketoconazole, itraconazole, atazanavir, and telithromycin). Doses of simvastatin and lovastatin >40mg should be avoided.
PLATELET AGGR. INHIBITORS / COMBO'S - MISC.	MC/DEL MC/DEL		CILOSTAZOL PENTOXIFYLLINE ER TBCR	MC/DEL MC/DEL MC/DEL MC MC		AGRYLIN CAPS ANAGRELIDE CAPS PLETAL TABS TRENTAL TBCR YOSPRALA	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
HEMATOLOGICALS								
MONOCLONAL ANTIBODY				MC/DEL MC/DEL MC MC/DEL MC		EMPAVELI ENSPRYNG GAMIFANT SOLIRIS ULTOMIRIS UPLIZNA	Use PA Form# 20420	A diagnosis of Paroxysmal nocturnal hemoglobinuria (PNH) using the HAM test or flow cytometry is required. In addition, the patient must show evidence of having received a meningitis vaccine at least 2 weeks prior to the start of therapy. Gamifant is recommended for the treatment of adult and pediatric (newborn and older) patients with primary hemophagocytic lymphohistiocytosis (HLH) with refractory, recurrent, or progressive disease or intolerance with conventional HLH therapy. Ultomiris is recommended for the treatment of adults with paroxysmal nocturnal hemoglobinuria (PNH).
IMMUNE GLOBULIN	MC MC/DEL MC MC MC/DEL MC/DEL MC		BIVIGAM ¹ CUTAQUIG ¹ GAMUNEX-C GAMMAGARD S-D ¹ HIZENTRA ¹ PANZYGA ¹ PRIVIGEN ¹	MC MC/DEL MC MC/DEL MC MC/DEL		ASCENIV ² CUVITRU GAMMAPLEX INJ HYQVIA OCTAGAM INJ ¹ XEMBIFY	Use PA Form# 20420 1. Clinical PA required 2. For the treatment of patients between 12 to 17 years of age.	Cutaquig is indicated as replacement therapy for primary humoral immunodeficiency (PI) in adults. Xembify is indicated for treatment of primary humoral immunodeficiency (PI) in patients 2 years of age and older. Asceniv indicated for the treatment of primary humoral immunodeficiency (PI) in adults and adolescents (12 to 17 years of age). PI includes but is not limited to the humoral immune defect in congenital agammaglobulinemia, common variable immunodeficiency (CVID), X-linked agammaglobulinemia, Wiskott-Aldrich syndrome, and severe combined immunodeficiencies (SCID).
HEREDITARY ANGIOEDEMA	MC MC MC MC/DEL MC/DEL MC MC/DEL		PROPHYLAXIS			PROPHYHLAXIS	1. Clinical PA is required to establish diagnosis and medical necessity. 2. For the treatment of patients ≥ 12 years of age.	Haegarda is indicated for routine prophylaxis to prevent Hereditary Angioedema (HAE) attacks in adolescent and adult patients
			TREATMENT			TREATMENT		
			CINRYZE ¹ HAEGARDA ¹ ORLADEYO ^{1,2} TAKHZYRO ¹ BERINERT KIT ¹ FIRAZYR ¹ RUCONEST VIAL ¹			KALBITOR VIAL		
HEMATOLOGICAL AGENTS- THROMBOPOIETIN RECEPTOR AGONISTS	MC MC		PROMACTA ¹ NPLATE ¹	MC/DEL MC/DEL		DOPTELET MULPLETA	Use PA Form# 20420 1. Clinical PA required. Must see prior trial with insufficient response to corticosteroids and immunoglobulins.	Doptelet and Mulpelta: For the treatment of thrombocytopenia in adults with chronic liver disease who are scheduled to undergo a procedure.

HEMATOLOGICAL AGENTS-IgAN				MC/DEL MC		FILSPARI ¹ TARPEYO	Use PA Form# 20420 1. PA required to confirm FDA approved indication.	All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists
ANEMIA- BETA THALASSEMIA				MC MC		REBLOZYL ZYNTEGLO	Use PA Form# 20420	Reblozyl is indicated for the the treatment of anemia in adult patients with beta thalassemia who require regular red blood cell (RBC) transfusion. It is not indicated for use as a substitute for RBC transfusions in patients who require immediate correction of anemia. Zynteglo is indicated for the treatment of adult and pediatric patients with β-thalassemia who require regular red blood cell (RBC) transfusions.
HEMATOLOGIC DISORDER TREATMENT AGENTS				MC/DEL MC		CABLIVI TAVALISSE	Use PA Form# 20420	Tavalisse is recommended for patients at risk of bleeding when one line of therapy (steroids, IVIG, splenectomy) has failed. Cablivi is recommended for the treatment of adult patients with acquired thrombotic thrombocytopenic purpura (aTTP), in combination with plasma exchange and immunosuppressive therapy.
COMPLEMENT RECEPTOR ANTAGONIST				MC		TAVNEOS	Use PA Form# 20420	
HEMOSTATIC								
HEMOSTATIC	MC/DEL MC		AMICAR AMINOCAPROIC ACID	MC MC		FIBRYGA RIASTAP	Use PA Form# 20420	Fibryga and Riastap are indicated for the treatment of acute bleeding episodes in adults and adolescents with congenital fibrinogen deficiency, including afibrinogenemia and hypofibrinogenemia. Fibryga® is not indicated for dysfibrinogenemia.
ACUTE HEPATIC PORPHYRIA (AHP)								
ACUTE HEPATIC PORPHYRIA (AHP)				MC		GIVLAARI	Use PA Form# 20420	Givlaari is indicated for the treatment of adults with acute hepatic porphyria (AHP).
PYRUVATE KINASE DEFICIENCY AGENTS								
PYRUVATE KINASE DEFICIENCY AGENTS				MC		PYRUKYND ¹	Use PA Form# 20420 1.PA required to confirm FDA approved indication.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s)
OP. - ANTIBIOTICS								
OP. - ANTIBIOTICS	MC MC MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC/DEL		AK-SPORE OINT BACITRACIN/NEOMYCIN/POLYM BACITRACIN/POLYMYXIN B OINT CHLOROPTIC SOLN ERYTHROMYCIN OINT NEOSPORIN SOLN POLYSPORIN TRIMETHOPRIM SULFATE/POLY TOBRAMYCIN SULFATE SOLN	MC MC MC MC MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC		AK-POLY-BAC OINT AK-SULF OINT AK-TOB SOLN AZASITE BACITRACIN OINT BLEPH-10 SOLN GATIFLOXACIN DROPS GENTAMICIN SULFATE GENTAK ILOTYCIN OINT LEVOFLOXACIN DROPS NEOMYCIN/BACI/POLYM OINT NEOMYCIN/POLYMYXIN/GRAMIC NEOSPORIN OINT OCUSULF-10 SOLN OCUTRICIN SOLN POLYTRIM DROPS SULFACETAMIDE SODIUM DROPS SULFACETAMIDE SODIUM OINT TERAK OINT	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. -ANTI-PARASITIC				MC		XDEMYV ¹	Use PA Form# 20420 1. For the treatment of Demodex biopharitis.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

OP. - RHO KINASE INHIBITORS	MC		RHOPRESSA					on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s)
							Use PA Form# 20420	
OP. - QUINOLONES	MC/DEL MC/DEL MC/DEL MC/DEL		CILOXAN OINT CIPROFLOXACIN SOL 0.3% OFLOXACIN QUIXIN SOLN	MC/DEL MC/DEL MC		BESIVANCE CILOXAN SOLN OCUFLOX SOLN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - QUINOLONES-4TH GENERATION	MC/DEL		MOXIFLOXACIN 0.5% SOLN (Generic Vigamox)	MC		ZYMAXID	Use PA Form# 20420	
OP. - ARTIFICIAL TEARS AND LUBRICANTS	MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC MC MC MC		ARTIFICIAL TEARS OINT ARTIFICIAL TEARS SOLN CELLUVISC SOLN EYE LUBRICANT OINT GENTEAL LIQUITEARS SOLN MAJOR TEARS SOLN PURALUBE OINT PURALUBE TEARS SOLN REFRESH SOLN OP REFRESH PLUS SOLN ¹ REFRESH PM OINT	MC/DEL MC MC MC MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC MC/DEL MC		ARTIFICIAL TEARS SOLN OP BION TEARS SOLN DRY EYES OINT DURATEARS OINT HYPO TEARS ISOPTO TEARS SOLN LACRI-LUBE LUBRIFRESH P.M. OINT MURINE SOLN MUROCEL SOLN NATURE'S TEARS SOLN REFRESH SOLN REFRESH TEARS SOLN ¹ TEARGEN SOLN TEARISOL SOLN TEARS NATURALE TEARS PURE SOLN TEARS RENEWED OINT THERATEARS SOLN V-R ARTIFICIAL TEARS SOLN	Use PA Form# 20420 1. Dosing limits apply, please see dose consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - BETA - BLOCKERS	MC/DEL MC/DEL MC/DEL MC/DEL		BETOPTIC-S SUSP CARTEOLOL HCL SOLN LEVOBUNOLOL HCL SOLN METIPRANOLOL SOLN	MC MC/DEL MC MC/DEL MC MC/DEL MC/DEL		BETAGAN SOLN BETAXOLOL HCL SOLN ISTALOL OCUPRESS SOLN OPTIPRANOLOL SOLN TIMOPTIC SOLN TIMOLOL DROP TIMOLOL SOL-GEL TIMOPTIC-XE SOLG	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - ANTI-INFLAMMATORY / STEROIDS OPTH.	MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL		AK-SPORE HC OINT ALREX SUSP DEXAMETH SOD PHOS SOLN FLAREX SUSP FLUOROMETHOLONE SUSP FML DROPS SUSP 1% FML FORTE SUSP FML S.O.P. OINT LOTEMAX OINT	MC MC MC MC MC MC MC MC/DEL MC/DEL		AK-TROL SUSP BAC/POLY/NEOMY/HC OINT BLEPHAMIDE S.O.P. OINT BLEPHAMIDE SUSP BROMDAY EFLONE SUSP FLUOR-OP SUSP ILUVIEN IMPLANT INVELTYS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

	MC/DEL MC/DEL MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL	LOTEMAX SUSP LOTEMAX SM DROPS GEL 0.38% NEO/POLY/DEXAMETH OINT NEO/POLY/DEXAMETH SUSP PRED-G SUSP PRED FORTE SUSP 1% PRED MILD SUSP PREDNISOLONE TOBRADEX OINT TOBRADEX SUSP TOBEX OINT SULFACETAMIDE/PREDNISOLONE ZYLET SUSP	MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC	LOTEMAX GEL MAXITROL OPTH OINT 0.1% NEO/POLY/BAC/HC OINT NEOM/POLY/DEX OPTH OINT 0.1% OMNIPRED DROPS SUSP OZURDEX PRED-G S.O.P. OINT PREDNISOLONE SODIUM PHOSPHATE SOL RETISERT IMPLANT SULFACET SOD/PRED SOLN TRIESENCE VIAL TOBRADEX ST TOBRAMYCIN SUSP DEXAMETHASONE VASOCIDIN SOLN VEXOL SUSP XIPERE			
OP. - PROSTAGLANDINS	MC/DEL MC MC/DEL MC/DEL	LATANOPROST SOL 0.005% LUMIGAN SOLN ROCKLATAN TRAVATAN-Z	MC/DEL MC/DEL MC MC	7 8 8 8	ZIOPTAN BIMATOPROST 0.03% DROPS DURYSTA IYUZEH RESCULA ^{1,2,3}	1. All preferreds must be tried. 2. Dosing limits apply, please see dosing consolidation list. 3. Clinical PA is required to establish diagnosis and medical necessity. Use PA Form# 20420	Preferred drugs must be tried and failed, in step-order, due to lack of efficacy (failure to reach target IOP reduction) or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - CYCLOPLEGICS	MC MC/DEL MC/DEL MC/DEL	AK-PENTOLATE SOLN ATROPINE SULFATE CYCLOPENTOLATE HCL SOLN ISOPTO HYOSCINE SOLN	MC/DEL MC MC/DEL MC		CYCLOGYL SOLN ISOPTO ATROPINE SOLN ISOPTO HOMATROPINE SOLN MUROCOLL-2 SOLN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - MIOTICS - DIRECT ACTING	MC/DEL MC MC MC/DEL MC/DEL	ISOPTO CARBACHOL SOLN ISOPTO CARPINE SOLN PILOCAR SOLN PILOCARPINE HCL SOLN PILOPINE HS GEL				Use PA Form# 20420	
OP. - SELECTIVE ALPHA ADRENERGIC AGONISTS	MC MC MC MC/DEL MC/DEL	ALPHAGAN SOLN ALPHAGAN P 0.1% SOLN ALPHAGAN P 0.15% SOLN BRIMONIDINE DROPS 0.2 % SIMBRINZA	MC/DEL MC/DEL		BRIMONIDINE TARTRATE DROPS 0.15 % IOPIDINE SOLN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - ANTI-ALLERGICS	MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL	AZELASTINE HCL DROPS BEPREVE CROMOLYN SODIUM DROPS KETOTIFEN FUMARATE DROPS LASTACAFT OLOPATADINE HCL 0.1% OLOPATADINE HCL 0.2% ZADITOR SOLN	MC MC/DEL MC/DEL MC MC/DEL MC MC/DEL	8 8 8 8 8 8 9	ALOCRIOL SOLN ALOMIDE SOLN EMADINE SOLN OPTICROM SOLN PATANOL SOLN ZERVIAE EPINASTINE	Use PA Form# 20420	All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. ANTI-ALLERGICS- MASTCELL STABILIZER CLASS			MC/DEL		ALAMAST SOLN	Use PA Form# 20420	

				MC/DEL BENZAMYCIN GEL MC/DEL BENZAMYCINPAK PACK MC BENZEFOAM MC BENZOYL PEROXIDE MC BREVOXYL MC/DEL CLEOCIN-T ² MC CLINAC BPO GEL MC CLINDAGEL GEL MC/DEL CLINDAMYCIN PHOSPHATE CREAM ² MC CLINDETS SWAB MC DESQUAM-E GEL MC DESQUAM-X MC DIFFERIN 0.3% GEL MC DIFFERIN MC EMGEL GEL MC EPIDUO MC EPSOLAY MC ERYCETTE PADS MC FINEVIN CREA MC/DEL KLARON LOTN MC METROCREAM CREA ² MC METROGEL GEL ² MC METROLOTION LOTN ² MC NEOBENZ MICRO MC/DEL NORITATE CREA MC ONEXTON ⁵ MC/DEL PLIXDA MC RETIN-A GEL ² MC RETIN-A CREA ² MC RETIN-A MICRO GEL MC RHOFADE MC/DEL SODIUM SULFACET/SULF LOTN MC SOOLANTRA ⁴ MC/DEL TRIAZ MC TWYNEO MC VELTIN MC WINLEVI ⁵ MC ZENCIA WASH MC ZETACET MC/DEL ZIANA MC ZILXI	please see dosing consolidation list. 5. Not approved for use in children <12 years of age 6. For the treatment of patients ≥ 9 years of age. Use PA Form# 10220 for Brand Name requests Use PA Form# 20420 for all other requests		
TOPICAL- ATOPIC DERMATITIS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC	1 1 1 2 2 2	ELIDEL CREA PIMECROLIMUS CRE (AUTH GENERIC LABELER 68682 Oceanside Pharmaceuticals) PROTOPIC OINT TACROLIMUS OINT ADBRY ^{2,4} DUPIXENT ^{1,2,4} EUCRISA ^{2,4}	MC/DEL MC	CIBINQO OPZELURA ³	1. Avoid live vaccines if treated with Dupixent 2. Clinical PA required. 3. For the treatment of patients ≥ 12 years of age. 4. Preferred after a trial and failure of TCs and TCIs. Use PA Form# 20420	Preferred drugs also indicated for this condition, including topical steroids, cyclosporin AND calcineurin inhibitors must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Note: If unable to use TCIs then a trial of Eucrisa could be recommended before Dupixent.

	MC		MISCELLANEOUS PROCTO-KIT CREA 1%	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC	DIPROLENE IMPEKLO ⁴ LEXETTE OLUX FOAM PSORCON PSORCON E SERNIVO SPRAY ² TEMOVATE ULTRAVATE		
TOPICAL - STEROID LOCAL ANESTHETICS				MC	EPIFOAM FOAM	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - STEROID COMBINATIONS	MC		DERMA-SMOOTHIE-FS SCALP	MC	CARMOL-HC CREA	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - EMOLLIENTS	MC/DEL MC MC		AMMONIUM LACTATE CREA ¹ AMMONIUM LACTATE LOTN 12% ¹ VITAMIN A & D MEDICATED OINT	MC MC MC MC MC	LAC-HYDRIN CREA ¹ LAC-HYDRIN LOTN 12% MEDERMA GEL MIMYX RENOVA CREA	Use PA Form# 20420 1. Dosing limits still apply. Please see dose consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - ENZYMES / KERATOLYTICS / UREA				MC MC MC	CARMOL 40 CREA SALEX CREA SALEX LOTN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Ziox, Panafil and Papain products have been removed from the PDL due to FDA safety concerns regarding drugs containing Papain.
TOPICAL - GENITAL WARTS	MC/DEL		IMIQUIMOD 5% ²	MC/DEL MC/DEL MC/DEL MC MC MC MC	5 8 8 8 8 8 8	PODOFILOX SOLN CONDYLOX ¹ ALDARA ¹ PICATO VEREGEN ¹ ZYCLARA ¹	Use PA Form# 20420 1. Non-preferred products must be used in specified order. 2. Dosing limits still apply. Please see dose consolidation list.
TOPICAL - LOCAL ANESTHETICS	MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL		AF CAPSICUM OLEORESIN CREA CAPSAICIN CREA CAPSAICIN PATCH DIBUCAINE OINT ELA-MAX ¹ LIDOCAINE/PRILOCAINE CREA ¹ LIDOCAINE CREAM LIDOCAINE GEL LIDOCAINE PTCH 5%	MC/DEL MC/DEL MC MC MC MC MC MC/DEL	EMLA PADS EMLA CREA LIDA MANTLE CREA LIDODERM PTCH PONTOCAINE SOLN SYNERA ZOSTRIX ZTLIDO ²	Use PA Form# 20420 1. Lidocaine/Prilocaine cream and Ela-Max products require PA for users over 18 years of age. 2. Dosing limits still apply. Please see dose consolidation list. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - DEPIGMENTING AGENTS				MC MC MC MC/DEL MC/DEL MC MC MC	8 8 8 8 8 8 8 9	ALUSTRA CREA EPIQUIN MICRO GLYQUIN CREA HYDROQUINONE CREA HYDROQUINONE/SUNSCREENS SOLAQUIN FORTE CREA TRI-LUMA CREA ELDOQUIN	Use PA Form# 20420 As per Medicaid Policy, cosmetic drugs are not covered. Non-cosmetic clinical applications will be considered by prior authorization on a case by case basis.
TOPICAL - SCABICIDES AND PEDICULICIDES	MC/DEL MC MC/DEL MC/DEL		ACTICIN CREA LICE KILLING SHAM LICE TREATMENT CREME RINS LIQD PERMETHRIN LOTN	MC MC MC/DEL MC	ELIMITE CREA EURAX LINDANE MALATHION	Use PA Form# 20420 1. Dosing limits apply, please refer to dosage consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

	MC		NATROBA ¹	MC MC/DEL		OVIDE LOTN SPINOSAD SUSP		
TOPICAL - WOUND / DECUBITUS CARE				MC MC		REGRANEX GEL VYJUVEK	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Regranex will be approved for diabetic patients in good control (HgbA1c <8), who are not smoking, with a stage III or IV WOCN AND NPUAP lower extremity diabetic ulcer and with an adequate blood supply (TcP 02 >30, ABI >0.7 or ASP > 70), and where the underlying cause has been corrected. The wound must be free of infection and have been previously treated with preferred standard therapies for at least 2 months. Maximum approval for 20 weeks. Vyjuvek: For the treatment of wounds in patients 6 months of age and older with dystrophic epidermolysis bullosa (DEB) with mutation(s) in the collagen type VII alpha 1 chain (COL7A1) gene. Accuzyme and Ethezyme products have been removed from the PDL due to FDA concerns regarding drugs containing Papain.
TOPICAL - ASTRINGENTS / PROTECTANTS	MC		XERAC AC SOLN	MC MC MC MC		LOWILA BAR MOISTURIN DRY SKIN CREA PROSHIELD PLUS SKIN PROTE CREA SURGILUBE GEL	Use PA Form# 20420 1. Dosing limits apply, please refer to dosage consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - ANTISEPTICS / DISINFECTANTS	MC/DEL		POVIDONE-IODINE SOLN	MC MC MC MC		BETADINE OINT FORMALYDE-10 AERS IODOSORB LAZERFORMALYDE SOLUTION SOLN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MISCELLANEOUS EYE								
OP. - EYE	MC MC MC MC MC MC/DEL		AK-DILATE SOLN EYE WASH SOLN NAPHAZOLINE HCL SOLN PHENYLEPHRINE HCL SOLN PONTOCAINE SOLN SODIUM CHLORIDE	MC MC/DEL MC		LENS PLUS REWETTING DROPS MURO 128 NEO-SYNEPHRINE SOLN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MISCELLANEOUS EAR								
EAR	MC/DEL MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL		A/B OTIC SOLN ACETASOL SOLN ACETASOL HC SOLN ACETIC ACID ACETIC ACID/HYDROCORTISON ALLERGEN SOLN CARBAMIDE PEROXIDE 6.5% OTIC SOLN. CIPRO HC SUSP CORTISPORIN-TC SUSP CORTOMYCIN COLY-MYCIN-S SUSP DERMOTIC EAR DROPS SOLN EAR DROPS RX SOLN EAR WAX REMOVAL DROPS NEOMYCIN/POLYMYXIN/HC OFLOXACIN 0.3% OTIC	MC MC MC/DEL MC/DEL MC MC MC MC MC MC MC MC MC/DEL MC/DEL		ANTIBIOTIC EAR SOLN ANTIBIOTIC EAR SUSP CIPRODEX CIPROFLOXACIN HCL DEBROX SOLN FLOXIN FLUOCINOLONE ACETONIDE OIL DROPS 0.01% OTIPRIO OTOVEL	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MOUTH ANTISEPTICS								
MOUTH ANTI-INFECTIVES	MC MC/DEL		NILSTAT SUSP NYSTATIN SUSP	MC MC		MYCELEX TROC ORAVIG	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MOUTH ANTISEPTICS	MC/DEL MC/DEL MC MC		CHLORHEXIDINE GLUCONATE LIDOCAINE VISCOUS SOLN TRIAMCINOLONE IN ORABASE PSTE TRIAMCINOLONE ORADENT PSTE	MC MC MC MC		APHTHASOL PSTE ¹ PERIOGARD SOLN ¹ TRIAMCINOLONE ACETONIDE PSTE ¹	Use PA Form# 20420 1. Must fail all preferred products before non-preferred.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

DENTAL PRODUCTS							
DENTAL PRODUCTS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		ETHEDENT CREA GEL-KAM CONC GEL-KAM GEL 0.4% PHOS FLUR SOLN SF 5000 PLUS CREA SF GEL STANNOUS FLUORIDE ORAL RI CONC	MCOMC MC/DEL MC/DEL MC		APF GEL GEL DENTAGEL GEL PHOS-FLUR GEL THERA-FLUR-N GEL Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ARTIFICIAL SALIVA/STIMULANTS							
ARTIFICIAL SALIVA/STIMULANTS	MC		SALIVA SUBSTITUTE SOLN	MC MC MC		EVOXAC CAPS RADIACARE SOLR SALAGEN TABS Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MISCELLANEOUS ANORECTAL							
ANORECTAL - MISC.	MC MC MC/DEL MC/DEL MC/DEL		CORTENEMA ENEM ELA-MAX 5 CREA HYDROCORTISONE ENEM PROCTOSOL HC CREA PROCTOZONE-HC CREA	MC/DEL MC/DEL MC/DEL MC/DEL MC		ANUSOL-HC CREA CORTIFOAM FOAM PROCTOFOAM HC FOAM PROCTO-KIT CREA 2.5% RECTIV OINT Use PA Form# 20420	
T-CELL ACTIVATION INHIBITOR							
PSORIASIS BIOLOGICALS	MC MC MC MC MC		ENBREL ^{1,5} ENBREL SURECLICK ¹ HUMIRA ^{1,5} OTEZLA TALTZ ²	MC MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC MC MC MC		AMJEVITA COSENTYX ⁴ CYLTEZO HADLIMA HULIO HYRIMOZ IDACIO ILUMYA ³ SKYRIZI SOTYKTU SPEVIGO SILIQ STELARA TREMIFYA YUFLYMA YUSIMRY Use PA Form# 20910	1. Dosing limits apply, please refer to dosage consolidation list. 2. Clinical PA required and will be preferred for the indication of plaque psoriasis, psoriatic arthritis and ankylosing spondylitis. 3. For the treatment of adults with moderate-to-severe plaque psoriasis who are candidates for systemic therapy or phototherapy. 4. Please see criteria section 5. Will not require a PA if at least one systemic drug such as methotrexate, cyclosporine, methoxsalen or acitretin is in members drug profile. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Cosentyx approvals for 300mg dose(s) must use "300DOSE" package (containing 2 x 150mg pens or syringes). It is recommended to assess for TB infection prior to starting treatment with Taltz®.
ALTERNATIVE MEDICINES							
ALTERNATIVE MEDICINES	MC MC		DIMETHYL SULFOXIDE SOLN MELATONIN	MC/DEL		CO-ENZYME Q-10 Use PA Form# 20420	Will only be approved for specific conditions supported by at least two double-blinded, placebo-controlled randomized trials that are not contradicted by other studies of similar quality.
CHELATING AGENTS							
CHELATING AGENTS	MC/DEL		CUPRIMINE CAPS	MC MC		CLOVIQUE DEPEN TITRATABS TABS Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

				MC/DEL MC MC/DEL		EXJADE ¹ SYPRINE TRIENTINE CAPS	treatment of chronic iron overload due to blood transfusions in membes ?	another drug and the preferred drug(s) exists. Clovique® should be used when continued treatment with penicillamine is no longer possible because of intolerable or life endangering side effects.
ANTILEPROTIC								
ANTILEPROTIC				MC		THALOMID CAPS ¹	1. All PA requests for 150mg dosing will require use of Thalomid 100mg and 50mg capsules. Use PA Form# 20420	Approved for indications of leprosy, treatment-resistant multiple myeloma and AIDS.
ANTINEOPLASTIC AGENTS								
ANTINEOPLASTIC AGENTS - ANTIADNDROGENS	MC/DEL		BICALUTAMIDE	MC/DEL		CASODEX	Use PA Form# 20420	
ANTINEOPLASTIC AGENTS- LHRH ANALOGS	MC/DEL MC/DEL MC/DEL		LUPRON DEPOTSYPHNGEKIT ¹ LUPRON DEPOT- PED KIT ¹ (1-month) LUPRON DEPOT-PED SYPHNGEKIT (3-month) TRIPTODUR VIAL	MC/DEL MC/DEL MC/DEL MC		LUPRON DEPOT SYPHNGEKIT FIRMAGON ² SUPPRELIN LA (IMPLANT) KIT TRELSTAR VANTAS ²	1. Dosing limits apply, please refer to dosage consolidation list. 2. PA required to confirm FDA approved indication. Use PA Form# 20420	
ANTINEOPLASTIC AGENTS - TYROSINE KINASE INHIBITORS				MC MC/DEL MC		SPRYCEL ¹ TYKERB ² GLEEVEC ¹	Use PA Form# 20420 1. Verification of diagnosis is required. 2. PA required to confirm FDA approved indication and to monitor for potential drug-drug interactions.	
ANTINEOPLASTICS-MISCELLANEOUS	MC MC/DEL MC/DEL		AMIFOSTINE MERCAPTOPYRINE OXALIPLATIN	MC MC/DEL MC/DEL MC MC/DEL MC/DEL		DOCEFREZ ELOXATIN ETHYOL LEUPROLIDE PURINETHOL ZOLINZA	Use PA Form# 20420	
ANTINEOPLASTICS- MONOCLONAL ANTIBODIES	MC/DEL		TRAZIMERA	MC/DEL MC/DEL MC MC MC/DEL		ENHERTU HERCEPTIN HERZUMA KANJINTI OGIVRI ONTRUZANT	Use PA Form# 20420	
CANCER								
CANCER	MC MC/DEL MC MC MC/DEL MC		ALIMTA ANASTROZOLE TABS ERBITUX IMATINIB MESYLATE LETROZOLE RUXIENC	MC MC MC/DEL MC MC MC/DEL		ABECMA AKEEGA ALECENSA ALIQOPA ³ ALUNBRIG ¹ ALYMSYS ARIMIDEX	1. PA required to confirm appropriate diagnosis and testing. 2. Avoid CYP3A drug drug interaction.	All non-preferred: A clinical PA is required to confirm appropriate clinical indication for the individual drug request. Specific to each drug all age, clinical testing requirements, previous step therapies, adjunctive drug therapy requirements, and response without disease progression will be also be evaluated for clinical appropriateness. The standard for the appropriate indication will include the FDA label as well as current NCCN guidelines Scemblix is for the treatment of adult patients with: Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP), previously treated with two or more

MC/DEL	LORBRENA
MC	LUMAKRAS
MC/DEL	LUMOXITI ¹
MC	LUNSUMIO ¹
MC	LYNPARZA ¹
MC	LYTGOBI
MC	NEXAVAR ¹
MC	NERLYNX ³
MC	NINLARO(PO)
MC/DEL	NUBEQA
MC	MARGENZA
MC/DEL	MEKINIST ^{3,4}
MC/DEL	MEKTOVI ¹
MC	MONJUVI
MC/DEL	MYLOTARG ³
MC/DEL	MVASI
MC	ODOMZO ^{1,2,5}
MC	OJJAARA
MC	OMISIRGE
MC	ONUREG
MC/DEL	OPDIVO ³
MC	OPDUALAG
MC	ORGOVYX
MC	ORSERDU ^{2,3}
MC	PADCEV
MC	PEMAZYRE
MC	PEPAXTO
MC	PHESGO
MC/DEL	PIQRAY
MC	POLIVY
MC	POMALYST
MC	PORTRAZZA ³
MC	QINLOCK
MC	RETEVMO
MC	REZLIDHIA
MC/DEL	ROZLYTREK
MC	RUBRACA
MC	RITUXAN
MC	RYBREVANT
MC	RYDAPT
MC	RYLAZE
MC/DEL	SARCLISA
MC	SCEMBLIX ¹
MC/DEL	STIVARGA
MC/DEL	SUTENT ^{1,2}
MC/DEL	SYLATRON
MC	TABRECTA
MC	TALVEY
MC/DEL	TAFINLAR ^{3,4,5,6}
MC	TAZVERIK
MC/DEL	TALZENNA ¹
MC/DEL	TAGRISSO
MC	TECARTUS
MC	TECENTRIQ ¹
MC	TEPMETKO
MC/DEL	TIBSOVO ¹

				MC	TIVDAK		
				MC	TRODELVY		
				MC	TRUSELTIQ		
				MC/DEL	TRUXIMA		
				MC	TUKYSA		
				MC	UKONIQ		
				MC/DEL	VANFLYTA		
				MC	VEGZELMA		
				MC	VENCLEXTA ³		
				MC	VERZENIO ³		
				MC/DEL	VITRAKVI		
				MC/DEL	VIZIMPRO ¹		
				MC	VONJO		
				MC/DEL	WELIREG		
				MC/DEL	XALKORI		
				MC/DEL	XPOVIO		
				MC/DEL	XOSPATA		
				MC/DEL	XTANDI		
				MC/DEL	YERVOY		
				MC	YESCARTA ³		
				MC/DEL	ZALTRAP		
				MC	ZEJULA ¹		
				MC/DEL	ZELBORAF		
				MC	ZEPZELCA		
				MC	ZYDELIG		
				MC/DEL	ZYKADIA		
				MC	ZYNLONTA		
				MC	ZYNYZ ¹		
				MC	ZYTIGA		
IMMUNOSUPPRESSANTS							
IMMUNOSUPPRESSANTS	MC/DEL		CYCLOSPORINE MODIFIED	MC/DEL	CELLCEPT	1. For the treatment of adult and pediatric patients 12 years and older with chronic graft-versus-host disease (chronic GVHD) after failure of at least 2 prior lines of systemic therapy	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC		GENGRAF CAPS	MC/DEL	CYCLOSPORINE CAPS		
	MC/DEL		MYCOPHENOLATE	MC/DEL	CYCLOSPORINE SOL. MODIFIED		
	MC/DEL		MYFORTIC	MC	ENVARUSUS XR		
	MC/DEL		NEORAL SOL	MC/DEL	NEORAL CAP		DDI: Cyclosporine will now be non-preferred and require prior authorization if it is currently being used in combination with either Lipitor (doses greater than 20mg/day), Crestor, or lovastatin (doses greater than 20mg).
	MC/DEL		RAPAMUNE	MC	PROGRAF CAPS		
	MC/DEL		SANDIMMUNE	MC	REZUROCK ¹		
	MC/DEL		TACROLIMUS CAPS	MC/DEL	ZORTRESS		DDI: Cyclosporine will require prior authorization when used with Livalo. DDI: All preferred immunosuppressants will require clinical PA for patients over 60 that are currently on fluoroquinolone therapy.
						Use PA Form# 20420	
IMMUNOSUPPRESSANTS- Misc.				MC	HYFTOR ^{1,2}	1. For the treatment of patients ≥ 6 years of age. 2. Clinical PA required for appropriate diagnosis and clinical parameters.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
						Use PA Form# 20420	
PURINE ANALOG							
PURINE ANALOG	MC		AZASAN TABS	MC/DEL	IMURAN TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		AZATHIOPRINE TABS				
K REMOVING RESINS							
K REMOVING RESINS	MC/DEL		LOKELMA	MC/DEL	SPS SUSP	Use PA Form# 20420	
	MC/DEL		SODIUM POLYSTYRENE SULFON	MC/DEL	SPS 30GM/120ML ENEMA SUSP		
				MC	VELTASSA		

|--|--|--|--|--|--|--|--|

New drugs are initially non-preferred until reviewed by the DUR Committee and the State. According to State policy, any drug requiring specific diagnosis still requires the specific diagnosis unless otherwise noted within this document.

Last update 01/17

PDL DOSAGE CONSOLIDATION LIST

Tabs/Caps/Patches: Quantities in units

Shaded areas are non-preferred agents - Quantities of these

Sprays/Inhalers/Nebulizers: Quantities in GM, ML, OR MCG

non-preferred agents are available up the limit only with

Injectibles: Quantities in ML

prior authorization

Drug Name	Strength	Limit/Day	Limit/Days	Drug Name	Strength	Limit/Day	Limit/Days
ABILIFY SOLUTION	1MG/ML	30ML	1020/34	ATROVENT HFA	17MCG	12 INHALATIONS	25.8/34
ACCUPRIL	5MG	1	35/35	ATROVENT 30ML	0.03%	12 SPRAYS	30/30
ACCUPRIL	10MG	1	35/35	ATROVENT 15ML	0.06%	16 SPRAYS	45/30
ACCUPRIL	20MG	1	35/35	AVANDIA	2MG	1.5	53/35
ACEON	2MG	1	35/35	AVANDIA	4MG	1	35/35
ACEON	4MG	1	35/35	AVAPRO	75MG	1.5	53/35
ACTONEL	5MG	1	35/35	AVAPRO	150MG	1	35/35
ACTONEL	35MG	1/WK	5/35	AXERT (Step 8)	6.25MG		12/30
ACTOS	All Strengths	1	35/35	AXERT (Step 8)	12.5MG		12/30
ADDERALL XR	5MG	3	90/30	AZELEX	20%		1 TUBE/18
ADDERALL XR	10MG	3	90/30	AZILECT	All Strengths	1	35/35
ADDERALL XR	15MG	3	90/30	BACTROBAN CREAM			1 TUBE/30
ADDERALL XR	20MG	2	60/30	BECONASE AQ	42MCG	8 INHALATIONS	50/30
ADDERALL XR	30MG	1	35/35	BENICAR-HCT	All Strengths	1	30/30
ADEMPAS	All Strengths	1	35/35	BENAZEPRIL	5MG	1	35/35
ADVAIR DISKUS	All Strengths	2	60/30	BENAZEPRIL	10MG	1.5	53/35
ADVAIR HFA	All Strengths	4	120/30	BENAZEPRIL	20MG	1	35/35
ADZENYS XR	All Strengths	1	30/30	BENAZEP/HCTZ	5-6.25	1	35/35
AEROBID	250MCG	8 INHALATIONS	21/35	BENAZEP/HCTZ	10/12.5	1	35/35
AEROBID-M	250MCG	8 INHALATIONS	21/35	BEVESPI AERO		4 INHALATIONS	120/30
ALAVERT-NON DROW	TAB	1	96/96	BONIVA	2.5MG	1	35/35
ALENDRONATE	All Strengths	1/WK	35/35	BOTOX (ADULTS)	100U/ML	1 session/90 days	600U/90
ALTABAX	5GM		1 TUBE/30	BOTOX (CHILDREN>12)	100U/ML	1 session/90 days	400U/90
ALTABAX	15GM		1 TUBE/30	BREO ELLIPTA	100/25MCG	1 INHALATIONS	60/60
ALTABAX	30GM		1 TUBE/30	BRILINTA	All Strengths	2	70/35
ALTACE	1.25MG	1	35/35	BRINTELLIX	All Strengths	1	35/35
ALTACE	2.5MG	1	35/35	BUTRANS		1 patch/WK	4/28
ALTACE	5MG	1	35/35	BYETTA	5mcg inj	0.04ML	1.2ML/30
AMARYL	1MG	1	35/35	BYETTA	10mcg inj	0.08ML	2.4ML/30
AMARYL	2MG	1	35/35	CALAN SR	120MG	1	35/35
AMBIEN	5MG		12/34	CALAN SR	180MG	2	70/35
AMBIEN	10MG		12/34	CALAN SR	240MG	2	70/35
AMBIEN CR	6.25MG		12/34	CARDIZEM CD	120MG/24	1	35/35
AMBIEN CR	12.5MG		12/34	CARDIZEM CD	180MG/24	1	35/35
AMERGE (Step 8)	1MG		12/30	CARDIZEM CD	240MG/24	1	35/35
AMERGE (Step 8)	2.5MG	2.5MG	12/30	CARDIZEM CD	300MG/24	1	35/35
AMLODIPINE	2.5MG	1.5	53/35 DAYS	CARDIZEM CD	360MG/24	1	35/35
AMLODIPINE	5MG	1.5	53/35 DAYS	CARDIZEM LA	120MG/24	1	35/35
AMMONIUM LACTATE CREA	12%		1 TUBE/10	CARDIZEM LA	180MG/24	1	35/35
AMMONIUM LACTATE LOTN	12%		1TUBE/8	CARDIZEM LA	240MG/24	1	35/35
AMPHETAMINE/DEXTROAMPHET ER	5MG	3	90/30	CARDIZEM LA	300MG/24	1	35/35
AMPHETAMINE/DEXTROAMPHET ER	10MG	3	90/30	CARDIZEM LA	360MG/24	1	35/35
AMPHETAMINE/DEXTROAMPHET ER	15MG	3	90/30	CARDURA	1MG	1	35/35
AMPHETAMINE/DEXTROAMPHET ER	20MG	2	60/30	CARDURA	2MG	1.5	53/35
AMPHETAMINE/DEXTROAMPHET ER	30MG	1	90/90	CARDURA	4MG	1.5	53/35
AMPHETAMINE SALT	5,10,15MG	3	105/35	CARTIA XT	120MG	1	90/90
AMPHETAMINE SALT	20MG	2	70/35	CARTIA XT	180MG	1	90/90
AMPHETAMINE SALT	30MG	1	35/35	CARTIA XT	240MG	1	90/90
ANDRODERM	2.5MG	2	60/30	CARTIA XT	300MG	1	90/90
ANDRODERM	5MG	1	30/30	CATAPRES-TTS1	0.1 MG/24HR		5/35
ARAVA	10MG	1	35/35	CATAPRES-TTS2	0.2 MG/24HR		5/35
ARCAPTA	75MCG	1 INHALATION	35/35	CATAPRES-TTS3	0.3 MG/24HR		5/35
ARICEPT	5MG	1	35/35	CEFIXIME	400MG	2	2/7
ARICEPT	10MG	1	35/35	CELEBREX	100MG	1	35/35
ARIPIRAZOLE	2MG	2	180/90	CELEBREX	200MG	2	70/35
ARIPIRAZOLE	5MG	2	180/90	CELEBREX	400MG	1	35/35
ARIPIRAZOLE	10MG	2	180/90	CELEXA	20mg	0.5	17/34
ARIPIRAZOLE	15MG	2	180/90	CELEXA	40mg	1	51/34
ARIPIRAZOLE	20MG	1.5	135/90	CITALOPRAM	10MG	2	180/90
ARIPIRAZOLE	30MG	1	90/90	CITALOPRAM	20MG	2	180/90
ARIXTRA INJECTION	2.5MG/0.5ML		7/30	CITALOPRAM	40MG	1	90/90
ARIXTRA INJECTION	5MG/0.4ML		7/30	CLARINEX	REDI TAB	1	35/35
ARIXTRA INJECTION	7.5MG/0.6ML		7/30	CLEOCIN-T		1 PACKAGE	1/30
ARIXTRA INJECTION	10MG/0.8ML		7/30	CLINDAMYCIN PHOSPHATE		1 PACKAGE	1/30
ARONAIR	All Strengths	1 INHALATION	60U/30	COMBIVENT	103-18MCG	12 INHALATIONS	30/35
ASMANEX 30 UNITS	220MCG	1 INHALATION	30U/30	Drug Name	Strength	Limit/Day	Limit/Days
ASMANEX 60 UNITS	220MCG	2 INHALATIONS	60U/30	EFFEXOR XR	37.5MG	1	35/35
ASMANEX 120 UNITS	220MCG	4 INHALATIONS	120U/30	EFFEXOR XR	75MG	1	35/35
ATACAND	4MG	1.5	53/35	EMSAM	All Strengths	1	34/34
				ENALAPRIL	2.5	1	90/90

ATACAND	8MG	1.5	53/35
ATACAND	16MG	1	35/35
ATRIPLA	600MG	1	35/35
Drug Name	Strength	Limit/Day	Limit/Days
COMETRIQ	80MG	1	35/35
COMETRIQ	20MG	3	105/35
CONCERTA	18MG	1	30/30
CONCERTA	27MG	1	30/30
CONCERTA	36MG	2	60/30
COPAXONE INJ	20MG		1/32
COPAXONE KIT	20MG/ML		1/30
COREG CR	All Strengths	1	34/34
COSENTYX	150MG	1	1/30
CRESTOR	5MG	1	35/35
CRESTOR	10MG	1	35/35
CRESTOR	20MG	1	35/35
CRESTOR	40MG	1	35/35
CYMBALTA	All Strengths	1	35/35
DALMANE	15MG		10/30
DALMANE	30MG		10/30
DAYPRO	600MG	2	70/35
DAYTRANA	10mg/9hr (27.5mg)	1	34/34
DAYTRANA	15mg/9hr (41.3mg)	1	34/34
DAYTRANA	20mg/9hr (55.0mg)	1	34/34
DAYTRANA	30mg/9hr (82.5mg)	1	34/34
DDAVP	5ML		15/34
DENAVIR CREAM			2gm/30
DEPO-PROVERA	150MG/ML		1/90
DEPO-PROVERA	400MG/ML		2.5/90
DEPO-TESTOSTERONE	200MG/ML		20/90
DESMOPRESSIN	0.1MG	12	420/35
DESMOPRESSIN	0.2MG	6	210/35
DESONIDE	0.05%		2 TUBES/30
DESOWEN	0.05%		2 TUBES/30
DETROL LA	2MG	1	35/35
DEXEDRINE	All Strengths	3	90/30
DEXILANT	All Strengths	1	35/35
DEXTROAMPHETAMINE	All Strengths	3	90/30
DICLOFENAC 1% GEL	1% GEL		2 TUBES/30
DIFLUCAN	150MG		1/7
DILACOR XR	240MG/24	1	35/35
DILACOR XR	120MG/24	1	35/35
DILACOR XR	180MG/24	1	35/35
DILTIA - XT	120MG/24	1	90/90
DILTIA - XT	180MG	1	90/90
DILTIA - XT	240MG/24	1	90/90
DILTIAZEM CAP ER	120MG	1	90/90
DILTIAZEM CAP XR	120MG	1	90/90
DILTIAZEM CAP	120MG/24	1	90/90
DILTIAZEM CAP	180MG/24	1	90/90
DILTIAZEM CAP ER	240MG	1	90/90
DILTIAZEM CAP XR	240MG	1	90/90
DILTIAZEM XR CAP	240MG/24	1	90/90
DILTIAZEM CAP	240MG/24	1	90/90
DILTIAZEM CAP	300MG/24	1	90/90
DILTIAZEM CAP	360MG/24	1	90/90
DIOVAN	80MG	1	35/35
DIOVAN - HCT	80 - 12.5	1	35/35
DITROPAN XL	5MG	1	35/35
DITROPAN XL	10MG	2	70/35
DORAL	7.5MG		10/30
DOXAZOSIN	1MG	1	90/90
DOXAZOSIN	2MG	1.5	135/90
DOXAZOSIN	4MG	1.5	135/90
DRYSOL SOL	20%		1 BOTTLE/30DAYS
DURAGESIC PATCHES	12.5MCG/HR		11/33
DURAGESIC PATCHES	25MCG/HR		11/33
DURAGESIC PATCHES	50MCG/HR		11/33
DURAGESIC PATCHES	75MCG/HR		11/33
DURAGESIC PATCHES	100MCG/HR		22/33
DULOXETINE	20MG	3	270/90
DULOXETINE	30MG	3	270/90
DULOXETINE	60MG	2	180/90
EDEX	All Strengths		1/30
Drug Name	Strength	Limit/Day	Limit/Days

ENALAPRIL	5MG	1.5	135/90
ENALAPRIL	10MG	1.5	135/90
ENALAPR/HCTZ	5-12.5	1	90/90
ENBREL	25MG/ML		8/28
ENBREL SURECLICK			8/28
ESTAZOLAM	1MG		10/30
ESTAZOLAM	2MG		10/30
ESTRING MIS	2MG		1/90
EVENITY		12 DOSES/LIFETIME	12 DOSES/LIFETIME
EVOTAZ	All Strengths	1	30/30
FELODIPINE	2.5MG	1	90/90
FELODIPINE	5MG	1.5	135/90
FENTANYL	25MCG/HR		11/33
FENTANYL	50MCG/HR		11/33
FENTANYL	75MCG/HR		11/33
FENTANYL	100MCG/HR		22/33
FETZIMA	All Strengths	1	35/35
FINASTERIDE	5MG	1	90/90
FLONASE	50MCG	4 SPRAYS	32/34
FLOVENT HFA 44MCG	44MCG	4 INHALATIONS	10.6/30
FLOVENT HFA 110MCG	110MCG	4 INHALATIONS	12/30
FLOVENT HFA 220MCG	220MCG	8 INHALATIONS	24/30
FLOVENT DISKUS	50MCG, 100MCG	4 INHALATIONS	60/30
FLOVENT DISKUS	250MCG	3 INHALATIONS	120/30
FLUCONAZOLE	150MG		1/7
FLUNISOLIDE SOLN	0.025%	16 SPRAYS	75/30
FLUOXETINE CAP	40MG	2	180/90
FLUOXETINE CAP	20MG	4	360/90
FLUOXETINE CAP	10MG	3	270/90
FLURAZEPAM	15MG		10/30
FLURAZEPAM	30MG		10/30
FLUTICASONE SPR		4 SPRAYS	48/90
FLUVOXAMINE	25MG	3	270/90
FLUVOXAMINE	50MG	3	270/90
FOCALIN	All Strengths	3	105/35
FOCALIN XR	All Strengths	1	35/35
FORFIVO XL	All Strengths	1	35/35
FOSAMAX	5MG	1	35/35
FOSAMAX	10MG	1	35/35
FOSAMAX	70MG	1/WK	5/35
FOSAMAX	40MG	2/WK	10/35
FOSINOPRIL	10MG	1.5	135/90
FOSINOPRIL	20MG	2	180/90
FRAGMIN INJ	10000U/ML	2ML	14/7
FRAGMIN INJ	2500U/.2ML	0.4ML	2.80/7
FRAGMIN INJ	25000U/ML	0.8ML	5.6/7
FRAGMIN INJ	5000U/.2ML	0.4ML	2.80/7
FRAGMIN INJ	7500U/.3ML	0.6ML	4.2/7
FROVA TAB (Step 8)	2.5MG		12/30
FULYZAQ	125MG	2	70/35
FUZEON	KIT	1	1/30
FYCOMPA	All Strengths	1	35/35
GABAPENTIN	300MG	9	810/90
GABAPENTIN	400MG	9	810/90
GABAPENTIN	600MG	6	540/90
GABAPENTIN	800MG	4	360/90
GEODON	20MG	2	70/35
GEODON	40MG	2	70/35
GEODON	60MG	2	70/35
GEODON	80MG	2	70/35
GEODON	INJ	2	70/35
GILOTRIF	All Strengths	1	35/35
GLIMEPIRIDE	1MG	1	90/90
GLIMEPIRIDE	2MG	1	90/90
GLUCOSE TES STRP		12	420/35
GLUCAGEN INJ. HYPOKIT			2/30
GLYCOLAX*	255GM		255GM/90
* Available for once daily dosing to members under the age of 18 years			
Drug Name	Strength	Limit/Day	Limit/Days
LUNESTA	2MG		12/34
LUNESTA	3MG		12/34
LUPRON DEPOT INJ	11.25MG	KIT	1/90
LUPRON DEPOT INJ	22.5	KIT	1/90
LUPRON DEPOT INJ	30MG		1/90

ILARIS			2/28
HALCION	0.125MG		10/35
HALCION	0.25		10/35
HUMIRA	40mg/0.8ml		4/28
HYDROXYZINE TAB	All Strengths	3	270/90
HYTRIN	1MG	1	35/35
HYTRIN	5MG	1	35/35
HYZAAR	50-12.5	1	35/35
IMDUR	30MG	1.5	53/35
IMDUR	60MG	1.5	53/35
IMITREX (step 8)	25MG		12/30
IMITREX (step 8)	50MG		12/30
IMITREX (step 8)	100MG		12/30
IMITREX VIAL	All Strengths		6 boxes/30
IMITREX CARTRIDGE	All Strengths		12/30
IMITREX NASAL SPRAY	All Strengths		12/30
IMITREX PEN INJCTR	All Strengths		12/30
IMIQUIMOD	5%		12/30
IMIQUIMOD	5%		12/30
INTAL	800MCG	8 INHALATIONS	28.4/34
INVOKANA	All Strengths	1	35/35
IPRATROPIUM 30ML	0.03%	12 SPRAYS	90/90
IPRATROPIUM 15ML	0.06%	16 SPRAYS	135/90
ISOPTIN SR	180MG	2	70/35
IRBESARTAN	All Strengths	1	90/90
ISOPTIN SR	240MG	2	70/35
ISOSORBIDE MONO	30MG	2	180/90
ISOSORBIDE MONO	60 MG	1.5	135/90
JANUMET	All Strengths	2	70/35
JANUVIA	All Strengths	1	35/35
JUVISYNC	All Strengths	1	35/35
KETOPROFEN	100MG	2	180/90
KETOPROFEN	200MG	1	90/90
KETOROLAC	10MG	4.8	24/30
KHEDEZLA	All Strengths	1	35/35
LAC-HYDRIN CREAM	12%		1TUBE/30
LAMICTAL	25MG	6	210/35
LAMICTAL	25MG CHW	6	210/35
LAMICTAL	100MG	2	70/35
LAMISIL	250MG	1	35/35
LAMOTRIGINE	25MG	6	540/90
LAMOTRIGINE	100MG	2	180/90
LANSOPRAZOLE CAPS	All Strengths	1	90/90
LEFLUNOMIDE	10MG	1	90/90
LESCOL	20MG	1	35/35
LEVAQUIN	250MG	1	35/35
LEXAPRO	5MG	0.5	15/30
LIPITOR	10MG	1	35/35
LIPITOR	20MG	1	35/35
LIPITOR	40MG	1.5	53/35
LISINOP/HCTZ	10/12.5MG	1	90/90
LOSARTAN	All Strengths	1	90/90
LOSARTAN- HCT	All Strengths	1	90/90
LOTENSIN	5MG	1	35/35
LOTENSIN	10MG	1.5	35/35
LOTENSIN	20MG	1	53/35
LOTENSIN - HCT	5 - 6.25	1	35/35
LOTENSIN - HCT	10 - 12.5	1	35/35
LOVASTATIN	10MG	1.5	135/90
LOVASTATIN	20MG	1.5	135/90
LOVENOX INJ	30MG/.3ML	0.6	14 injections/7
LOVENOX INJ	40MG/.4ML	0.8	14 injections/7
LOVENOX INJ	60MG/.6ML	1.2	14 injections/7
LOVENOX INJ	80MG/.8ML	1.6	14 injections/7
LOVENOX INJ	100MG/ML	2	14 injections/7
LOVENOX INJ	120MG/.8ML	1.6	14 injections/7
LOVENOX INJ	150MG/ML	2	14 injections/7
LUNESTA	1MG		12/34
Drug Name	Strength	Limit/Day	Limit/Days
NIFEDIPINE ER	90MG	1	90/90
NIFEDIPINE ER,CR	30MG	1	90/90
NORVASC	2.5MG	1.5	53/35 DAYS
NORVASC	5MG	1.5	53/35 DAYS
NURTEC ODT	All Strengths		8/30
NUVARING		1/MO	1/28

LUPRON DEPOT INJ	30MG	KIT	1/90
LYRICA	25,50,75MG	3	102/35
LYRICA	100,150,200MG	3	102/35
LYRICA	225,300MG	2	70/35
MAVIK	1MG	1	35/35
MAVIK	2MG	1	35/35
MAXAIR AUTO	200MCG	12 INHALATIONS	14/30
MAXALT (step 8)	5MG		12/30
MAXALT (step 8)	10MG		12/30
MAXALT MLT (step 1)	5MG		12/30
MAXALT MLT (step 1)	10MG		12/30
MEDROXYPR AC	150MG/ML		1/90
MELOXICAM TABS	All Strengths	1	90/90
METADATE ER	10,20MG	3	90/30
METFORMIN ER	500MG	4	360/90
METHYLIN	All Strengths	3	90/30
METHYLPHENIDATE ER	36mg	2	180/90
METHYLPHENIDATE	All Strengths	3	90/30
METROCREAM		1 PACKAGE	1/30
METROGEL		1 PACKAGE	1/30
METROLOTION		1 PACKAGE	1/30
METRONIDAZOLE CREAM		1 PACKAGE	1/30
METRONIDAZOLE GEL		1 PACKAGE	1/30
METRONIDAZOLE LOTION		1 PACKAGE	1/30
MEVACOR	10MG	1.5	53/35
MEVACOR	20MG	1.5	53/35
MIACALCIN		3.75ml	1 bottle/34
MICARDIS	All Strengths	1	30/30
MICARDIS-HCT	All Strengths	1	30/30
MIGRANAL NASAL SPRAY	All Strengths		12/30
MIRALAX	255G	8.5G	1 bottle/30
MIRALAX	17G/PACKET	0.5 packet	15 packets/30
MIRTAZAPINE	15mg	3	270/90
MOBIC	7.5 MG	1	35/35
MOBIC	15MG	1	35/35
MOEXIPRIL	7.5	1.5	135/90
MONOPRIL	10MG	1.5	53/35
MONOPRIL	20MG	2	70/35
MUPIROCIN			1 TUBE/30
NABUMETONE	500MG	2	180/90
NABUMETONE	750MG	2	180/90
NARATRIPTAN			12/30
NASACORT AERS	55 MCG	4 SPRAYS	9.3/25
NASONEX	50MCG	4 SPRAYS	17/30
NATROBA		120ML	1 bottle/30
NAYZILAM	All Strengths		5/30
NEUPOGEN INJ	300MCG/ML		10/30
NEUPOGEN INJ	480MCG/1.6		16/30
NEUPOGEN INJ	300MCG/.5ML		5/30
NEUPOGEN INJ	480MCG/.8ML		8/30
NEURONTIN	300MG	9	315/35
NEURONTIN	600MG	9	315/35
NEXIUM	20MG	1	35/35
NEXIUM	40MG	2	70/35
NEXIUM SUS	All Strengths	1	30/30
NIFEDIPINE CR	90MG	1	90/90
NIFEDIPINE ER	60MG	1	90/90
NIFEDIPINE ER	30MG	1	90/90
NIFEDIPINE ER	60MG	1	90/90
Drug Name	Strength	Limit/Day	Limit/Days
RELPAX	All Strengths		12/30
REMODULIN	All Strengths		1 MDV/30
RESTORIL	7.5MG		10/30
RESTORIL	15MG		10/30
RESTORIL	30MG		10/30
RETIN-A		1 TUBE	1 TUBE/30
REVLIMID	All Strengths	1	35/35
REYVOW	All Strengths		4/30
RHINOCORT AQ	32MCG	8 SPRAYS	18/30
REFRESH PLUS		15 ML	1 bottle/30
REFRESH PLUS		30 ML	2 bottles/30
REFRESH TEARS		15 ML	1 bottle/30
REFRESH TEARS		30 ML	2 bottles/30
RESCULA			2 bottles/35
REYATAZ	All Strengths	1	35/35

ODOMZO	200mg	1	30/30
OLMESARTAN	All Strengths	1	90/90
OLANZAPINE	2.5MG	3	270/90
OLANZAPINE	5MG	3	270/90
OLANZAPINE	7.5MG	3	270/90
OLANZAPINE	10MG	3	270/90
OLANZAPINE	15MH	2	180/90
OLANZAPINE	20MG	1.5	135/90
OLANZAPINE ODT	All Strengths	1	90/90
OMEPRAZOLE	10MG	1	90/90
OMEPRAZOLE	20MG	1	90/90
OMEPRAZOLE	40MG	1	90/90
OMNARIS	50MCG	4 sprays	12.5/30
ONGLYZA	All Strengths	1	35/35
OPSUMIT	All Strengths	1	35/35
ORUVAIL	100MG	2	70/35
ORUVAIL	200MG	1	35/35
OXAPROZIN	600MG	2	180/90
OXYCODONE ER	10,20,40MG	2	70/35
OXYCODONE ER	80MG	4	140/35
OXYCONTIN**	10,20,30,40MG	2	70/35
OXYCONTIN**	80MG	4	140/35
PANTOPRAZOLE	All Strengths	1	90/90
PAROXETINE	10MG	2	180/90
PAROXETINE	20MG	2	180/90
PAXIL	10MG	1.5	53/35
PAXIL	20MG	1	35/35
PEGASYS KIT		KIT	1/28
PLAN B			2/15 or 4/30
PLENDIL	2.5MG	1	35/35
PLENDIL	5MG	1.5	53/35
PRAVACHOL	10MG	1	35/35
PRAVACHOL	20MG	1	35/35
PRAVACHOL	40MG	1	35/35
PRAVACHOL	80MG	1	35/35
PRAVASTATIN	10MG	1	35/35
PRAVASTATIN	20MG	1	35/35
PRAVASTATIN	40MG	2	180/90
PRAVASTATIN	80MG	1	35/35
PREVPAC MIS	500MG-30MG		14/30
PRILOSEC OTC	20MG	2	168/84
PRINIVIL	2.5MG	1	35/35
PRINIVIL	5MG	1	35/35
PRINIVIL	10MG	1.5	53/35
PRINIVIL	20MG	1.5	53/35
PRINZIDE	10-12.5	1	35/35
PROAIR HFA	90mcg	12 INHALATIONS	17/34
PROTONIX	20MG	2	70/35
PROTONIX	40MG	2	70/35
PROZAC	10MG	1.5	53/35
PULMICORT	200MCG	8 INHALATIONS	1/25
PULMICORT FLEX	All Strengths	8 Inhalations	2/30
QUETIAPINE	25MG	3	270/90
QUETIAPINE	50MG	3	270/90
QUETIAPINE	100MG	3	270/90
QUETIAPINE	200MG	3	270/90
QUINAPRIL	5MG	1	90/90
QUINAPRIL	10MG	1	90/90
QUINAPRIL	20MG	1	90/90
QVAR AERS	All Strengths	8 Inhalations	14.6/25
RANITIDINE SYRUP***	15MG/ML	20ML	700ML/35
RELAFEN	500MG	2	70/35
RELAFEN	750MG	2	70/35
REMERON	15MG	1.5	53/35
Drug Name	Strength	Limit/Day	Limit/Days
SULAR	10MG	1.5	53/35
SULAR	20MG	1	35/35
SUMATRIPTAN PEN INJ	All Strengths		12/30
SUMATRIPTAN NASAL SPRAY	All Strengths		12/30
SUMATRIPTAN SYRINGE	All Strengths		12/30
SUMATRIPTAN TAB	All Strengths		12/30
SYNVISC INJ	8MG/ML		2/30
SYRINGES		10	1000/100
TAFINLAR	50MG	6	210/35
TAFINLAR	75MG	4	140/35

RISPERDAL	0.5MG	1.5	53/35
RISPERDAL	0.25MG	1.5	53/35
RISPERDAL	1MG	1.5	53/35
RISPERDAL	2MG	1.5	53/35
RISPERDAL	3MG	2	70/35
RISPERDAL	4MG	2	70/35
RISPERDAL INJ	25MG		2/28
RISPERDAL INJ	37.5		2/28
RISPERDAL INJ	50MG		2/28
RISPERDAL M-TAB	0.5MG	1.5	53/35
RISPERDAL M-TAB	1MG	1.5	53/35
RISPERDAL M-TAB	2MG	4	140/35
RISPERDAL SOL.	1MG/ML	8ML	280/35
RISPERIDONE	0.5MG	3	270/90
RISPERIDONE	0.25MG	3	270/90
RISPERIDONE	1MG	3	270/90
RISPERIDONE	2MG	3	270/90
RISPERIDONE	3MG	2	180/90
RISPERIDONE	4MG	2	180/90
RISPERIDONE SOL.	1MG/ML	8ML	280/35
RITALIN LA	All Strengths	1	35/35
RITALIN LA	30mg	2	70/35
SAVELLA	All Strengths	2	70/35
SEREVENT DISKUS	50MCG	2 INHALATIONS	60/30
SEROQUEL	100MG		45/30
SEROQUEL XR	150MG	1	35/35
SEROQUEL XR	200MG	1	35/35
SEROQUEL XR	300MG	2	70/35
SEROQUEL XR	400MG	2	70/35
SERTRALINE	25MG	3	270/90
SERTRALINE	50MG	3	270/90
SERTRALINE	100MG	3	270/90
SIMVASTATIN	5MG	1	35/35
SIMVASTATIN	10MG	1.5	53/35
SIMVASTATIN	20MG	1.5	53/35
SIMVASTATIN	40MG	1.5	53/35
SIMVASTATIN	80MG	1	35/35
SINGULAIR	4MG	1	35/35
SINGULAIR	5MG	1	35/35
SINGULAIR	10MG	1	35/35
SONATA	5MG		12/34
SONATA	10MG		12/34
SPIRIVA	HANDHLR	1 INHALTION	30/30
SPORANOX SOL	10MG/ML	10ML/ML	300cc/30
SPORANOX PULSEPAK	F		30/30
SPORANOX	100MG		30/30
STADOL INJ	1MG/ML		9/35
STADOL INJ	2MG/ML		9/35
STRATTERA	All Strengths	1	35/35
SUPRAX	400MG	1	1/7

Drug Name	Strength	Limit/Day	Limit/Days
XOPENEX HFA		12 INHALATIONS	2 INHALERS/34
XOPENEX NEB		12CC	408/34
ZALEPLON	All Strengths		30/30
ZECUITY	6.5		4/28
ZEMBRACE	All Strengths		3boxes/30
ZESTORETIC	10-12.5	1	35/35
ZESTRIL	2.5MG	1	35/35
ZESTRIL	5MG	1	35/35
ZESTRIL	10MG	1.5	53/35
ZESTRIL	20MG	1.5	53/35
ZETONNA	37MCG	2	60/30
ZIPRASIDONE	20MG	3	270/90
ZIPRASIDONE	40MG	3	270/90
ZOCOR	5MG	1	35/35
ZOCOR	10MG	1.5	53/35
ZOCOR	20MG	1.5	53/35
ZOCOR	40MG	1.5	53/35
ZOFRAN*	4MG	3	90/30
ZOFRAN*	8MG	1.5	45/30
ZOFRAN*	24MG	0.5	15/30
ZOFRAN*	4MG/5ML	15ML	450/30
ZOLMITRIPTAN TAB	All Strengths		12/30
ZOLOFT	25MG	0.5	18/35

TAMIFLU CAPS	75MG		10/30
TAZTIA XT CAP	120MG/24	1	90/90
TAZTIA XT CAP	180MG/24	1	90/90
TAZTIA XT CAP	240MG/24	1	90/90
TAZTIA XT CAP	300MG/24	1	90/90
TAZTIA XT CAP	360MG/24	1	90/90
TELMISARTAN	All Strengths	1	90/90
TEMAZEPAM	7.5MG		10/30
TEMAZEPAM	15MG		10/30
TEMAZEPAM	30MG		10/30
TEQUIN	200MG	1	35/35
TERAZOSIN	1MG	1	90/90
TERAZOSIN	5MG	1	90/90
TERBINAFINE	250MG	1	35/35
TEST STRIPS	Blood Glucose	12	420/35
TIAZAC	120MG/24	1	35/35
TIAZAC	180MG/24	1	35/35
TIAZAC	240MG/24	1	35/35
TIAZAC	300MG/24	1	35/35
TIAZAC	360MG/24	1	35/35
TIAZAC	420MG/24	1	35/35
TILADE	1.75MG	8 INHALATIONS	48.6/35
TOPAMAX SPRINKLES	All Strengths	1	35/35
TOPROL XL	25MG	1.5	53/35
TOPROL XL	50MG	1.5	53/35
TRADJENTA	All Strengths	1	35/35
TRAMADOL	50MG	8	720/90
TRAMADOL/ APAP	37.5/325MG	8	720/90
TRETINOIN		1 TUBE	1 TUBE/30
TRELEGY ELLIPTA	All Strengths	1 INHALATION	30U/30
TREXIMET	85/500	2.5	12/30
TRIAZOLAM	0.125MG		10/30
TRIAZOLAM	0.25MG		10/30
TROKENDI XR	25MG	1	35/35
TROKENDI XR	50MG	1	35/35
TROKENDI XR	100MG	1	35/35
TROKENDI XR	200MG	2	70/35
UBRELVY	All Strengths		10/30
ULTRAM	50MG	8	280/35
UNIVASC	7.5MG	1.5	53/35 DAYS
UTIBRON	7.5mcg/15.6mc	2 INHALATIONS	60/30
VALTOCO	All Strengths		10/30
VALSARTAN-HCT	All Strengths	1	90/90
VASERETIC	5-12.5MG	1	35/35
VASOTEC	2.5MG	1	35/35
VASOTEC	5MG	1.5	53/35
VASOTEC	10MG	1.5	53/35
VENLAFAXINE TABS	25	3	270/90
VENLAFAXINE TABS	37.5	3	270/90
VENLAFAXINE TABS	100	3	270/90
VENLAFAXINE ER CAPS	37.5	3	270/90
VENLAFAXINE ER CAPS	75	3	270/90
VENLAFAXINE ER	150	2	180/90
VENTOLIN HFA	90MCG	12 INHALATIONS	36/34
VERAPAMIL ER, SR	120MG	1	90/90
VERAPAMIL ER, CR, SR	180MG	2	90/90
VERAPAMIL ER, CR, SR	240MG	2	90/90
VERELAN	180MG	1	35/35
VERELAN SR	120MG	1	35/35
VERELAN SR	180MG	1	35/35
VERELAN SR	240MG	2	70/35
VERAMYST	27.5MCG	4 sprays	10/30
VYEPTI	All Strengths		4/30
VYVANSE	All Strengths	1	35/35
VYVANSE CHEW	All Strengths	1	35/35

ZOLOFT	50MG	0.5	18/35
ZOLOFT	100MG	3	105/35
ZOLPIDEM (step 1)	5MG		30/30
ZOLPIDEM (step 1)	10MG		30/30
ZOMIG (Step 8)	5MG		12/30
ZTLIDO	All Strengths	3	90/30
ZYPREXA	2.5MG	1.5	53/35
ZYPREXA	5MG	1	35/35
ZYPREXA	7.5MG	1	35/35
ZYPREXA	10MG	1	35/35
ZYPREXA	15MG	1	35/35
ZYPREXA	20MG	1	35/35
ZYPREXA ZYDIS	5MG	1	35/35
ZYPREXA ZYDIS	10MG	1	35/35
ZYPREXA ZYDIS	15MG	1	35/35
ZYPREXA ZYDIS	20MG	1	35/35

*Cancer diagnosis with non-daily chemotherapy required

**Available without pa with CA and HO diag.

*** Ranitidine syrup available without PA to users less than 6 years old.

MDV=Multidose Vial

Pain Management Policy

[Back to Index](#)

Beginning January 2017, all current opiate users who are above the maximum combined daily dose of 100 MME must titrate their total daily dose of opioid medications below 300 MME. Also, the maximum daily supply of an opiate prescription for acute pain will be limited to 7-day supplies. The maximum day supply of an opiate prescription for chronic pain will be limited to 30-day supplies. As of July 1, 2017 all users of opioid medications must comply with the maximum combined daily dose of 100 MME.

However, for MaineCare members, effective January 1, 2017, opioid prescription(s) for more than a 7-day supply and/or more than 30 MME/ day will require a prior authorization. Please note that MaineCare implemented a 30 MME limit January 1, 2013 that is still effective.

The following are general exceptions: pain associated with cancer treatment, end-of-life and hospice care, palliative care, and symptoms related to HIV/AIDS. Per MaineCare criteria, the diagnosis of cancer must be written on the prescription. A palliative care exception for any MaineCare opioid prescription will require prior authorization (PA) with appropriate clinical documentation.

Post-surgical members may receive prior authorizations for opiates up to a 60 days in length if medical necessity is provided by the surgical provider.

An MME conversion chart is available at www.mainearepd.org. Click on "General Pharmacy Info."