

| CATEGORY | Coverage Indicator | Step Order | PREFERRED DRUGS | Coverage Indicator | Step Order | NON-PREFERRED DRUGS | PA Required | Criteria |
|----------|--------------------|------------|-----------------|--------------------|------------|---------------------|-------------|----------|
|----------|--------------------|------------|-----------------|--------------------|------------|---------------------|-------------|----------|

PDL Effective November 8, 2024

***PLEASE NOTE: For a search box hit Ctrl F**

*** PLEASE NOTE: All cost effective generics applicable to DEL are considered PREFERRED Drugs. "BASIC" Covered Drugs are bolded with the Coverage Indicator of "MC / DEL".**

General Criteria for all PDL categories- For more information or help using the PDL, providers may call 1-888-445-0497; members should call 1-866-796-2463. To access PDL and PA materials via the internet: www.mainearepdl.org

A: Preferred Drugs- Unless otherwise specified, preferred drugs are available without prior authorization. Step order may apply for preferred drugs in some drug categories as indicated on the PDL. (See item "D" below for explanation of step order.)

C: Adequate Drug Trials- 1. The minimum trial period for each preferred and step order drug is two weeks, unless otherwise stated within specific PDL drug categories; trials with less than a two week duration will be reviewed on a case-by-case basis; 2. A trial will not be considered valid if preferred or non-preferred products were readily available (by override, individual purchase, samples, etc.); 3. Certain drug trials, such as with controlled substances, may require evidence that the preferred drugs were actually tried (example: with random pill counts and with random urine drug tests, using the methods of GC/MS with no lower threshold); 4. Adequate trials require documentation of attempts to titrate dose of preferred agents toward desired clinical response. 5. Adequate trials include prevention/treatment of common adverse effects associated with preferred agents (example: antinausea, antipruritics, etc.)

D: Step Order- When numbers appear in the "step order" column, it means drugs in this category must be used in the order specified, with the lower numbers having preference over the higher numbers. Chart notes should be provided to confirm drug trials that do not appear in the member's MaineCare drug profile.

E. The Department will institute strategies to ensure cost effectiveness through the use of an enhanced Drug Benefit Preferred brand drugs will no longer be preferred in any PDL drug category where preferred generic drugs are also available. It is expected that preferred generics will be used prior to any preferred brands. This will be operated as a form of step care. Preferred brands in these categories will require prior authorization for these high utilization / high cost members.

F: Brand Name Medication Requests- (Must be submitted on the Brand Name PA request form)- According to MaineCare Benefits Manual Chapter II (80.07-5), when medically necessary covered brand-name drugs have an A-rated generic equivalent available, the most cost effective medically necessary version will be approved and reimbursed, since the brand-name and A-rated generic drugs have been determined by the FDA to be chemically and therapeutically equivalent. The Bureau does not make determinations as to whether or not a generic drug is clinically inferior or inequivalent to its brand version. This is the proper role of the FDA. Physicians should submit their reports of generic inequivalence directly to the FDA via the MEDWATCH.

G: PA requests for non- FDA Approved Indications- Decisions will be made on a case-by-case basis until the DUR committee is able to review the evidence and make a recommendation. Interim approvals and DUR recommendations for approval of a drug for a non- FDA approved indication will require a minimum of two published, peer reviewed, non contradicted, double- blind, placebo-controlled randomized clinical studies establishing both safety and efficacy.

H: Dose Consolidation Requirements- Some drugs may also be affected by dose consolidation requirements. Please see Dose Consolidation List and/or Splitting Tables provided in the PDL.

I. Trials from Multiple Drug Classes - Trial/failure/intolerance to preferred agents from multiple classes within the same category or other categories of drugs may be required prior to the approval of non-preferred agents (e.g., Cymbalta, Zofran, Elidel and others).

J. Drug-specific PA Forms- Drug-specific PA forms contain medical necessity documentation requirements and/or criteria that may not be repeated in the PDL. Drug-specific PA forms may be obtained on the web at www.mainearepdl.org.

K. PA Exemptions for Prescribers- According to MaineCare Benefits Manual Chapter II (80.07-4), providers may receive a three (3) month exemption from prior authorization requirement for certain categories of drugs when they demonstrate high compliance with the Department's PDL. The Department will notify providers in writing which drug categories are included and what dates apply to the exemption. If a provider loses his/ her exemption, members who previously were not required to obtain a PA while the prescriber was exempt will be required to do so, and criteria for approval of that medication will need to be met.

L: Drug-Drug Interactions (DDI)- The DUR Committee has implemented new drug-drug interaction edits requiring prior authorization. Several drug-drug combinations and PDL drug categories are affected by new PA requirements. These will be indicated in the PDL with DDI notation. Please see the DDI document provided in the PDL.

ASSORTED ANTIBIOTICS

| | | | | | | |
|------------------------------------|-------------|---------------------------------|--------|--------------------------------|---|--|
| BETA-LACTAMS / CLAVULANATE COMBO'S | MC/DEL | AMOXICILLIN | MC/DEL | AUGMENTIN ³ | 3. Chewable 125mg & 250mg and Solution 125mg/5ml and 250mg/5ml available without PA. 4. Use preferred generic amoxicillin/clavulanate potassium alternatives. Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Ampicillin will now be non-preferred and require prior authorization if it is currently being used in combination with either Prevacid, pantoprazole, Prilosec, or any currently non preferred PPI. |
| | MC/DEL | AMOXICILLIN/POTASSIUM CLA CHEW | MC/DEL | AUGMENTIN XR TB12 ⁴ | | |
| | MC/DEL | AMOXICILLIN/POTASSIUM CLA SUSR | | | | |
| | MC/DEL | AMOXICILLIN/POTASSIUM CLA TABS | | | | |
| | MC/DEL | AMPICILLIN | | | | |
| | MC | BICILLIN L-A SUSP | | | | |
| | MC/DEL | DICLOXACILLIN SODIUM CAPS | | | | |
| | MC | OXACILLIN SODIUM SOLR | | | | |
| | MC/DEL | PENICILLIN V POTASSIUM | | | | |
| | MC | TIMENTIN SOLR | | | | |
| MC | UNASYN SOLR | | | | | |
| MC/DEL | ZOSYN | | | | | |
| CEPHALOSPORINS | MC/DEL | CEFADROXIL HEMIHYDRATE | MC | CEDAX | 1. Both brand and generic are clinically non-preferred. 2. Dosing limits apply, please see Dosage Consolidation List. 3. Approvals will only be considered for patients 18 years of age or older who have limited or no alternative | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Vantin will now be non-preferred and require prior authorization if it is currently being used in combination with either Prevacid, pantoprazole, Prilosec, or any currently non |
| | MC/DEL | CEFAZOLIN SODIUM SOLR | MC/DEL | CEFACTOR ¹ | | |
| | MC/DEL | CEFDINIR | MC/DEL | CEFADROXIL MONOHYDRATE TABS | | |
| | MC/DEL | CEFEPIME | MC/DEL | CEFIXIME SUS | | |
| | MC/DEL | CEFPODOXIME | MC/DEL | CEPHALEXIN TABS | | |
| | MC/DEL | CEFPODOXIME PROXETIL SUS | MC | CEPHALEXIN 750MG CAPS | | |
| | MC/DEL | CEFPODOXIME PROXETIL TAB | MC/DEL | CEFTIN | | |
| | MC/DEL | CEFIXIME 400MG ² CAP | MC | DAXBIA | | |
| | MC/DEL | CEFPROZIL | MC | FETROJA ³ | | |

| | | | | | | | |
|---|--|--|---|--|--|--|--|
| | | | | | | | Arikayce will require clinical PA to confirm MAC lung disease and for use in adults who have limited or no alternative treatment options. Zemdri will be reserved for patients with limited or no alternative treatment of care. |
| ANTI-MYCOBACTERIALS / ANTI-TUBERCULOSIS | MC/DEL MC/DEL MC/DEL MC/DEL | | ETHAMBUTOL HCL TABS MYAMBUTOL TABS RIFABUTIN CAPS RIFAMPIN | MC/DEL MC/DEL MC | | MYCOBUTIN CAPS PRETOMANID RIFADIN CAPS | Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Pretomanid is indicated as part of a combination regimen with bedaquiline and linezolid for the treatment of adults with pulmonary extensively drug resistant (XDR) or treatment-intolerant or non-responsive multidrug-resistant (MDR) tuberculosis (TB). Approval of this indication is based in limited clinical safety and efficacy data. This drug is indicated for use in a limited and specific population of patients. DDI: Preferred rifampin will be non-preferred and require prior authorization if it is currently being used in combination with either Pradaxa or Latuda. |
| ANTIMALARIAL AGENTS | MC/DEL MC MC/DEL MC/DEL | | DARAPRIM TABS KRINTAFEL ² MEFLOQUINE HCL TABS QUININE SULFATE | MC MC/DEL MC/DEL MC MC MC/DEL | | ARALEN TABS CHLOROQUINE PHOSPHATE TABS ³ HYDROXYCHLOROQUINE TABS ³ ISONARIF ¹ MALARONE TABS PLAQUENIL TABS | Use PA Form# 20420 1. Ingredients available as preferred without PA. 2. Krintafel is preferred for ≥ 16 years of age. 3. Established users will be grandfathered Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Avoid coadministration of Krintafel® with Organic Cation Transporter 2 (OCT2) and Multidrug and Toxin Extrusion (MATE) substrates (e.g. dofetilide, metformin). |
| ANTHELMINTICS | MC/DEL MC/DEL MC/DEL | | ALBENDAZOLE PRAZIQUANTEL TAB STROMECTOL TABS | MC MC MC/DEL | | ALBENZA TABS EMVERM BILTRICIDE TABS | Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ANTIBIOTICS - MISC. | MC MC MC MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC | | AZACTAM SOLR COLY-MYCIN-M SOLR COLISTIMETHATE SODIUM SOLR FIRVANQ ⁴ FUROXONE TABS METRONIDAZOLE ¹ PENTAMIDINE ISETHIONATE SOLR SOLOSEC TRIMETHOPRIM TABS VANCOMYCIN 5GM INJ. VANCOMYCIN CAPS XIFAXAN 200mg | MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC MC | | AEMCOLO COLISTIMETHATE SODIUM SOLR CAYSTON ³ FLAGYL CAPS FLAGYL TABS FLAGYL ER TBCR KETEK LIKMEZ METRONIDAZOLE 375MG CAPS ¹ METRONIDAZOLE 750MG TABS ¹ NEBUPENT SOLR REBYOTA ⁵ TINDAMAX VANCOMYCIN 10GM INJ. ² XENLETA XIFAXAN VOWST ⁵ | 1. 375mg caps and 750mg tabs are non-preferred. Please use available preferred strengths(250mg & 500mg tabs) to obtain required dose without PA. 2. Please use multiple 5gm which are preferred to obtain dose without PA. 3. Clinical PA is required to establish CF diagnosis and medical necessity. Prior trial and failure of preferred Tobi before approval will be granted. 4. Quantity limit of one per 150ml bottle. 5. For the treatment of patients 18 years of age and older. Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. 1. For macrolide resistant infections when quinolones inappropriate DDI: Ketek is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with either Enblex 15mg or Vesicare 10mg or carbamazepine. Cayston is only indicated to improve respiratory symptoms in CF patients with Pseudomonas aeruginosa. Dosing limits, as should be given TID X28 days (followed by 28 days OFF Cayston therapy). A bronshodilator should be used before administration of Cayston. Xenleta will be considered for the treatment of adults with community-acquired bacterial pneumonia (CABP) caused by the following susceptible microorganisms: Streptococcus pneumoniae, Staphylococcus aureus (methicillin-susceptible isolates), Hemophilus influenzae, Legionella pneumophila, Mycoplasma pneumoniae, and Chlamydia pneumoniae. Vowst: To prevent the recurrence of Clostridioides difficile infection (CDI) in individuals 18 years of age and older following antibacterial treatment for recurrent CDI (rCDI). Likmez: patient has a medical necessity for a non-solid oral dosage form. Rebyota: For the prevention of recurrence of Clostridioides difficile infection (CDI) in individuals 18 years of age and older following antibiotic treatment for recurrent CDI. The limitation of use is that Rebyota® is not indicated for treatment of CDI. |
| CARBAPENEMS | | | | MC MC MC/DEL MC/DEL | | INVANZ SOLR MERREM SOLR PRIMAXIN RECARBRIO | Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |

| | | | | | | | |
|--|--|--|--|---|--|--|---|
| LINCOSAMIDES / OXAZOLIDINONES / LEPROSTATICS | MC/DEL MC/DEL MC/DEL MC MC/DEL | CLEOCIN SOLN CLEOCIN SUSR CLINDAMYCIN HCL 150CAPS DAPSONE TABS LINEZOLID 600mg TABS | MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL | 8 8 8 8 8 9 9 | CLEOCIN CAPS CLINDAMYCIN HCL 300CAPS ¹ SIVEXTRO VIBATIV LINEZOLID TABS ZYVOX SUSR ZYVOX TABS | 1. Use multiple 150's for Clindamycin instead of 300's. 2. Quantity limit of 14 days supply within a 60day period. Use PA Form# 30820 for Zyvox & Vibativ Use PA Form# 20420 for all others | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. For Zyvox or Vibativ, please see the criteria listed in the Antibacterial Antibiotics PA form. |
| ANTI INFECTIVE COMBO'S - MISC. | MC/DEL MC/DEL MC/DEL MC/DEL | ERYTHROMYCIN/SULF SUSR SEPTRA/DS TABS SULFAMETHOXAZOLE/TRIMETH TRIMETHOPRIM/SULFAMETHOXA | MC MC | | BACTRIM DS TABS VABOMERE ¹ | Use PA Form# 20420 1. For the treatment of patients ≥ 18 years of age. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ANTIPROTOZOALS | MC/DEL MC/DEL | BENZNIDAZOLE ² LAMPIT ² | MC | | ALINIA ¹ | 1. Alina is preferred for children less than 12 years of age. 2. Clinical PA required for appropriate diagnosis. Use PA Form# 20420 | Benznidazole is indicated for pediatric patients 2 to 12 years of age for the treatment of Chagas disease (American trypanosomiasis) caused by Trypanosoma cruzi. |
| ANTI - FUNGALS | | | | | | | |
| ANTIFUNGALS - ASSORTED | MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | ANCOBON CAPS FLUCONAZOLE ¹ KETOCONAZOLE TABS ⁷ NYSTATIN TERBINAFINE TABS ⁴ VORICONAZOLE TABS | MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC | 6 6 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 | LAMISIL TABS ⁴ ITRACONAZOLE BREXAFEMME CRESEMBA ⁹ GRIFULVIN V TABS GRISEOFULVIN SUSP GRISEOFULVIN ULTRAMICROSI TABS GRIS-PEG TABS REZZAYO ⁹ SPORANOX SOLN ² SPORANOX PULSEPAK CAPS ³ SPORANOX CAPS ³ DIFLUCAN ERAXIS INJ ⁶ GRIFULVIN SUSP ONMEL NOXAFIL ⁵ TOLSURA VFEND TABS VIVJOA | See quantity limit table. Non-preferred products must be used in specified step order. Continue to use Anti-Fungal PA form for non-preferred products. 1. QL--1/every 7-day period (150mg only). 2. Sporanox QL 300cc/month with PA. See quantity limit table. 3. Sporanox QL 30/month with PA. 4. Quantity limit of one tablet daily. Please see dosage consolidation list. 5. Approved if immuno suppressed/ HIV or if the member has failed a 7 day trial of a preferred antifungal therapy. 6. Eraxis will be approved if submitting with documentation that it was initiated during a hospitalization and this request is to finish the hospital course. 7. Quantity limits allowing 30 day supply without PA. PA will be required if using > 30 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. The other criteria are listed on the Antifungal PA form including the required proof of a non-cosmetic fungal infection. DDI: Any Griseofulvin will now be non-preferred and require prior authorization if it is currently being used in combination with either Prevacid, pantoprazole, Prilosec, or any currently non preferred PPI. DDI: Sporanox is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for current use with Enablex 15mg, Vesicare 10mg, Prandin, Prevacid, pantoprazole, Prilosec, or any currently non preferred PPI, due to a significant drug-drug interaction. DDI: Vfend is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with Warfarin. DDI: Fluconazole (except 150mg strength) will now be non-preferred and require prior authorization if it is currently being used with glimepiride (Amaryl), Enablex 15mg, or Vesicare 10mg. Diflucan is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with either glimepiride (Amaryl), Enablex 15mg, or Vesicare 10mg. DDI: Fluconazole will require prior authorization if being used in combination with Plavix or Warfarin. DDI: Ketoconazole will now be non-preferred and require prior authorization if they are currently being used in combination with any of the following medications: Prevacid, Pantoprazole, Plavix, Onglyza, Enablex 15mg, Vesicare 10mg, Latuda, Cometriq, Tafilnar or Ormeprazole. Rezzayo: In patients 18 years of age or older who have limited or no alternative options for the treatment of candidemia and invasive candidiasis. |

| ANTI - VIRALS | | | | | | |
|-------------------------|--------|--|--------|---|--------------------------|--|
| ANTIRETROVIRALS | MC/DEL | ABACAVIR TABS | MC/DEL | 8 | ABACAVIR SOL | days. |
| | MC | APRETUDE | MC/DEL | 8 | APTIVUS | 8. For children < 18, quantity limits allows 8 weeks supply without PA. PA will be required if using > than 8 weeks. If 18 and older PA will be required for any quantity. Not approving for Onychomycosis indication. |
| | MC/DEL | ATAZANAVIR | MC/DEL | 8 | CIMDUO | 9. For patients ≥ 18years of age |
| | MC | ATRIPLA ¹ | MC/DEL | 8 | COMBIVIR TABS | Use PA Form# 10120 |
| | MC | BIKTARVY | MC/DEL | 8 | EDURANT | |
| | MC | CABENUVA | MC/DEL | 8 | EPZICOM ¹ | |
| | MC | COMPLERA ¹ | MC/DEL | 8 | FUZEON | |
| | MC/DEL | DELSTRIGO | MC/DEL | 8 | INTELENCE | |
| | MC | DESCOVY ¹ | MC/DEL | 8 | ISENTRESS ³ | |
| | MC | DIDANOSINE | MC/DEL | 8 | ISENTRESS HD | |
| | MC/DEL | DOVATO | MC | 8 | JULUCA | |
| | MC | EFAVIRENZ TAB | MC | 8 | KALETRA | |
| | MC/DEL | EFAVIRENZ CAP | MC/DEL | 8 | LAMIVUDINE SOLN | |
| | MC | EFAVIRENZ-EMTRICITABINE-TENOFOVIR DF TAB | MC/DEL | 8 | LEXIVA | |
| | MC | EMTRICITABINE-TENOFOVIR | MC/DEL | 8 | NEVIRAPINE | |
| | MC | EMTRIVA ¹ | MC | 8 | NORVIR | |
| | MC | EPIVIR SOL | MC/DEL | 8 | PIFELTRO | |
| | MC/DEL | EVOTAZ ¹ | MC | 8 | RETROVIR | |
| | MC | GENVOYA ^{1,5} | MC | 8 | REYATAZ | |
| | MC/DEL | ISENTRESS 400MG ⁶ | MC/DEL | 8 | SELZENTRY | |
| | MC/DEL | ISENTRESS CHEW ³ | MC | 8 | STAVUDINE | |
| | MC/DEL | ISENTRESS POWDER | MC | 8 | STRIBILD ¹ | |
| | MC/DEL | LAMIVUDINE TABS | MC | 8 | SUNLENCA ⁵ | |
| | MC/DEL | LAMIVUDINE/ZIDOVUDINE | MC/DEL | 8 | SYMFI ⁵ | |
| | MC/DEL | LOPINAVIR-RITONAVIR SOL | MC/DEL | 8 | SYMFI LO ⁵ | |
| | MC | LOPINAVIR-RITONAVIR TAB | MC/DEL | 8 | SYM TUZA | |
| | MC | ODEFSEY ¹ | MC | 8 | TRIUMEQ ^{1,4} | |
| | MC/DEL | PREZCOBIX | MC/DEL | 8 | TRIZIVIR TABS | |
| | MC | PREZISTA ² | MC | 8 | TRUVADA ¹ | |
| | MC/DEL | RITONAVIR TAB 100MG | MC/DEL | 8 | VIRACEPT TABS | |
| | MC | RUKOBIA ⁹ | MC | 8 | VITEKTA | |
| | MC | SUSTIVA ¹ | MC | 8 | ZERIT | |
| | MC | TIVICAY | MC | 8 | VIDEX EC | |
| | MC | TIVICAY PD | MC | 8 | VIREAD TABS ¹ | |
| | MC | TROGARZO ⁵ | MC/DEL | 8 | ZIAGEN TABS | |
| | MC | TYBOST | MC/DEL | 8 | ZIAGEN SOL | |
| | MC | VIREAD POW | MC/DEL | 9 | VIRAMUNE XR | |
| | MC/DEL | ZIDOVUDINE | | | | |
| CYTO-MEGALOVIRUS AGENTS | MC | CIDOFOVIR | MC | | VALCYTE TABS | Use PA Form# 20420 |

| | | | | | | | | |
|--------------------------|--|--|--|--|-----------------------|---|---|--|
| | MC MC/DEL MC/DEL | | FOSCARNET SODIUM GANCICLOVIR VALGANCICLOVIR | MC/DEL MC/DEL MC/DEL | | FOSCAVIR LIVTENCITY ¹ PREVYMIS | 1. Must show failure or contraindication to all the following ganciclovir, valganciclovir, cidofovir and foscarnet before Livtency will be approved. | on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Prevymis: Documentation that member is high-risk for CMV reactivation as defined by transplant guidelines or that there has been significant myelosuppression by one of the preferred agents. DDI: Livtency is a substrate of CYP3A4. Coadministration of Livtency® with strong inducers of CYP3A4 is not recommended, except for selected anticonvulsants. |
| HERPES AGENTS | MC/DEL MC/DEL | | ACYCLOVIR VALACYCLOVIR HCL | MC/DEL MC MC/DEL MC MC/DEL | 8 8 8 8 9 | FAMCICLOVIR ¹ SITAVIG ZOVIRAX ¹ VALTRES TABS ¹ FAMVIR TABS ¹ | 1. Must fail Acyclovir and Valacyclovir before non-preferred products in step order. Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| INFLUENZA AGENTS | MC MC MC/DEL | | AMANTADINE CAPS RELENZA DISKHALER AEPB OSELTAMIVIR ¹ | MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL | | AMANTADINE TABS FLUMADINE TABS FLUMIST RIMANTADINE HCL TABS TAMIFLU ¹ TAMIFLU SUS XOFLUZA | 1. Tamiflu and Osetamivir 10 caps or 60cc's per month. Will be audited for presence of positive influenza tests in patient or family member. Use PA Form# 20420 for all others | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| IMMUNE SERUMS | | | | | | | | |
| IMMUNE SERUMS | MC | | HYPERRHO INJ | | | | | |
| HEPATITIS AGENTS | | | | | | | | |
| HEPATITIS C AGENTS | MC MC MC/DEL MC/DEL MC/DEL MC MC/DEL | | SOFOSBUVIR/VELPATASVIR ² (Authorized generic labeler 72626 Asegua Therapeutics) MAVYRET ² PEGASYS KIT ¹ PEGASYS SOLN PEG-INTRON KIT ¹ RIBAVIRIN RIBASPHERE | MC/DEL MC/DEL MC MC/DEL MC MC MC MC MC/DEL | | COPEGUS TABS DAKLINZA EPCLUSA ² HARVONI ² REBETOL CAPS RIBAPAK SOVALDI ² VIEKIRA PAK ² VIEKIRA XR ² VOSEVI ZEPATIER ² | 1. Dosing limits apply, please see dosage consolidation list. 2. Approvals will require clinical PA. Please see the Hepatitis PA form for criteria Use PA Form #10700 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Olysio will require a prior authorization if it is currently being used in combination with drugs known to be significant CYP3A4 inhibitors (ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, nelfinavir, ritonavir, atazanavir, saquinavir and telithromycin). |
| HEPATITIS AGENTS - MISC. | | | | MC | | ACTIMMUNE | Use PA Form# 20420 | Approved for chronic granulomatous disease, osteopetrosis and idiopathic pulmonary fibrosis. |
| HEPATITIS B ONLY | MC/DEL MC | | ENTECAVIR TENOFVIR | MC MC MC MC | | BARACLUDE HEPSERA TABS TYZEKA VEMLIDY | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Baraclude is indicated for treatment of chronic Hep B virus (HBV) in adults with: evidence of active viral replication AND either evidence of persistent elevation in serum aminotransferases (ALT or AST) or histologically active disease, Patient is 16 years of age or older. Boxed warning: Use not recommended for those co-infected with HIV and HBV who are not also receiving highly active antiretroviral therapy (HAART). Vemlidy® remain non-preferred and require prior authorization and be available to those who have evidence of bone loss or renal insufficiency or who are unable to tolerate or who have failed on preferred medications. |
| RSV PROPHYLAXIS | | | | | | | | |
| RSV PROPHYLAXIS | | | | MC | | SYNAGIS ¹ | Use PA Form# 30120 1. MaineCare will approve Synagis PA's for start date of | Please see the criteria listed on the Synagis PA form. |

STEROIDS

| | | | | | | | | |
|--|--------|------------------|------------------------------|--------|--|--------------------------|------------------------------------|--|
| GLUCOCORTICOIDS/ MINERALOCORTICOIDS | MC/DEL | | BUDESONIDE EC 3mg DR CAPS | MC | | ALKINDI SPRINKLE | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | MC | | CELESTONE SUSP | MC | | CORTEF 10 and 20 TABS | | |
| | MC/DEL | | CORTEF 5 | MC/DEL | | FLORINEF TABS | | |
| | MC/DEL | | CORTISONE ACETATE TABS | MC | | HEMADY | | |
| | MC/DEL | | DELTASONE TABS | MC/DEL | | MEDROL TABS | | |
| | MC/DEL | | DEPO-MEDROL SUSP | MC | | MEDROL DOSEPAK TABS | | |
| | MC/DEL | | DEXAMETHASONE | MC | | MILLIPRED | | |
| | MC | | DEXPAK | MC | | ORTIKOS | | |
| | MC/DEL | | FLUDROCORTISONE ACETATE TABS | MC | | ORAPRED SOLN | | |
| | MC/DEL | | HYDROCORTISONE | MC | | PEDIAPRED LIQD | | |
| | MC | | KENALOG | MC | | PREDNISONE INTENSOL CONC | | |
| | MC/DEL | | METHYLPREDNISOLONE TABS | MC | | STERAPRED TABS | | |
| | MC/DEL | | PREDNISOLONE | MC | | ZILRETTA | | |
| | MC/DEL | | PREDNISONE | | | | | |
| | MC/DEL | | SOLU-CORTEF SOLR | | | | | |
| MC/DEL | | SOLU-MEDROL SOLR | | | | | | |

DDI: All preferred steroids will require clinical PA for patients over 60 that are currently on fluoroquinolone therapy.

HORMONE REPLACEMENT THERAPIES

| | | | | | | | | |
|-----------------------|--------|--|---------------------|--------|--------------------------|------------------------|------------------------------------|--|
| ANDROGENS / ANABOLICS | MC/DEL | | ANDRODERM PT24 | MC | | ANADROL-50 | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Additionally, laboratory evidence of a testosterone deficiency must be supplied. One of each dosage form should be tried (tablet, injection, and topical) |
| | MC/DEL | | ANDROGEL 1% | MC | | ANDRO LA 200 OIL | | |
| | MC/DEL | | ANDROGEL PUMP 1.62% | MC/DEL | | ANDROGEL PACKETS 1.62% | | |
| | MC/DEL | | DANAZOL CAPS | MC | | ANDROID CAPS | | |
| | MC/DEL | | TESTOSTERONE CYP | MC | | AXIRON | | |
| | | | | MC | | DELATESTRYL OIL | | |
| | | | | MC/DEL | | DEPO-TESTOSTERONE OIL | | |
| | | | | MC | | FORTESTA | | |
| | | | | MC | | HALOTESTIN TABS | | |
| | | | | MC/DEL | | JATENZO | | |
| | | | | MC/DEL | | METHITEST TAB | | |
| | | | | MC/DEL | | METHYLTESTOSTERONE CAP | | |
| | | | | MC/DEL | | OXANDROLONE | | |
| | | | | MC/DEL | | STRIANT MUC ER | | |
| | | | | MC | | TESTIM | | |
| | | | MC/DEL | | TESTOSTERONE GEL PACKETS | | | |
| | | | MC/DEL | | TESTOSTERONE SOL | | | |
| | | | MC | | TESTRED CAPS | | | |
| | | | MC | | TLANDO | | | |
| | | | MC/DEL | | VOGELXO | | | |
| | | | MC/DEL | | XYOSTED | | | |

Oxandrolone: Weight gain (adjunctive therapy): Adjunctive therapy to promote weight gain after weight loss following extensive surgery, chronic infections, or severe trauma, and in some patients who, without definite pathophysiologic reasons, fail to gain or to maintain normal weight. Other indications included in manufacturer labeling: Adjunctive therapy to offset protein catabolism with prolonged corticosteroid administration. Requirement for documentation of weight loss over two readings- Patient has involuntary weight loss of more than 10% of total body weight in less than four months) and, BMI < 18.5 (Normal BMI = 18.5 to 24.9)

| | | | | | | | | |
|-------------------------------|--------|--|-----------------|--------|---|-----------------------|------------------------------------|---|
| ESTROGENS - PATCHES / TOPICAL | MC | | EVAMIST | MC/DEL | 5 | ESTRADIOL PTWK | Use PA Form# 20420 | Approved for failures on multiple oral estrogen agents after 90 day trials or if unable to swallow any oral medication. |
| | MC/DEL | | MINIVELLE PATCH | MC/DEL | 8 | DIVIGEL ¹ | | |
| | | | | MC/DEL | 8 | CLIMARA PTWK | | |
| | | | | MC/DEL | 8 | ELESTRIN ¹ | | |
| | | | | MC/DEL | 8 | MENOSTAR PATCH | | |
| | | | | MC/DEL | 8 | VIVELLE-DOT PTTW | | |

| | | | | | | | | |
|------------------|--------|--|---------------|--------|----------------|-------------------------|------------------------------------|---|
| ESTROGENS - TABS | MC/DEL | | ESTRADIOL | MC/DEL | | ENJUVA | Use PA Form# 20420 | Preferred drugs must be tried for at least 90 days and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | MC/DEL | | PREMARIN TABS | MC/DEL | | ESTRADIOL-NORETHINDRONE | | |
| | | | | MC/DEL | | ESTRACE TABS | | |
| | | | | MC | | ESTRATAB TABS | | |
| | | | | MC/DEL | | MENEST TABS | | |
| | | | | MC/DEL | | NORETHINDRON-ETHINYL | | |
| | | | MC | | ORTHO-EST TABS | | | |

| | | | | | | | |
|--|--|---|--|--|--|---|---|
| | MC MC MC/DEL MC/DEL MC | TRI-LO-ESTARYLLA TAB TRI-ESTARYLLA TRI-SPRINTEC TAB TRI-LO-SPRINTEC TRINESSA | | | | | If member experienced adverse reactions, consider using Oral Contraceptives from other groups. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with Tracleer. |
| CONTRACEPTIVES - MULTI-PHASIC COMBINATIONS | | | MC | | NATAZIA | Use PA Form# 20420 Use PA Form# 20420 | |
| VASOMOTOR SYMPTOMS AGENTS | | | | | | | |
| VASOMOTOR SYMPTOMS AGENTS | | | MC/DEL | | VEOZAH | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Avoid concomitant use of Veozah with drugs that are weak, moderate or strong CYP1A2 inhibitors. Veozah: Approval requires at least one preferred Hormone Replacement Therapy (HRT) and two preferred non-hormonal therapies (i.e., SSRIs, SNRIs, gabapentin, pregabalin, clonidine). |
| DIABETES SUPPLIES | | | | | | | |
| DIABETIC - SUPPLIES | | CONTINUOUS GLUCOSE MONITORING ^{1,2} DIABETIC- LANCETS DIABETIC- LANCING DEVICES DIABETIC- LANCING DEVICES DIABETIC- PEN NEEDLES DIABETIC- SYRINGES DIABETIC- TEST STRIPS DIABETIC- METERS | | | | Use PA Form#20420 | 1. Clinical PA is required to establish diagnosis and medical necessity. 2. Dosing limits apply. Please refer to Dose consolidation list. Please refer to the MaineCare Preferred Diabetic Supply List available at www.mainearepdl.org Continuous Glucose Monitoring Criteria: Patient has a diagnosis of Diabetes Mellitus AND Practitioner feels patient has sufficient training to use CGM • 2 years of age or older for Dexcom G6 and Dexcom G7, ≥ 14 years for Medtronic Guardian, or ≥ 4 years for Freestyle Libre 2. • At least one of the following are documented: o Hypoglycemic unawareness o Treated with insulin (at least 1X day) o Has history of problematic hypoglycemia with documentation of at least one recurrent level 2 hypoglycemic events, or 1 level 3 hypoglycemic event • Approval of non-preferred products will be limited to cases where the CGM is directly integrated with the patient's insulin pump. The make and model of pump must be documented on the prior authorization. |
| DIABETES THERAPIES | | | | | | | |
| DIABETIC - INSULIN | MC/DEL MC MC MC MC MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | APIDRA HUMALOG KWIKPEN INJ 100/ML HUMALOG JUNIOR KWIKPEN 100/ML HUMALOG MIX 75/25 HUMALOG 50/50 VIAL HUMULIN INJ 70/30 KWIKPEN HUMULIN INJ 70/30 HUMULIN R INJ U-500 INSULIN ASPART PROT MIX 70-30 INSULIN ASPART INSULIN LISPRO LANTUS SOLN LEVEMIR NOVOLOG NOVOLOG MIX NOVOLOG MIX 70/30 FLEXPEN | MC/DEL MC/DEL MC MC/DEL MC MC MC MC MC/DEL MC | | ADMELOG AFREZZA ¹ BASAGLAR FIASP HUMALOG KWIKPEN U-200 HUMULIN INJ 50/50 HUMULIN N INJ U-100 HUMULIN R U-100 LYUMJEV NOVOLIN RELION | Use PA Form# 20420 1. Not to be as a monotherapy. Obtain lab values of pulmonary function and recent smoking history 2. For the treatment of patients ≥3 years of age | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| DIABETIC - PENFILLS | MC MC | HUMALOG MIX KWIK 50/50 HUMALOG MIX INJ 75/25 KWP | MC MC/DEL | | APIDRA OPTICLIK PEN NOVOLIN 70/30 PEN | | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |

| | | | | | | |
|--|--|--|--|---|---|---|
| | <p>MC MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL</p> | <p>HUMALOG KWIK INJ 100/ML HUMALOG KWIK INJ 200/ML HUMULIN R U-500 KWP INSULIN ASPART PROT MIX 70-30 PEN INSULIN ASPART PEN INSULIN LISPRO KWIKPEN U-100 LANTUS SOLOSTAR LEVEMIR FLEXTOUCH LEVEMIR FLEXPEN NOVOLOG MIX PENFILL NOVOLOG PENFILL SOLN NOVOLOG FLEXPEN NOVOLOG MIX 70/30 VIAL TOUJEO MAX SOLOSTAR TOUJEO SOLOSTAR</p> | <p>MC MC/DEL</p> | <p>REZVOGLAR KWIKPEN TRESIBA</p> | <p>another drug and the preferred drug(s) exists.</p> | |
| DIABETIC - DPP- 4 ENZYME INHIBITOR | <p>MC/DEL MC/DEL</p> | <p>JANUVIA^{1,2} TRADJENTA²</p> | <p>MC/DEL MC/DEL MC/DEL MC</p> | <p>NESINA ONGLYZA² QTERN ZITUVIO</p> | <p>1. Preferred if therapeutic doses of metformin are seen in members drug profile for at least 60 days within the past 18 months or if phosphate binder is currently seen in the members drug profile.</p> <p>2. Dosing limits apply. Please refer to Dose consolidation list.</p> <p>Use PA Form# 20420</p> | <p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>DDI: Onglyza 5mg will require a prior authorization if it is currently being used in combination with drugs known to be significant CYP3A4 inhibitors (ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, nelfinavir, ritonavir, atazanavir, saquinavir and telithromycin).</p> |
| DIABETIC - DPP- 4 ENZYME INHIBITOR-COMBO | <p>MC/DEL MC/DEL MC/DEL</p> | <p>JANUMET^{1,2} JANUMET XR^{1,2} JENTADUETO¹</p> | <p>MC/DEL MC/DEL MC MC/DEL</p> | <p>JENTADUETO XR KAZANO KOMBIGLYZE XR OSENI</p> | <p>1. Preferred if therapeutic doses of metformin are seen in members drug profile for at least 60 days within the past 18 months or if phosphate binder is currently seen in the members drug profile.</p> <p>2. Dosing limits apply. Please refer to Dose consolidation list.</p> <p>Use PA Form# 20420</p> | |
| DIABETIC - LANCET-LANCET DEVICE | | | | | <p>Use PA Form# 20420</p> | Please refer to the MaineCare Preferred Diabetic Supply List available at www.mainearepdl.org |
| DIABETIC - SYRINGES-NEEDLES | | | | | <p>Use PA Form# 20420</p> | Please refer to the MaineCare Preferred Diabetic Supply List available at www.mainearepdl.org |
| DIABETIC - OTHER | | | <p>MC/DEL MC</p> | <p>CYCLOSET SYMLIN</p> | <p>Use PA Form #20420 for all others</p> | |
| SGLT 2 INHIBITORS | <p>MC/DEL MC/DEL MC/DEL</p> | <p>FARXIGA INVOKANA¹ JARDIANCE</p> | <p>MC/DEL</p> | <p>STEGLATRO</p> | <p>1.Dosing limits apply please refer to Dose Consolidation List</p> <p>Use PA Form# 20420</p> | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |

| | | | | | | | | |
|-------------------------------|--|--|--|--|----------------------------|--|--|--|
| SGLT 2 INHIBITOR COMBINATIONS | MC/DEL MC/DEL MC/DEL | | INVOKAMET SYNJARDY XIGDOU XR | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | | GLYXAMBI INVOKAMET XR SEGLUROMET STEGLUJAN SYNJARDY XR TRIJARDY XR | | Preferred drugs must be tried for at least 3 months at full therapeutic doses and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Glyxambi /Xigduo XR- Verify prior trials and failures or intolerance of preferred treatments from other diabetic categories Synjardy® XR is not recommended for patients with type 1 DM or for the treatment of diabetic ketoacidosis. Use PA Form# 20420 |
| DIABETIC MONITOR | MC MC MC MC | | ONE TOUCH ULTRA 2 KIT ONE TOUCH ULTRA MINI KIT TRUE METRIX TRUETRACK | MC MC MC MC MC MC MC MC MC | | ACCUCHECK ASCENSIA ASSURE CONTOUR BREEZE Z EXACTECH FREESTYLE INSULINX FREESTYLE LITE SYSTEM KIT ONE TOUCH ULTRA SMART KIT PRECISION XTRA METER PRODIGY | Use PA Form# 20420 | Effective October 25th 2007, approvals for all non preferred meters/ test strips will require medical necessity documenting clinically significant features that are not available on any of the preferred meters. |
| DIABETIC TEST STRIPS | MC MC MC | | ONE TOUCH ULTRA ¹ TRUE METRIX TRUETRACK | MC MC MC MC MC MC MC MC MC MC | | ACCUCHECK ASCENSIA ASSURE CONTOUR BREEZE Z EXACTECH FREESTYLE FREESTYLE LITE FREESTYLE INSULINX ONE TOUCH DELICA PRECISION XTRA PRODIGY | 1. Only 50 ct & 100 ct package size. Use PA Form# 20420 | Effective October 25th 2007, approvals for all non preferred meters/ test strips will require medical necessity documenting clinically significant features that are not available on any of the preferred meters. |
| INCRETIN MIMETIC | MC MC MC/DEL | | BYETTA TRULICITY VICTOZA | MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL | 5 5 8 8 8 8 | OZEMPIC RYBELSUS ADLYXIN BYDUREON BCISE MOUNJARO SOLIQUA XULTOPHY | | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Soliqua must try both insulin and a preferred incretin mimetic and have a medical necessity for use that is not based on convenience or simply due to the fact that one injection is needed instead of two. Use PA Form# 20420 |
| DIABETIC - ORAL SULFONYLUREAS | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | | CHLORPROPAMIDE TABS GLIMEPIRIDE GLIPIZIDE TABS GLIPIZIDE ER TABS GLYBURIDE MICRONIZED TABS GLYBURIDE TABS ¹ TOLAZAMIDE TABS TOLBUTAMIDE TABS | MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL | | AMARYL TABS DIABETA TABS GLUCOTROL TABS GLUCOTROL XL TBCR GLYNASE TABS MICRONASE TABS | Use PA Form# 20420 1. Pa required for members ≥65. Glyburide has a greater risk of severe prolonged hypoglycemia in older adults. | Preferred drugs must be tried for at least 3 months at full therapeutic doses and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: All sulfonylureas (except glyburide) will now be non-preferred and require prior authorization if it is currently being used with either ranitidine or cimetidine. DDI: Glimepiride will now be non-preferred and require prior authorization if it is currently being used with either fluconazole (except 150mg strength) or fluvoxamine. Amaryl is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with either fluconazole or fluvoxamine. |
| DIABETIC -ORAL BIGUANIDES | MC/DEL | | METFORMIN HCL TABS | MC | | GLUCOPHAGE TABS | Use PA Form# 20420 | Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered |

| | | | | | | | |
|--------------------------------------|--------|---|------------------------------------|------------------------------|---|---|--|
| | MC/DEL | | METFORMIN ER | MC MC MC/DEL | GLUCOPHAGE XR TB24 FORTAMET METFORMIN ER OSMOTIC | | on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| DIABETIC - THIAZOL / BIGUANIDE COMBO | | | | MC/DEL MC/DEL MC MC | ACTOPLUS MET ¹ ACTOPLUS MET XR AVANDARYL ¹ AVANDAMET TABS ¹ | Use PA Form# 20420 1. Requires use of Actos, Metformin, or other preferred anti-diabetics. | DDI: Actos, Avandia, or any combination product with Actos or Avandia will now be non-preferred and require prior authorization if it is currently being used with gemfibrozil. |
| DIABETIC - / THIAZOL | MC/DEL | | PIOGLITAZONE HCL ¹ | MC/DEL MC | ACTOS TABS ³ AVANDIA TABS ² | 1. Pioglitazone HCL is non-preferred as monotherapy. Pioglitazone HCL is preferred if therapeutic doses of metformin, sulfonylurea or insulin are seen in members drug profile for at least 60 days within the past 18 months. 2. Current users of Avandia who have tried Actos will be able to continue use of Avandia. 3. Dosing limits apply please refer to Dose Consolidation List Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Actos, Avandia, or any combination product with Actos or Avandia will now be non-preferred and require prior authorization if it is currently being used with gemfibrozil. |
| DIABETIC - ALPHAGLUCOSIDASE | MC/DEL | | | MC | PRECOSE TABS | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| DIABETIC - SULFONYLUREA / BIGUANIDE | MC/DEL | | GLYBURIDE/METFORMIN | MC MC MC/DEL | GLUCOVANCE TABS ¹ METAGLIP TABS ¹ DUETACT ² | 1. Use individual ingredients. 2. Use Actos with generic glimepiride. Use PA Form# 20420 | Approved for patients failing to achieve good diabetic control with maximal doses of individual components. |
| DIABETIC - MEGLITINIDES | MC | | NATEGLINIDE | MC/DEL MC/DEL | PRANDIN TABS STARLIX TABS | Use PA Form# 20420 | Preferred drugs from other diabetic sub-categories must be tried and failed due to lack of inadequate diabetic control or intolerable side effects before non-preferred drug will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Prandin is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for current use with both Sporanox and gemfibrozil, due to a significant drug-drug interaction. |
| GLUCOSE ELEVATING AGENTS | | | | | | | |
| GLUCOSE ELEVATING AGENTS | MC/DEL | 1 | GLUCAGEN INJ. HYPOKIT ¹ | MC | GLUCAGON DIAGNOSTIC KIT | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | MC/DEL | 2 | BAQSIMI ^{2,4} | MC MC/DEL MC | GLUCAGEN DIAGNOSTIC KIT GVOKE ³ ZEGALOGUE ⁵ | 1. Dosing limits apply, please see dose consolidation list. 2. For the treatment of patients ≥ 4 years of age. 3. For the treatment of patients ≥ 2 years of age. 4. Baqsimi will require a step through Glucagen. 5. For the treatment of patients ≥ 6 years of age. | |

| THYROID | | | | | | | |
|--|--|--|--|--|--|--|--|
| THYROID EYE DISEASE | | | | MC | | TEPEZZA Use PA Form# 20420 | |
| THYROID HORMONES | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | | ARMOUR THYROID TABS CYTOMEL TABS ERMEZA ¹ LEVOTHROID TABS LEVOTHYROXINE SODIUM TABS LEVOXYL TABS UNITHROID TABS | MC MC/DEL MC MC/DEL | | LEVOTHYROXINE SODIUM SOLR LIOETHYRONINE SYNTHROID TABS THYQUIDITY Use PA Form# 20420 1. Clinical PA is required to confirm diagnosis of dysphagia. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ANTITHYROID THERAPIES | MC/DEL MC/DEL | | METHIMAZOLE TABS PROPYLTHIOURACIL TABS | MC/DEL | | TAPAZOLE TABS Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| CUSHING DISEASE AGENTS | | | | | | | |
| CUSHING DISEASE AGENTS | | | | MC MC | | ISTURISA ¹ RECORLEV Use PA Form #20420 1. For the treatment of adult patients with Cushing's disease for whom pituitary surgery is not an option or has not been curative. | Recorlev® is associated with dose-related QT interval prolongation. QT interval prolongation may lead to life-threatening ventricular dysrhythmias such as Torsades de pointes. |
| OSTEOPOROSIS / BONE AGENTS | | | | | | | |
| OSTEOPOROSIS | MC/DEL | | ALENDRONATE | MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC MC MC MC/DEL | | ACTONEL TABS ARELIA SOLR BINOSTO BONIVA INJECTION KIT BONIVA TABS ^{2,4} CALCITONIN NS DUAVEE DIDRONEL TABS EVISTA TABS ¹ EVENTY ² FORTEO FORTICAL FOSAMAX TABS AND PLUS D ³ PROLIA SOHONOS ⁶ STRENSIQ ⁵ TYMLOS XGEVA ZOMETA Use PA Form# 20420 1. Approval only requires failure of Alendronate. 2. Quantity limits apply, please see dosage consolidation list. 3. Please use Alendronate and Vitamin D. 4. Please use other preferred agents. 5. Obtain baseline ophthalmology exams and renal ultrasounds and then periodically during treatment 6. Clinical PA for indication required. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Binosto use preferred generic alendronate tablets Evenity® should be limited to 12 monthly doses Sohonos: For the reduction in volume of new heterotopic ossification in adults and pediatric patients aged 8 years and older for females and 10 years and older for males with fibrodysplasia ossificans progressiva (FOP). |
| FIBROBLAST GROWTH FACTOR 23 INHIBITORS | MC | | CRYSVITA ¹ | | | Use PA Form #20420 1. Preferred for patients <21 years for the treatment of X-linked hypophosphatemia. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| CALCIMIMETIC AGENTS | | | | | | | |
| CALCIMIMETIC AGENTS | | | | MC MC | | PARSABIV SENSIPAR Use PA Form# 30115 | For Sensipar baseline PTH, Ca, and phosphorous levels are required and initial approvals will be limited to 3 months. Subsequent approvals will require additional levels being done to assess changes. Will not approve if baseline Ca is less than 8.4. |

Parsabiv is for the treatment of secondary hyperparathyroidism (HPT) in adults with chronic kidney disease (CKD) on hemodialysis. Parsabiv® has not been studied in adults with parathyroid carcinoma, primary hyperparathyroidism, or with chronic kidney disease who are not on hemodialysis and is not recommended for use in these populations.

GROWTH HORMONE

| | | | | | | | |
|----------------|--------|--|------------|---|----------------|--|---|
| GROWTH HORMONE | MC/DEL | GENOTROPIN ¹ NORDITROPIN SOLN ¹ NUTROPIN AQ ¹ | MC | 8 | HUMATROPE SOLR | Use PA Form# 10710 1. Clinical PA is required to establish diagnosis and medical necessity. | See Growth Hormone PA form for criteria. Step-order will still apply unless clinical contraindication supplied. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | MC/DEL | | MC | 8 | INCRELEX | | |
| | | | MC | 8 | NUTROPIN | | |
| | MC/DEL | | MC/DEL | 8 | NGENLA | | |
| | | | MC | 8 | OMNITROPE | | |
| | | | MC | 8 | SAIZEN SOLR | | |
| | | | MC | 8 | SKYTROFA | | |
| | | | MC/DEL | 8 | SOGROYA | | |
| | MC/DEL | 8 | TEV-TROPIN | | | | |

| | | | | | | | |
|--------------------------|--|--|----|--|----------------------|---|---|
| ACHONDROPLASIA TREATMENT | | | MC | | VOXZOGO ¹ | 1. Pediatric patients with achondroplasia who are 5 years of age and older with open epiphyses. Use PA Form# 20420 | Voxzogo: To increase linear growth in pediatric patients with achondroplasia who are 5 years of age and older with open epiphyses. This indication is approved under accelerated approval based on an improvement in annualized growth velocity. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s). |
|--------------------------|--|--|----|--|----------------------|---|---|

| | | | | | | | |
|---------------------|--|--|--------|---|-----------------------------|---|--|
| SOMATOSTATIC AGENTS | | | MC/DEL | 7 | OCTREOTIDE INJ ¹ | Use PA Form# 10710 1. Non-preferred products must be used in specified step order. | |
| | | | MC | 8 | BYNFEZIA ¹ | | |
| | | | MC | 8 | MYCAPSSA ¹ | | |
| | | | MC/DEL | 8 | SANDOSTATIN ¹ | | |
| | | | MC | 8 | SOMATULINE ¹ | | |

GROWTH HORMONE ANTAGONISTS

| | | | | | | | |
|----------------|--|--|----|--|----------|------------------------------------|--|
| GH ANTAGONISTS | | | MC | | SOMAVERT | Use PA Form# 10710 | Approved for acromegaly patients failing surgery/radiation/drug therapy including bromocriptine and sandostatin. |
|----------------|--|--|----|--|----------|------------------------------------|--|

VASOPRESSIN RECEPTOR ANTAGONIST

| | | | | | | | |
|---------------------------------|--|--|--------|--|-----------------------|---|---|
| VASOPRESSIN RECEPTOR ANTAGONIST | | | MC | | JYNARQUE ¹ | Use PA Form# 20420 1. Clinical PA required for appropriate diagnosis | Samsca Drug Warning- Avoid use in patients with underlying liver disease, including cirrhosis, because the ability to recover from liver injury may be impaired. Limit duration of therapy to 30 days to minimize the risk of liver injury. DDI: Jynarque- Concomitant use with strong CYP3A inhibitors is contraindicated. Avoid concomitant use of Jynarque® with OATP1B1/B3 and OAT3 substrates (e.g. statins, bosentan, glyburide, nateglinide, repaglinide, methotrexate, furosemide). |
| | | | MC/DEL | | SAMSCA | | |

URINARY INCONTINENCE

| | | | | | | | |
|--------------|--------|---------------------------------|--------|---|--|---|---|
| VASOPRESSINS | MC/DEL | DESMOPRESSIN TABS DDAVP SOLN | MC/DEL | 5 | DDAVP TABS | 1. Products must be used in specified step order. Nocturnal enuresis patients will be encouraged to periodically attempt stopping DDAVP. 2. Patients with a diagnosis of hemophilia or Von Willebrands disease will be exempt from prior authorization. Use PA Form# 20420 | Approved for central diabetes insipidus and for nocturnal enuresis. For nocturnal enuresis- must be over 6 years old, must fail an adequate trial of alarm training (higher success rate, lower relapse rate) and must periodically attempt weaning (at 6 month intervals). |
| | MC/DEL | | MC/DEL | 6 | DESMOPRESSIN SPRAY ¹ | | |
| | | | MC | 8 | DESMOPRESSIN ACETATE SOLN ¹ | | |
| | | | MC/DEL | 8 | NOCDURNA ¹ | | |
| | | | MC | 8 | NOCTIVA ¹ | | |
| | | | MC/DEL | 8 | STIMATE SOLN ^{1,2} | | |

| | | | | | | | |
|----------------|--------|---|--------|-------------------|--------------------|------------------------------------|--|
| ANTISPASMODICS | MC/DEL | DETROL TABS DETROL LA CAPS OXYBUTYNIN | MC/DEL | 8 | DARIFENACIN ER TAB | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | MC/DEL | | 8 | DITROPAN | | | |
| | MC/DEL | | 8 | FLAVOXATE HCL TAB | | | |
| | | | 8 | TOLTERODINE | | | |

| | | | | | | | |
|------------------------------|--------|---------------------|----|---|------------------|------------------------------------|--|
| ANTISPASMODICS - LONG ACTING | MC/DEL | GELNIQUE GEL PACKET | MC | 8 | DITROPAN XL TBCR | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered |
|------------------------------|--------|---------------------|----|---|------------------|------------------------------------|--|

| | | | | | | | | | |
|---|--|--|--|------------------------------------|-----------------------|--|---|---|--|
| | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | | MYRBETRIQ OXYBUTYININ ER TABS OXYTROL SOLIFENACIN SUCCINATE TAB TOVIAZ TROSPIUM | MC/DEL MC MC/DEL MC MC | 8 8 8 8 8 | ENABLEX ^{1,2} GEMTESA ² TOLTERODINE TAB VESICARE ¹ VESICARE ³ LS | 1. See Criteria Section. 2. Use a preferred long acting antispasmodic. 3. For the treatment of patients ≥ 2 years of age. | on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. 1. Vesicare 5mg and Enablex 7.5mg maximum doses if given with drugs known to be significant CYP3A4 inhibitors.(Ketoconazole, Sporanox, Erythromycin, Fluconazole, Nefazodone, Nelfinavir, and Ritonavir) DDI: Enablex 15mg and Vesicare 10mg will now be non-preferred and require prior authorization if they are currently being used in combination with any of the following medications: clarithromycin, erythromycin, Ketek, Crixivan, Norvir, ketoconazole, fluconazole (except 150mg strength), Sporanox, nefazodone, or diltiazem. | |
| CHOLINERGIC | MC/DEL | | BETHANECHOL | MC/DEL | | URECHOLINE | Use PA Form# 20420 | | |
| HYPERAMMONIA TREATMENTS | MC | | CARGLUMIC ACID TABS | MC | | CARBAGLU TABS | | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. | |
| | | | | | | | Use PA Form# 20420 | | |
| UREA CYCLE DISORDER | MC MC | | BUPHENYL TABLET PHEBURANE GRANULES | MC MC MC MC/DEL MC/DEL | | BUPHENYL POWDER RAVICTI LIQUID OLPRUVA SODIUM PHENYL BUTYRATE POWDER SODIUM PHENYL BUTYRATE TAB | | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Olpruva: As adjunctive therapy to standard of care, which includes dietary management, for the chronic management of adult and pediatric patients weighing 20kg or greater and with a body surface area (BSA) of 1.2m2 or greater, with urea cycle disorders (UCDs) involving deficiencies of carbamylphosphate synthetase (CPS), ornithine transcarbamylase (OTC), or argininosuccinic acid synthetase (AS). | |
| | | | | | | | Use PA Form# 20420 | | |
| METABOLIC MODIFIER | | | | | | | | | |
| HERED. TYROSINEMIA | | | | MC | | ORFADIN | Use PA Form# 20420 | Approved for Type 1 hereditary tyrosinemia patients. Must include laboratory evidence of dx at first PA. | |
| FABRY DISEASE AGENTS | | | | MC MC MC/DEL | | ELFABRIO ¹ FABRAZYME ² GALAFOLD ¹ | 1.Clinical PA to verify appropriate diagnosis. 2.For the treatment of patients 2 years of age and older. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Elfabrio and Galfold: For the treatment of adults with confirmed Fabry disease. | |
| | | | | | | | Use PA Form# 20420 | | |
| ANTIHYPERTENSIVES / CARDIAC | | | | | | | | | |
| CARDIAC GLYCOSIDES | MC/DEL MC/DEL MC/DEL | | DIGITEK TABS DIGOXIN LANOXIN | | | | Use PA Form# 20420 | | |
| CARDIAC MYOSIN INHIBITORS | | | | MC | | CAMZYOS | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Camzyos: For the treatment of adults with symptomatic New York Heart Association (NYHA) class II-III obstructive hypertrophic cardiomyopathy (HCM) to improve functional capacity and symptoms. DDI: Concomitant use of Camzyos® with a moderate to strong CYP2C19 inhibitor or a strong CYP3A4 inhibitor is contraindicated. | |
| CARDIAC - SINUS NODE INHIBITORS | | | | MC | | CORLANOR | Use PA Form#20420 | In patients with stable, symptomatic chronic heart failure with left ventricular ejection fraction ≤35%, who are in sinus rhythm with resting heart rate ≥70 beats per minute (bpm) and | |
| CARDIAC - SOLUBLE GUANYLATE CYCLASE STIMULATORS | | | | MC/DEL | | VERQUVO | | | |

| | | | | | | | |
|--|--|--------|---|--|--|---|--|
| | | | | | | Use PA Form# 20420 | |
| CARDIAC- SODIUM- GLUCOSE COTRANSPORTER 2 (SGLT2) INHIBITOR | | | | MC | | INPEFA ¹ | 1. To reduce the risk of cardiovascular death, hospitalization for heart failure, and urgent heart failure visit in adults with: Heart failure or Type 2 diabetes mellitus, chronic kidney disease, and other cardiovascular risk factors. Other Preferred SGLT inhibitors must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ANTIANGINALS--Isosorbide Di-nitrate/ Mono-Nitrates | MC/DEL MC/DEL | | ISOSORBIDE MONONITRATE TABS ISOSORBIDE MONONITRATE ER | MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC | | DILATRATE SR CPR ISORDIL TABS ISORDIL TITRADOSE TABS ISOSORBIDE DINITRATE SUBL ISOSORBIDE DINITRATE TABS ISOSORBIDE DINITRATE CR TBCR ISOSORBIDE DINITRATE ER TBCR ISOSORBIDE DINITRATE TD TBCR IMDUR TB24 ISMO TABS MONOKET TABS | Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| NITRO - OINTMENT/CAP/CR | MC/DEL MC/DEL MC MC | | NITROBID OINT NITROGLYCERIN CPR NITROL OINT NITRO-TIME CPR | | | | Use PA Form# 20420 |
| NITRO - PATCHES | MC/DEL MC/DEL | 1 1 | NITROGLYCERIN PT24 ¹ NITRO-DUR PT 24 0.8MG ¹ | MC MC/DEL | | NITRODISC PT24 NITRO-DUR PT24 | 1. At least 2 step 1's and step 3 of the preferred products must be used in specified order or PA will be required. Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| NITRO - SUBLINGUAL/ SPRAY | MC/DEL | | NITROSTAT SUBL | MC/DEL MC MC | | NITROQUICK SUBL NITROLINGUAL SOLN NITROLINGUAL TABS | Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| BETA BLOCKERS - NON SELECTIVE | MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL | | CARVEDILOL LEVATOL TABS NADOLOL TABS PINDOLOL TABS PROPRANOLOL HCL SOLN ¹ PROPRANOLOL HCL TABS ¹ PROPRANOLOL HCL 60MG TABS PROPRANOLOL LA CAPS RANOLAZINE ER TABS SOTALOL AF SOTALOL HCL TABS TIMOLOL MALEATE TABS | MC MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC MC MC MC | | ASPRUZYO BETAPACE TABS BETAPACE AF TABS COREG CR ³ COREG TABS CORGARD TABS INDERAL TABS HEMANGEOL SOL INDERAL XL CAP INDERAL LA CPR INNOPRAN XL RANEXA | 1. Recommend using BID since its effects do not last 24 hours. 2. Please use other strengths in combination to obtain this dose. 3. Dosing limits still apply. Please see dose consolidation list Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Concomitant use of Ranolazine products with strong CYP3A inhibitors, including ketoconazole, itraconazole, clarithromycin, nefazodone, nelfinavir, ritonavir, indinavir, and saquinavir, is contraindicated. |
| BETA BLOCKERS - CARDIO SELECTIVE | MC/DEL MC/DEL MC/DEL | | ACEBUTOLOL HCL CAPS ATENOLOL TABS ¹ BETAXOLOL HCL TABS | MC MC/DEL MC | | KERLONE TABS LOPRESSOR TABS SECTRAL CAPS | 1. Recommend using Atenolol (and metoprolol) BID since its effects do not last 24 hours Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |

| | | | | | | | | | | |
|---|--|--------------------------------------|---|--|--|---|---|------------------------------------|--|---|
| | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | | BISOPROLOL FUMARATE TABS BYSTOLIC METOPROLOL TARTRATE TABS ¹ METOPROLOL ER NEBIVOLOL HCL TAB | MC/DEL MC/DEL MC/DEL | | TENORMIN TABS TOPROL XL TB24 ZEBETA TABS | | Use PA Form# 20420 | | |
| BETA BLOCKERS - ALPHA / BETA | MC/DEL | | LABETALOL HCL TABS | MC | | TRANDATE TABS | | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. | |
| BETA BLOCKERS & DURECTIC COMBOS | MC/DEL | | METOPROLOL-HYDROCHLOROTHIAZIDE TAB | MC/DEL | | DUTOPROL | | Use PA Form# 20420 | | |
| CALCIUM CHANNEL BLOCKERS-- Amlodipines, Bepridil, Diltiazems, Felodipines, Isradipines, Nifedipines, Nisoldipine, and Verapamils | MC/DEL | | AMLODIPINE ¹ | MC/DEL MC MC/DEL | | KATERZIA NORLIQVA NORVASC TABS ¹ | | Use PA Form# 20420 | 1. Dosing limits apply, please see dose consolidation list. | |
| | MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL | | DILTIA XT CP24 DILTIAZEM HCL ER CP24 DILTIAZEM HCL XR CP24 DILTIAZEM CD 300MG CP24 DILTIAZEM CD 360MG CP24 CARTIA XT CP24 ¹ DILTIAZEM CD CP24 ¹ DILTIAZEM HCL ER CP24 ¹ DILTIAZEM XR CP24 ¹ TIAZAC CP24 ¹ | MC/DEL MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL | 5 6 8 8 8 8 8 8 8 | | DILACOR XR CP24 ¹ TAZTIA ¹ CARDIZEM TABS ¹ CARDIZEM CD CP24 ¹ CARDIZEM LA TB24 ¹ CARDIZEM SR CP12 ¹ DILTIAZEM HCL TABS ¹ DILTIAZEM HCL ER CP12 ¹ DILTIAZEM HCL ER CP12 ¹ | | Use PA Form# 20420 | 1. Products must be used in specified order or PA will be required. Just write "Diltiazem 24-hour" and the pharmacy will use a preferred long acting diltiazem that does not require PA. Preferred drugs must be tried and failed (in step-order) due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: All preferred diltiazems will now be non-preferred and require prior authorization if they are currently being used in combination with either Enablex 15mg or Vesicare 10mg. All non-preferred diltiazems require prior authorization, but with any prior authorization request, the member's drug profile will also be monitored for current use with Enablex 15mg or Vesicare 10mg. |
| | | | | MC/DEL MC/DEL | | | PLENDIL TB24 FELODIPINE | | Use PA Form# 20420 | Other Preferred calcium channel blockers must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | | | | MC MC | | | DYNACIRC CAPS DYNACIRC CR TBCR ¹ | | Use PA Form# 20420 | 1. Established users will be grandfathered Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | | | | MC MC | | | CARDENE SR CPR NICARDIPINE HCL CAPS | | Use PA Form# 20420 | Other Preferred calcium channel blockers must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | | | AFEDITAB CR NIFEDIAC CC NIFEDICAL XL TBCR NIFEDIPINE TBCR NIFEDIPINE ER TBCR | MC/DEL MC/DEL MC/DEL MC/DEL | | ADALAT CC TBCR ¹ NIFEDIPINE CAPS PROCARDIA CAPS PROCARDIA XL TBCR | | Use PA Form# 20420 | 1. Established users of Adalat CC are grandfathered. Preferred drug must be tried and failed in step order due to lack of efficacy or intolerable side effects before non-preferred drugs in step order will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | | | | | MC MC | | SULAR TB24 SULAR CR ¹ | | Use PA Form# 20420 | 1. Established users of 10MG and 20MG strengths are grandfathered. |
| | MC/DEL MC/DEL MC/DEL | 1 1 1 | | VERAPAMIL HCL CR TBCR VERAPAMIL HCL ER TBCR VERAPAMIL HCL SR TBCR | MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL | | CALAN TABS CALAN SR TBCR COVERA-HS TBCR ISOPTIN-SR VERAPAMIL HCL ER CP24 VERAPAMIL HCL SR CP24 VERAPAMIL HCL TABS VERELAN CP24 VERELAN PM CP24 | | Use PA Form# 20420 | Products must be used in specified order or PA will be required. Just write "Verapamil 24-hour" and the pharmacy will use a preferred long acting generic that does not require PA. Preferred drugs must be tried and failed (in step-order) due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | ANTIARRHYTHMICS | MC/DEL MC/DEL MC/DEL MC/DEL | | AMIODARONE HCL DISOPYRAMIDE FLECAINIDE MEXILETINE HCL | MC/DEL MC/DEL MC/DEL MC/DEL | | CORDARONE DISOPYRAMIDE MULTAQ NORPACE | | Use PA Form# 20420 | 1. Prescription must be written by Cardiologist. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |

| | | | | | | | | |
|--|--|--|---|--|--|---|---|---|
| | MC/DEL MC/DEL MC MC/DEL MC/DEL | | PROCAINAMIDE PROPAFENONE QUINAGLUTE QUINIDINE GLUCONATE QUINIDINE SULFATE | MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL | | PACERONE QUINIDEX TAMBOCOR TIKOSYN ¹ RYTHMOL SR RYTHMOL | Use PA Form# 20420 | DDI: Amiodarone will now be non-preferred and require prior authorization if it is currently being used in combination with either Lovastatin (doses greater than 40mg/day) or Lipitor (doses greater than 20mg/day) or Levofloxacin or Gemifloxacin, or Moxifloxacin, or Ofloxacin. DDI: Multaq will be preferred unless the following medications are seen in the member's drug profile within the last 35 days for brand name medications or 90 days for generic medications: Erythromycin, Amiodarone and other antiarrhythmics, TCA's, Phenothiazine, Ketoconazole, Itraconazole, Voriconazole, Cyclosporine, Telithromycin, Clarithromycin, Nefazodone, Ritonavir. |
| ACE INHIBITORS | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | | BENAZEPRIL HCL CAPTOPRIL TABS ENALAPRIL MALEATE TABS FOSINOPRIL SODIUM LISINOPRIL TABS RAMIPRIL QUINAPRIL HCL | MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL | 5 5 8 8 8 8 8 8 8 8 8 8 8 8 | MAVIK TABS ACCUPRIL TABS ACEON TABS ¹ ALTACE CAPS ¹ EPANED LOTENSIN TABS ¹ MOEXIPRIL HCL ¹ MONOPRIL HCT TABS ¹ PRINIVIL TABS ¹ QBRELIS UNIVASC ¹ VASOTEC TABS ¹ ZESTRIL TABS ¹ | 1. Non-preferred products must be used in specified order. Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Non-preferred products are subject to step-order requirements unless clinical circumstances warrant exception. |
| ANGIOTENSIN RECEPTOR BLOCKER | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | | AMLODIPINE-OLMESARTAN TAB ³ IRBESARTAN ¹ LOSARTAN ¹ MICARDIS TABS ³ OLMESARTAN ¹ TELMISARTAN ¹ | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC | 8 8 8 8 8 8 | ATACAND TABS AVAPRO BENICAR TABS COZAAR DIOVAN EDARBI TEVETEN TABS | Use PA Form# 20420 1. Dosing limits apply, please see dose consolidation list. 2. Use preferred active ingredients which are available without PA. 3. Preferred without a PA only if patient on a diabetic therapy or prior ACE therapy. | Per best practices patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy |
| DIRECT RENIN INHIBITOR | | | | MC/DEL MC/DEL MC/DEL | | AMTURNIDE TEKTURNA ¹ TEKAMLO | 1. Must show failure of single and combination therapy from all preferred antihypertensive categories. Use PA Form# 20420 | |
| ANTIHYPERTENSIVES - CENTRAL | MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL | | CLONIDINE HCL TABS GUANFACINE HCL TABS HYDRALAZINE HCL TABS HYLOREL TABS METHYLDOPA TABS MINOXIDIL TABS PRAZOSIN HCL CAPS RESERPINE TABS | MC/DEL MC/DEL MC MC MC/DEL MC MC/DEL | | CLONIDINE PATCH CLONIDINE TTS GUANABENZ ACETATE TABS ISMELIN TABS MINIPRESS CAPS NEXICLON TENEX TABS | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ACE INHIBITORS AND CA CHANNEL BLOCKERS | | | | MC/DEL MC MC MC/DEL | 8 8 8 9 | AMLODIPINE/BENAZEPRIL PRESTALIA ¹ TARKA TBCR LOTREL CAPS | 1. Prestalialia will only be approved for patients ≥ 18 years of age. Use individual preferred generic medications. Use PA Form# 20420 | |
| ACE AND THIAZIDE COMBO'S | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | | BENAZEPRIL HCL/HYDROCHLOR CAPTOPRIL/HYDROCHLOROTHIA ENALAPRIL MALEATE/HCTZ TABS LISINOPRIL-HCTZ TABS LOTENSIN HCT TABS | MC/DEL MC MC/DEL MC/DEL MC MC/DEL | | ACCURETIC TABS MONOPRIL HCT TABS PRINZIDE TABS UNIRETIC TABS VASERETIC TABS ZESTORETIC TABS | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |

| | | | | | | | | |
|--|--|--|---|--|----------------------------|--|---|--|
| BETA BLOCKERS AND DIURETIC COMBO'S | MC/DEL MC/DEL MC/DEL | | ATENOLOL/CHLORTHALIDONE BISOPROLOL FUMARATE/HCTZ PROPRANOLOL/HCTZ | MC/DEL MC/DEL MC MC MC/DEL | | CORZIDE TABS LOPRESSOR HCT TABS TENORETIC TIMOLIDE 10/25 TABS ZIAC TABS | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ARB'S AND CA CHANNEL BLOCKERS | MC/DEL MC/DEL MC/DEL | | AMLODIPINE/VALSARTAN AMLODIPINE/VALSARTAN HCT TRIBENZOR | MC/DEL MC MC/DEL MC/DEL | | AZOR BYVALSON EXFORGE EXFORGE HCT | Use PA Form# 20420 | DDI: Byvalson will be non-preferred and require a prior authorization if it is currently being used in combination with drugs known to be significant CYP2D6 inhibitors (e.g. quinidine, propafenone, fluoxetine, paroxetine). Per best practices patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy |
| ARB'S AND DIURETICS | MC/DEL MC/DEL MC/DEL MC/DEL | | BENICAR HCT ¹ LOSARTAN HCT ¹ MICARDIS HCT TABS ¹ VALSARTAN-HCT ¹ | MC/DEL MC/DEL MC MC/DEL MC/DEL MC | 7 8 8 8 8 8 | IRBESARTAN HYDROCHLOROTHIAZIDE ATACAND HCT TABS AVALIDE TABS ¹ DIOVAN HCT TABS ¹ HYZAAR TABS TEVETEN HCT TABS | 1. Dosing limits apply, please see dose consolidation list. Use PA Form# 20420 | Per best practices patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy |
| ANGIOTENSIN MODULATORS-ARB COMBINATION | MC | | ENTRESTO | MC/DEL MC | | EDARBYCLOR ENTRESTO SPRINKLES | Use PA Form# 20420 | |
| ARB'S AND DIRECT RENIN INHIBITOR COMBINATION | | | | MC/DEL | | VALTURNA | Use PA Form# 20420 | |
| DIURETICS | MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC | | ACETAZOLAMIDE TABS BUMETANIDE CHLOROTHIAZIDE TABS CHLORTHALIDONE TABS EDECIN TABS EDECIN TABS HYDROCHLOROTHIAZIDE INDAPAMIDE TABS METHAZOLAMIDE TABS METHYCLOTHIAZIDE TABS SPIRONOLACTONE SPIRONOLACTONE/HYDRO TORSEMIDE TABS TRIAMTERENE/HCTZ ZAROXOLYN TABS | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC | | ALDACTAZIDE TABS ALDACTONE TABS AMILORIDE HCL BUMEX TABS DEMADEX TABS DIAMOX DIURIL DYAZIDE CAPS CAROSPIR ENDURON TABS FUROSCIX INSPIRA KERENDIA KEVEYIS LASIX TABS MAXZIDE MICROZIDE CAPS MIDAMOR TABS NAQUA TABS | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Furoscix: The indication for use is the treatment of congestion due to fluid overload in adults with NYHA Class II or Class III chronic heart failure AND the medication is being prescribed by or in consultation with a cardiologist AND the patient is experiencing symptoms despite compliance with oral loop diuretic therapy AND oral loop diuretic therapy will be resumed as soon as practical AND medical reasoning beyond convenience is provided for not pursuing therapy in an outpatient infusion setting. PA approval will be authorized for 1 month. DDI: The concomitant use of Keveyis® with high dose aspirin is contraindicated. |
| CCB / LIPID | | | | MC/DEL | | CADUET | Use PA Form# 20420 | |
| NEUROGENIC ORTHOSTATIC HYPOTENSION | | | | | | | | |
| NEUROGENIC ORTHOSTATIC HYPOTENSION | | | | MC | | NORTHERA | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| LIPID DRUGS | | | | | | | | |
| CHOLESTEROL - BILE SEQUESTRANTS | MC/DEL MC/DEL | | CHOLESTYRAMINE COLESTIPOL HCl | MC/DEL MC/DEL MC MC/DEL | | COLESTID PREVALITE QUESTRAN WELCHOL TABS | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| CHOLESTEROL - FIBRIC ACID DERIVATIVES | MC/DEL MC/DEL MC/DEL | | FENOFIBRATE TAB GEMFIBROZIL TABS NIACIN ER | MC MC/DEL MC/DEL MC/DEL | | ANTARA LOPID FENOFIBRATE 120mg TAB FENOFIBRATE CAP FIBRICOR | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Fenofibrate is preferred but will require a prior authorization requests if used concurrent with Warfarin. |

Additional criteria for the diagnosis of homozygous familial hypercholesterolemia (Repatha only): Total cholesterol levels > 290mg/dL or LDL-C > 190mg/dL (adults) OR Total cholesterol levels > 260mg/dL or LDL-C > 155mg/dL (children < 16 years) and TG within reference range OR Confirmation of diagnosis by gene testing.

[Use PA Form# 20420](#)

PULMONARY ANTI-HYPERTENSIVES

| | | | | | | | | |
|-------------------------------------|--|--|---|--|--|--|---|---|
| PULMONARY ANTI-HYPERTENSIVES | MC MC/DEL MC/DEL MC | | EPOPROSTENOL INJ ^{3,6} SILDENAFIL TADALAFIL VENTAVIS ³ | MC/DEL MC MC/DEL MC MC MC MC MC MC/DEL MC MC MC MC MC/DEL | | ADEMPAS ^{1,3} ADCIRCA ⁴ ALYQ TAB FLOLAN ³ LIQREV OPSUMIT ^{1,2} OPSYNVI ⁴ ORENITRAM REMODULIN ³ REVATIO ⁴ TADLIQ ⁴ TYVASO UPTRAVI VELVETRI ³ WINREVAIR⁴ | 1. Requires previous trials/failure of multiple preferred medications. 2. Dosing limits apply, please see the dose consolidation list. 3. Require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and NYHA functional class 3 or 4. 4. Require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and NYHA (WHO) functional class 2 or 3. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Sildenafil will be preferred with clinical PA for treatment of pulmonary arterial hypotension (WHO Group 1) in adults to improve exercise ability and delay clinical worsening. Avoid concomitant use of Sildenafil with moderate or strong Cyp3A inhibitors DDI: Upravi will require a prior authorization if it is currently being used in combination with strong inhibitors of CYP2C8 (gemfibrozil) DDI: Opsumit will require a prior authorization if it is currently being used in combination with drugs known to be significant CYP3A inhibitors (ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, nelfinavir, ritonavir, atazanavir, saquinavir and telithromycin). DDI: Adempas will require a prior authorization if it is currently being used in combination with drugs known to be PDE inhibitors should be avoided (including dipyridamole, addira and tadalafil) with adempas Liqrev: treatment of pulmonary arterial hypertension (WHO Group 1) in adults to improve exercise ability and delay clinical worsening. Avoid concomitant use of Liqrev with moderate or strong CYP3A inhibitors. |
|-------------------------------------|--|--|---|--|--|--|---|---|

[Use PA Form# 20420](#)

| | | | | | | | | |
|---|------------------------|--|-------------------------------------|--|--|--|---|---|
| ERA / ENDOTHELIN RECEPTOR ANTAGONIST | MC MC | | LETAIRIS ^{1,2} TRACLEER | | | | 1. Providers must be registered with LEAP Prescribing program, a restricted distribution program. 2. Clinical PA is required to establish diagnosis and medical necessity. | Tracleer approvals will require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and NYHA functional class 2 thru 4. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with Tracleer. Letairis approvals will require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and functional class 2 or 3 symptoms. |
|---|------------------------|--|-------------------------------------|--|--|--|---|---|

[Use PA Form# 20420](#)

IMPOTENCE AGENTS

| | | | | | | | | |
|-------------------------|--|--|--|--|--|--|---|---|
| IMPOTENCE AGENTS | | | | | | | As of January 1, 2006, per CMS (federal govt.), impotence agents are no longer covered. | As of January 1, 2006, per CMS (federal govt.), impotence agents are no longer covered. |
|-------------------------|--|--|--|--|--|--|---|---|

ANTI-EMETOGENICS

| | | | | | | | | |
|--|--|--|--|--|----------------------------|--|--|--|
| ANTIEMETIC - ANTICHOLINERGIC / DOPAMINERGIC | MC MC/DEL MC MC/DEL MC | | BONJESTA MECLIZINE HCL TABS PROMETHAZINE SUPP PROMETHAZINE TRANSDERM-SCOP PT72 | MC MC MC MC MC | | ANTIVERT TABS BARHEMSYS PHENERGAN SOLN PROMETHAZINE 50MG SUPP PROMETHEGAN SUPP TORECAN TABS | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Concomitant use of MAOIs and Bonjesta® is contraindicated. |
| ANTIEMETIC - 5-HT3 RECEPTOR ANTAGONISTS/ SUBSTANCE P NEUROKININ | MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | | DICLEGIS DRONABINOL CAPS GRANISETRON TAB ONDANSETRON TAB ONDANSETRON ODT TBDP ONDANSETRON SOL | MC MC MC MC MC MC | 8 8 8 8 8 8 | AKYNZEO ¹ APREPITANT ALOXI ANZEMET TABS APONVIE ⁴ CESAMET ¹ CINVANTI ⁴ | 1. Approvals will require diagnosis of chemo-induced nausea/vomiting and failed trials of all preferred anti-emetics, including 5-HT3 class (Ondansetron) and Marinol. | Preferred drugs and step therapy must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. * Ondansetron limits still apply as listed on the Ondansetron PA form for covered indications including chemotherapy, radiotherapy, post operative nausea & vomiting and hyperemesis gravidarum. Other medical indications will be approved or denied on a case by case basis. Hyperemesis and other medical indications approved are still subject to failure of multiple preferred antiemesis drugs. |

| | | | | | | | | |
|---|------------------------------|--|--|--|--|---|---|--|
| | | | | MC | 8 | EMEND ² | | Akynzeo- Concomitant use should be avoided in patients who are chronically using a strong CYP3A inducer such as rifampin. |
| | | | | MC | 8 | FOCINVEZ ^{1,2} | | |
| | | | | MC/DEL | 8 | KYTRIL | 2. Clinical PA is required for members on highly emetic anti-neoplastic agents. | Varubi – Available to the few who are unable to tolerate or who have failed on preferred medications |
| | | | | MC/DEL | 8 | MARINOL CAPS | | |
| | | | | MC | 8 | SANCUSO | | Aponvie is for the prevention of postoperative nausea and vomiting (PONV) in adults. |
| | | | | MC | 8 | SUSTOL | | |
| | | | | MC | 8 | SYNDROS | 3. Dosing limits apply, please see Dosage Consolidation List | |
| | | | | MC | 8 | TRIMETHOBENZAMIDE CAP | | |
| | | | | MC | 8 | VARUBI | | |
| | | | | MC/DEL | 8 | ZOFRAN ODT TBDP ³ | 4. Clinical PA required for appropriate diagnosis | |
| | | | | MC/DEL | 8 | ZOFRAN TABS ³ | | |
| | | | | MC/DEL | 8 | ZOFRAN INJ ³ | | |
| | | | | MC | 8 | ZUPLENZ | | |
| | | | | | | | Use PA Form# 20420 | |
| NON-SEDATING ANTIHISTAMINES / DECONGESTANTS | | | | | | | | |
| ANTIHISTIMINES - NON-SEDATING | MC MC/DEL MC/DEL MC | | ALAVERT TABS CETIRIZINE TABS LORATADINE TAVIST ND (OTC) | MC MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL | 5 5 5 5 8 8 8 8 8 9 | CLARINEX TABS ^{1,5} CLARINEX SYR ^{1,2} FEXOFENADINE ¹ ZYRTEC ¹ ZYRTEC SYR ^{1,2} ALLEGRA ³ CLARITIN ³ DESLORATADIN LORATADINE ODT ⁴ LEVOCETIRIZINE ⁴ XYZAL ³ | 1. Must fail preferred drugs, OTC loratidine and cetirizine before moving to non-preferred step order drugs. 2. Clarinex and Zyrtec syrup <6 yr w/o PA. 3. Must fail all step 5 drugs (Clarinex, Fexofenadine and Zyrtec) before moving to next step product. 4. All OTC versions of loratidine ODT are now non-preferred. 5. Pa's for Clarinex RediTabs will only be approved if between the ages of 6-11 years old. | Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. No combination product with decongestant will be approved since pseudoephedrine available without PA. Pseudoephedrine is available with prescription. |
| | | | | | | | Use PA Form# 20530 | |
| ANTIHISTIMINES - OTHER | MC/DEL MC/DEL MC/DEL | | CLEMASTINE CHLORPHENIRAMINE DIPHENHYDRAMINE | | | | Use PA Form# 20530 | |
| ALLERGY / ASTHMA THERAPIES | | | | | | | | |
| ANAPHYLACTIC DEVICES | MC/DEL MC/DEL MC/DEL | | EPINEPHRINE EPIPEN EPIPEN JR | MC MC/DEL | | TWINJECT SYMJEPI | | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | | | | | | | Use PA Form# 20420 | |
| ALLERGEN IMMUNOTHERAPY | | | | MC MC MC MC MC | | ODACTRA ORALAIR ¹ PALFORZIA RAGWITEK GRASTEK | 1. See criteria section | Prescriber must provide the testing to show that the patient is allergic to the components in the prescribed therapy and must provide a clinically valid rationale why single agent sublingual therapy is being chosen over subcutaneous therapy Palforzia® is approved for use in patients with a confirmed diagnosis of peanut allergy. Initial dose escalation may be administered to patients aged 4 through 17 years. Up-dosing and maintenance may be continued in patients 4 years of age and older. Odactra® is approved for use in persons 12 through 65 years of age. Note that Odactra® is not indicated for the immediate relief of allergic symptoms. |
| | | | | | | | Use PA Form# 20420 | Treatment must start 12 weeks before expected onset of pollen season and only after confirmed by positive skin test or in vitro testing for pollen-specific IgE antibodies for any of the 5 |

| | | | | | | | | |
|--|--|--|---|--|----------------------------|---|---|--|
| | | | | | | | grass species contained in Oralair Oralair: Patient age ≥10 years and ≤65 years Have an auto-injectable epinephrine on-hand | |
| ANTIASTHMATIC - ANTICHOLINERGICS - INHALER | MC MC/DEL MC/DEL | | INCRUSE ELLIPTA ³ SPIRIVA HANDIHALER ^{1,2} SPIRIVA RESPIMAT | MC/DEL MC MC/DEL | | FLUTICASON-SALMETEROL LONHALA MAGNAIR TUDORZA | Use PA Form# 20420 1. Quantity limit of 1 inhalation daily (1 capsule) 2. We ask physicians to write "asthma" on the prescription whenever Spiriva is primarily being used for that condition. 3. Quantity limit of 1 inhalation daily | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ANTIASTHMATIC - PHOSPHODIESTERASE 4 INHIBITORS | MC/DEL | | ROFLUMILAST | MC/DEL MC | | DALIRESP OHTUVAYRE ¹ | Use PA Form# 20420 1. For the maintenance treatment of chronic obstructive pulmonary disease (COPD) in adult patients | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ANTIASTHMATIC - ANTICHOLINERGICS - NEBULIZER | MC/DEL | | IPRATROPIUM BROMIDE SOLN | MC MC/DEL | | ATROVENT SOLN YUPELRI | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ANTIASTHMATIC - ANTIINFLAMMATORY AGENTS | MC/DEL MC/DEL MC/DEL MC MC/DEL | | CROMOLYN SODIUM NEBU DUPIXENT ^{2,4} FASENRA ² FASENRA ² AUTO INJCT NUCALA ² SYRINGE 40MG XOLAIR ¹ | MC MC | | CINQAIR ³ TEZSPIRE ⁵ | 1. Need max inhaled steroids and written by pulmonary or allergy specialist. Must have elevated IgE and ≥ to age 6. 2. For patients with severe asthma aged 12 years or older and eosinophilia. 3. For patients ≥ 18 years of age with eosinophilia. 4. Clinical PA required. 5. For adult and pediatric patients aged 12 years and older with severe asthma. Use PA Form# 20420 | All will require suboptimal response to maximal doses of inhaled steroid as evidenced by asthmatic ER/Hospital admissions and Allergy/Pulmonary specialist management. Dupixent limited to patient with asthma not controlled on high dose ICS-LABA who have eosinophil greater than or equal to 150 cells or the patient is depend on an oral corticosteroid Fasenra, Nucala and Cinqair are not indicated for treatment of other eosinophilic conditions and are not indicated for the relief of acute bronchospasm or status asthmaticus. |
| ANTIASTHMATIC - NASAL STEROIDS | MC/DEL MC/DEL MC MC/DEL MC/DEL MC | | BUDESONIDE SPRAY FLUTICASON SPR ³ OLOPATADINE SPRAY OMNARIS SPR ³ TRIAMCINOLONE NS QNASL | MC MC/DEL MC/DEL MC/DEL MC/DEL MC | 5 8 8 8 8 8 | BECONASE AQ INHA ^{1,3} DYMISTA FLONASE SUSP ^{2,3} FLUNISOLIDE SOLN ^{1,3} NASONEX SUSP RHINOCORT AERO ^{2,3} | Use PA Form# 20420 1. All preferred drugs must be tried before moving to non preferred steps. 2. All step 5 medications | Preferred drugs and step therapy must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |

| | | | | | | | | |
|--|--|--|--|--|---------------------------------|---|--|---|
| | | | | MC/DEL MC MC MC MC/DEL MC MC/DEL | 8 8 8 8 8 8 8 | RHINOCORT AQUA SUSP ^{2,3} RYALTRIS ⁴ TRI-NASAL SOLN ^{2,3} VANCENASE POCKETHALER AERS ^{2,3} VERAMYST ^{2,3} XHANCE ² ZETONNA ³ | need to be tried before moving to step 8's. 3. Dosing limits apply to whole category, please see dosage consolidation list. 4. Use of individual ingredients or other preferred agents. | Xhance will be considered for the treatment of nasal polyps in patients 18 years of age or older. The patient has had a documented side effect, allergy, or treatment failure of two preferred nasal glucocorticoids, one of which must be fluticasone. |
| ANTIASTHMATIC - NASAL MISC. | MC/DEL MC/DEL MC | | AZELASTINE CROMOLYN NASAL 4% IPRATROPIUM NASAL SOL ¹ | MC/DEL MC/DEL | 8 8 | ASTEPRO ² PATANASE | Use PA Form# 20420 1. Ipratropium will be approved if submitted with documentation supporting use of CPAP machine. 2. Utilize Multiple preferred, as well as step therapy Azelastine. | Approved if patient fails on non-sedating antihistamines and steroid nasal sprays. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ANTIASTHMATIC - BETA - ADRENERGICS | MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC | | ALBUTEROL NEB METAPROTERENOL PROAIR RESPICLICK PROVENTIL HFA SEREVENT TERBUTALINE SULFATE TABS ALBUTEROL 0.63mg/3ml VENTOLIN HFA AERS | MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC MC | | ACCUNEB NEBU ALBUTEROL HFA BRETHINE LEVALBUTEROL TARTRATE PROAIR DIGIHALER ⁴ STRIVERDI VOLMAX TBCR VOSPIRE ER TB12 XOPENEX HFA ³ XOPENEX NEBU ^{1,2} | 1. Xopenex users w/ prior asthma hospitalization due to albuterol nebulizer failure will be grandfathered. 2. Quantity Limit: 12 cc/day. 3. Dosing limits apply, please see dosage consolidation list. 4. For the treatment of patients ≥ 4 years of age. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ANTIASTHMATIC - ADRENERGIC COMBINATIONS | MC MC MC MC MC/DEL MC/DEL | | ADVAIR DISKUS ¹ ADVAIR HFA ¹ AIRDUO RESPICLICK ² BREQ ELLIPTA ¹ DULERA SYMBICORT | MC MC/DEL MC/DEL MC | | AIRDUO DIGIHALER ² AIRSUPRA BREZTRI AEROSPHERE TRELEGY ELLIPTA ¹ | 1. Dosing limits apply, please see dosage consolidation list. 2. For patients ≥ 12 years and older. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. AirDuo® Respiclick be non-preferred and require prior authorization and be available to those who are unable to tolerate or who have failed on preferred medications DDI: Avoid concomitant use of strong CYP3A4 inhibitors (e.g. ritonavir, atazanavir, clarithromycin, indinavir, itraconazole, nefazodone, nelfinavir, saquinavir, ketoconazole, telithromycin) with AirDuo® Respiclick is not recommended due to increased systemic corticosteroid and increased cardiovascular adverse effects |
| ANTIASTHMATIC - ADRENERGIC ANTICHOLINERGIC | MC/DEL MC MC/DEL MC/DEL | | ALBUTEROL/IPRATROPIUM NEB. SOLN ANORO ELLIPTA COMBIVENT RESPIMAT STIOLTO | MC/DEL MC/DEL MC/DEL | | BEVESPI AEROSPHERE ^{2,3} DUAKLIR PRESSAIR DUONEB SOLN ¹ | 1. Please use preferred individual ingredients Albuterol and Ipratropium. 2. Dosing limits apply, please see dosing consolidation list. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Duoneb components are available separately without PA. DDI: Avoid concomitant use of Bevespi with other anticholinergic-containing drugs, due to an increased risk of anticholinergic adverse events. Bevespi® should be used with extreme caution in patients being treated with MAO inhibitors, TCAs, or other drugs known to prolong the QTc interval. |

| | | | | | | | |
|-----------------------------------|--|--|---|--|----------------------------|--|--|
| | | | | | | 3. The safety and efficacy of use in children under the age of 18 years have not been established. Use PA Form# 20420 | Bevespi should be used with extreme caution in patients being treated with MAO inhibitors, TCAs, or other drugs known to prolong the QTc interval. |
| ANTIASTHMATIC - XANTHINES | MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL | | AMINOPHYLLINE TABS THEOCHRON TB12 THEOLAIR-SR TB12 THEOPHYLLINE CR TB12 THEOPHYLLINE ELIX THEOPHYLLINE SOLN THEOPHYLLINE ER CP12 THEOPHYLLINE ER TB12 | MC/DEL MC MC/DEL | | THEO-24 CP24 THEOLAIR TABS UNIPHYL TBCR | Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ANTIASTHMATIC - STEROID INHALANTS | MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC | | ARNUITY ELLIPTA ASMANEX TWISTHALER ^{3,4} ASMANEX HFA ⁵ BUDESONIDE NEB 0.25MG & 0.5MG ¹ FLOVENT DISKUS ³ PULMICORT FLEXHALER ³ QVAR AERS ³ | MC MC/DEL MC MC/DEL MC/DEL MC | 8 8 8 8 8 8 | AEROSPAN ALVESCO ³ ARMONAIR DIGIHALER BUDESONIDE NEB 1MG PULMICORT SUSP FLOVENT HFA ³ | 1. Budesonide Neb 0.25mg & 0.5mg will be preferred for members under the age of 8 years old. PA will be required for members 8 years of age and older, please consider other preferred options. 2. All preferreds must be Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |

| | | | | | | | | |
|--|----------------------------|--|---|--------------------------------|------------------|---|--|--|
| | | | | | | | <p>tried before moving to non preferred steps.</p> <p>3. Dosing limits apply, please see dosage consolidation list.</p> <p>4. Asmanex 110mcg will be limited to member between the ages of 4-11years old.</p> <p>5. Asmanex HFA will be preferred for members under the age of 6 years old. PA will be required for members 6 years of age and older, please consider other preferred options.</p> <p>Use PA Form# 20420</p> | |
| ANTIASTHMATIC - 5-Lipoxygenase Inhibitors | | | | MC | | ZYFLO CR TABS | | Other Preferred asthma controller drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ANTIASTHMATIC - LEUKOTRIENE RECEPTOR ANTAGONISTS | MC/DEL MC/DEL MC/DEL | | MONTELUKAST GRANULE ¹ MONTELUKAST SODIUM TAB MONTELUKAST SODIUM CHEW TAB | MC/DEL MC/DEL MC/DEL | 8 8 8 | ACCOLATE TABS SINGULAIR ² SINGULAIR GRANULES | Use PA Form# 20420 1.Montelukast Granules will only be approved if between ages of 6months-24 months. 2.Singulair Chewables 4mg from 2years-5years and Singulair Chewables 5mgs from 6years-14years old. | |
| ANTIASTHMATIC - ALPHA-PROTEINASE INHIBITOR | | | | MC MC/DEL MC MC | 8 8 8 8 | ARALAST ZEMAIRA GLASSIA PROLASTIN SUSR | Use PA Form# 20420 | Prolastin and Azemaira will be approved for members with A1AT deficiency and clinically demonstrable panacinar emphysema. |
| ANTIASTHMATIC - HYDRO-LYTIC ENZYMES | | | | MC/DEL | | PULMOZYME SOLN | Use PA Form# 20420 | Will be approved for cystic fibrosis patients. |
| ANTIASTHMATIC - MUCOLYTICS | MC/DEL | | ACETYLCYSTEINE ¹ | MC | | MUCOMYST | 1. Acetylcysteine is covered with diagnosis of CF. Use PA Form# 20420 | |
| ANTIASTHMATIC-CFTR POTENTIATOR AND COMBINATIONS | | | | MC MC MC MC MC/DEL | | BRONCHITOL ¹ ORKAMBI KALYDECO SYMDEKO TRIKAFTA | 1. For the treatment of patients ≥18 years of age with CF. Use PA Form# 20420 | <p>Kalydeco will be considered for patients with cystic fibrosis (CF) aged 1 month and older who have at least one mutation in the CFTR gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use.</p> <p>Symdeko will be considered for patients with cystic fibrosis (CF) aged 6 years and older who are homozygous for the <i>F508del</i> mutation or who have at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor based on in vitro data and/or clinical evidence. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use.</p> <p>Bronchitol will be considered as add-on maintenance therapy to improve pulmonary function in adult patients 18 years and older with cystic fibrosis (CF). Use Bronchitol® only for adults</p> |

who have passed the Bronchitol® Tolerance Test (BTT). (see Recommended Dosage section for further information)

Trikafta will be considered for the treatment of cystic fibrosis (CF) in patients aged 2 years and older who have at least one F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene or mutation in the CFTE gene that is responsive based on in vitro data. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to confirm the presence of at least one F508del mutation or a mutation that is responsive based on in vitro data.

Orkambi will be considered for patients with cystic fibrosis (CF) aged 1 year and older who are homozygous for the F508del mutation in the CFTR gene. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of the F508del mutation on both alleles of the CFTR gene. The efficacy and safety of Orkambi have not been established in patients with CF other than those homozygous for the F508del mutation.

[Use PA Form# 20420](#)

| | | | | | | | |
|--------------------------------------|--|--|--|--|--|---|--|
| IDIOPATHIC PULMONARY FIBROSIS | MC/DEL | | OFEV ¹ | MC MC | ESBRIET ¹ PIRFENIDONE | 1. Diagnosis required Use PA Form# 20420 | Ofev- Avoid concomitant use with P-gp and CYP4A inducers (e.g. carbamazepine, phenytoin, and St. John's wort) Esbriet- The concomitant use with strong CYP1A2 inhibitors (e.g. fluvoxamine, enoxacin) is not recommended |
| COUGH/COLD | | | | | | | |
| COUGH/COLD | MC/DEL MC/DEL MC/DEL MC MC | | DEXTROMETHORPHAN CAPS ¹ DEXTRO-GUAIF SYRP ¹ GUAIFENESIN SYRP ¹ PSEUDOEPHEDRINE ¹ ROBITUSSIN DM SYRP ¹ ROBITUSSIN SUGAR FREE SYRP ¹ | | | 1. All of cough cold preparations are not covered except these preferred products. Use PA Form# 20420 | All non-preferred products are not covered as permitted by Federal Medicaid regulations and MaineCare Policy. |
| DIGESTIVE AIDS / ASSORTED GI | | | | | | | |
| GI - ANTIPERISTALTIC AGENTS | MC/DEL MC/DEL MC/DEL MC | | DIPHENOXYLATE DIPHENOXYLATE/ATROPINE LOPERAMIDE HCL CAPS/LIQ OPIUM TINCTURE TINC PAREGORIC TINC | MC/DEL MC MC | LOFENE TABS LONOX TABS MOTOFEN TABS | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. |
| GI - ANTI-DIARRHEAL/ ANTACID - MISC. | MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | | ATROPINE SULFATE SOLN BISMATROL BISMUTH SUBSALICYLATE CALCIUM CARBONATE (ANTACID) CHEW DICYCLOMINE HCL GLYCOPYRROLATE TABS HYOSCYAMINE CAPS & TABS HYOSCYAMINE SULFATE KAOPECTATE MAGNESIUM OXIDE TABS MAG-OX 400 TABS PAMINE TABS PROPANTHELINE BROMIDE TABS SODIUM BICARBONATE TABS TUMS | MC/DEL MC/DEL MC MC MC MC MC/DEL MC MC MC MC MC MC MC MC MC MC MC MC MC | BELLADONNA ALKALOIDS & OP BENTYL TABS BENTYL SYRP CUVPOSA DARTISLA ODT ² ED-SPAZ MYTESI ¹ GLYCOPYRROLATE INJ LEVSIN TABS LEVSIN/SL SUBL NULEV TBDP OSCIMIN ROBINUL INJ ROBINUL TABS | Use PA Form# 20420 1. Dosing limits apply please refer to Dose Consolidation List 2. It is not indicated as monotherapy for treatment of peptic ulcer because effectiveness in peptic ulcer healing has not been established. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. Preferred products that used to require diag codes still require diag codes unless indicated otherwise. Mytesi requires a diagnosis of non-infectious diarrhea in patients with HIV/AIDS on anti-retroviral therapy, prior trials of preferred, more cost effective anti-diarrheals. |
| GI- BILE ACID | | | | MC | CHOLBAM | Use PA Form# 20420 | Indication of bile acid synthesis disorders due to single enzyme defects (SEDs) AND for adjunctive treatment of peroxisomal disorders (PDs) |
| GI- EOSINOPHILIC ESOPHAGITIS | MC | | EOHILIA ¹ | | | Use PA Form# 20420 1. Approvals will not be longer than 12 weeks of treatment in adult and pediatric patients 11 years of age and older | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Eohilia: Dietary modification, PPIs, and topical glucocorticoids are required as initial therapy. |
| GI - H2-ANTAGONISTS | MC MC/DEL MC/DEL | | ACID REDUCER TABS CIMETIDINE FAMOTIDINE | MC MC MC/DEL MC/DEL MC | AXID CAPS AXID AR TABS NIZATIDINE CAPS PEPCID PEPCID AC | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Cimetidine will now be non-preferred and require prior authorization if it is currently being used with any sulfonylurea (except for glyburide). DDI: Cimetidine will require prior authorization if being used in combination with Plavix. |

| | | | | | | | | |
|--|--|--|---|--|---|---|---|--|
| GI - IBAT INHIBITORS | | | | MC MC | | BYLVAY ^{1,2} LIVMARLI ^{1,2} | Use PA Form# 20420 1. For the treatment of patients ≥ 3months of age 2. Clinical PA required for appropriate diagnosis | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. |
| GI - PROTON PUMP INHIBITOR | MC/DEL MC/DEL MC/DEL | | OMEPRAZOLE CAPS ² PANTOPRAZOLE ² LANSOPRAZOLE CAPS ² | MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | 6 6 7 7 8 8 8 8 8 8 8 8 8 8 8 | NEXIUM CPDR ³ NEXIUM SUS ⁵ PRILOSEC OTC ³ ACIPHEX TBEC ³ DEXILANT (KAPIDEX) ² KONVOME ² OMEPRAZOLE-SODIUM BICARBONATE CAPS OMEPRAZOLE MAGNESIUM PREVACID CPDR ³ PREVACID SOLUTABS ^{1,4} PRILOSEC CPDR PROTONIX INJ PROTONIX ² VOQUEZNA TABS | 1. Prevacid Solutabs available without PA for children less than 9 years old. 2. Dosing limits apply, please see dosage consolidation list. 3. All preferreds and step therapy must be tried and failed. 4. Payment for Prevacid SoluTabs for patients 9 and older will be considered for those patients who cannot tolerate a preferred solid oral dosage form. 5. Nexium sus available without PA if member is < 12 yrs of age and ≤ 1 pack per day Use PA Form# 20720 | All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please refer to the PPI PA form for additional criteria on Non-Preferred PPIs DDI: Omeprazole will require prior authorization if being used in combination with Plavix. DDI: Lansoprazole will require prior authorization if being used in combination with Plavix. DDI: Prevacid, Omeprazole and pantoprazole will now be non-preferred and require prior authorization if they are currently being used in combination with any of the following medications: Ampicillin, B-12, Fe salts, Griseofulvin, Sporanox, Ketoconazole, Reyataz, or Vantin. DDI: All non-preferred PPIs require prior authorization, but with any prior authorization request, the member's drug profile will also be monitored for current use with ampicillin, B-12, Fe salts, griseofulvin, itraconazole, ketoconazole, Reyataz or Vantin due to a significant drug-drug interaction. |
| GI - ULCER ANTI-INFECTIVE | MC MC | | PYLERA TALICIA | | | VOQUEZNA DUAL PAK VOQUEZNA TRIPLE PAK | Use PA Form# 20420 | |
| GI - PROSTAGLANDINS | MC | | MISOPROSTOL TABS | MC/DEL | | CYTOTEC TABS | Use PA Form# 20420 | Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drug will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| GI - DIGESTIVE ENZYMES | MC/DEL MC | | CREON ¹ ZENPEP ¹ | MC/DEL MC/DEL MC/DEL | | PERTZYE ULTRESA VIOKACE | Use PA Form# 20420 1. Clinical PA is required to establish CF diagnosis and medical necessity. In all cases except cystic fibrosis patients, objective evidence of pancreatic insufficiency (fat malabsorption test etc...) must be supplied. | Non -Preferred drugs must be tried and failed in step-order due to lack of efficacy or intolerable side effects before other non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| GI - ANTI - FLATULENTS / GI STIMULANTS | MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL | | AMITIZA CALULOSE SYRP CONSTULOSE SYRP ENULOSE SYRP GASTROCROM CONC GENERLAC SYRP LACTULOSE SYRP METOCLOPRAMIDE HCL | MC MC/DEL MC MC/DEL | | CEPHULAC SYRP INFANTS GAS RELIEF SUSP GIMOTI SPRAY REGLAN TABS | | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. |

| | | | | | | | | |
|--------------------------|--------|--|--|--------|--|--|---|---|
| | MC/DEL | | SENOKOT GRAN | MC | | SENOKOT S TABS | | |
| | MC/DEL | | SENOKOT SYRP | MC/DEL | | SORBITOL | | |
| | MC/DEL | | SENOKOT CHILDRENS SYRP | MC | | STOOL SOFTENER PLUS CAPS | | |
| | MC | | SENOKOT XTRA TABS | MC | | SUFLAVE | | |
| | MC/DEL | | STOOL SOFTENER CAPS | MC | | SUTAB | | |
| | MC/DEL | | SUCRALFATE TABS | MC/DEL | | SYMPROIC ³ | | Use PA Form# 20420 |
| | MC/DEL | | SUPREP SOL | MC/DEL | | UNI-CENNA TABS | | |
| | MC | | TRULANCE ² | MC | | UNI-EASE PLUS CAPS | | |
| | MC | | UNI-EASE CAPS | MC | | V-R NATURAL SENNA LAXATIV TABS | | |
| | MC | | URSO FORTE | MC | | URSO 250 | | |
| | MC/DEL | | URSODIOL | MC | | XERMELO ² | | |
| MISC. UROLOGICAL | | | | | | | | |
| UROLOGICAL - MISC. | MC | | ACETIC ACID 0.25% SOLN | MC | | CITRIC ACID/SODIUM CITRAT SOLN | 1. Elmiron requires adequate proof of Dx with supportive testing. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | MC | | CYTRA-K SOLN | MC/DEL | | CYTRA-2 SOLN | | |
| | MC | | FOSFOMYCIN (NDC 82036427401 ONLY) | MC/DEL | | ELMIRON CAPS ¹ | | |
| | MC | | K-PHOS MF TABS | MC | | FURADANTIN SUSP | | Use PA Form# 20420 |
| | MC/DEL | | METHENAMINE MANDELATE TABS | MC/DEL | | MACROBID CAPS | | |
| | MC/DEL | | NEOSPORIN GU IRRIGANT SOLN | MC/DEL | | MACRODANTIN CAPS | | |
| | MC/DEL | | NITROFURANTOIN MONO CAPS | MC/DEL | | NITROFURANTOIN MACR SUSP | | |
| | MC/DEL | | PHENAZOPYRIDINE HCL TABS | MC | | POTASSIUM CITRATE/CITRIC SOLN | | |
| | MC/DEL | | PHENAZOPYRIDINE PLUS | MC/DEL | | PYRIDIUM PLUS TABS | | |
| | MC | | POT CITRATE TAB | MC | | PYRIDIUM TABS | | |
| | MC/DEL | | PROSED/DS TABS | MC/DEL | | RENACIDIN SOLN | | |
| | MC | | TRICITRATES SYRP | MC | | UROCIT-K | | |
| | MC/DEL | | URELIEF PLUS | | | | | |
| | MC | | UREX TABS | | | | | |
| | MC/DEL | | URISED TABS | | | | | |
| | MC/DEL | | UROQID #2 TABS | | | | | |
| PHOSPHATE BINDERS | | | | | | | | |
| PHOSPHATE BINDERS | MC/DEL | | CALCIUM ACETATE CAP ¹ | MC | | AURYXIA ¹ | Use PA Form# 20420 | Preferred drugs must be tried and failed in step-order due to lack of efficacy or intolerable side effects before less preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | MC/DEL | | FOSRENOL CHEW ¹ | MC/DEL | | CALCIUM ACETATE TAB ¹ | 1. Diag required. | |
| | MC/DEL | | MAGNEBIND - 400 ¹ | MC/DEL | | ELIPHOS ¹ | | |
| | MC | | PHOSLYRA ¹ | MC/DEL | | FOSRENOL PWDR ¹ | | |
| | MC/DEL | | REVELA ¹ | MC | | VELPHORO ¹ | | Xphozah to reduce serum phosphorus in adults with chronic kidney disease (CKD) on dialysis as add-on therapy in patients who have an inadequate response to phosphate binders or who are intolerant of any dose of phosphate binder therapy. |
| | MC/DEL | | | MC | | XPHOZAH | | |
| INTRA-VAGINALS | | | | | | | | |
| VAGINAL - ANTIBACTERIALS | MC/DEL | | CLEOCIN CREA | MC/DEL | | METROGEL VAGINAL GEL ¹ | 1. Dosing limits apply, please see Dosage Consolidation List. | Preferred drugs must be tried and failed in step-order due to lack of efficacy or intolerable side effects before less preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | MC/DEL | | CLEOCIN SUPP | MC/DEL | | VANAZOLE | | |
| | MC | | CLINDESSE CREA | MC | | XACIATO | | |
| | MC/DEL | | METRONIDAZOLE VAGINAL GEL ¹ | | | | | |
| | MC/DEL | | NUVESSA | | | | | Use PA Form# 20420 |
| VAGINAL - ANTI FUNGALS | MC/DEL | | CLOTRIMAZOLE CREA | MC | | AVC CREA | 1. Quantity limit: 1/script/2 weeks | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | MC/DEL | | CLOTRIMAZOLE-3 CREA | MC | | CLOTRIMAZOLE 3 DAY CREA | | |
| | MC/DEL | | GYNE-LOTRIMIN CREA | MC | | GYNAZOLE-1 CREA | | |
| | MC | | MICONAZOLE CREA | MC | | GYNE-LOTRIMIN 3 TABS | | |
| | MC | | MICONAZOLE 3 KIT CREA OTC | MC/DEL | | MICONAZOLE 3 COMBO PACK KIT ¹ | | |
| | MC/DEL | | MICONAZOLE 7 CREA | MC/DEL | | MICONAZOLE 3 SUPP | | DDI: Miconazole will require prior authorization if being used in combination with Warfarin. |
| | MC/DEL | | MICONAZOLE NITRATE CREA | MC | | TERAZOL 3 CREA | | |
| | MC | | NYSTATIN TABS | MC | | TERAZOL 7 CREA | | |
| | MC/DEL | | TERCONAZOLE CREAM | MC/DEL | | TERCONAZOLE SUPP | | |
| | MC | | VAGITROL | | | | | |

| | | | | | | | | | |
|---|--|--|---|--|---------------------------------|--|---|--|---|
| | MC | | V-R MICONAZOLE-7 CREA | | | | | | |
| VAGINAL - CONTRACEPTIVES | | | | | | | | | Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drug will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Use PA Form# 20420 |
| VAGINAL - ESTROGENS | MC/DEL MC/DEL | | ESTRING RING PREMARIN CREA | MC/DEL MC/DEL | | ESTRACE CREA ¹ VAGIFEM TABS ¹ | 1. Must fail all preferred products before non-preferred. Use PA Form# 20420 | | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| VAGINAL - OTHER | MC/DEL MC MC | | ACID JELLY GEL ACI-JEL GEL CERVICAL AMINO ACID CREA | MC | | AMINO ACID CERVICAL CREA | Use PA Form# 20420 | | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| BENIGN PROSTATIC HYPERPLASIA (BPH) | | | | | | | | | |
| BPH | MC/DEL MC/DEL MC/DEL MC/DEL | | DOXAZOSIN MESYLATE TABS FINASTERIDE ¹ 5mg TERAZOSIN HCL CAPS TAMSULOSIN HCL | MC/DEL MC/DEL MC MC MC/DEL MC/DEL | 5 8 8 8 8 8 | FLOMAX CP24 ALFUZOSIN AVODART ^{2,4} CARDURA TABS ⁴ ENTADFI ^{5,6} JALYN ^{3,4} PROSCAR TABS ⁴ RAPAFLO ⁴ UROXATRAL ⁴ | 1. There will be dosing limits of 1 tab per day with out PA. 2. Prior use of preferred agent prior to any approvals. 3. Use of preferred (tamsulosin and finasteride) and (tamsulosin and non-preferred Avodart). 4. Non-preferred products must be used in specified order. 5. Use of individual ingredients preferred (Finasteride and tadalafil). 6. Entadfi® is not recommended for more than 26 weeks Use PA Form# 20420 | | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approval of a non-preferred 5-alpha reductase inhibitor requires objective clinical evidence of a very enlarged prostate rather than just the presence of obstructive urinary outflow symptoms along with adequate trial of preferred Proscar. |
| ANXIOLYTICS | | | | | | | | | |
| ANXIOLYTICS - BENZODIAZEPINES | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | | ALPRAZOLAM TABS CHLORDIAZEPOXIDE HCL CAPS CLORAZEPATE DIPOTASSIUM TABS DIAZEPAM LORAZEPAM OXAZEPAM CAPS | MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL | 8 8 8 8 8 8 9 | ALPRAZOLAM ER ATIVAN LOREEV XR NIRAVAM SERAX TRANXENE XANAX TABS XANAX XR | Use PA Form# 20420 | | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ANXIOLYTICS - MISC. | MC/DEL MC MC MC/DEL MC/DEL MC/DEL | | BUSPIRONE HCL TABS HYDROXYZINE HCL SOLN HYDROXYZINE HCL SYRP HYDROXYZINE HCL TABS ¹ HYDROXYZINE PAMOATE CAPS MEPROBAMATE TABS | MC MC MC/DEL MC/DEL | | BUSPAR TABS DROPERIDOL SOLN DROPERIDOL SOLN DROPERIDOL SOLN | Use PA Form# 20420 1. Dosing limits apply, please refer to Dose consolidation list. | | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ANTI-DEPRESSANTS | | | | | | | | | |
| ANTIDEPRESSANTS - MAO INHIBITORS | MC/DEL | | NARDIL TABS | MC/DEL | | TRANLYCYPROMIINE | Use PA Form# 20420 | | |

| | | | | | | | | | | |
|--|--|--|--|--|--------|---|--|---|---|--|
| | | | | MC MC MC/DEL | 8 9 | ZYPREXA RELPREVV ZYPREXA ZYDIS TBDP ¹ SEROQUEL XR | | 3. Dosing limits apply please refer to the dose consolidation list. 4.Requires step through 1 preferred drug for all indications except AMDD. AMDD requires insufficient response from two antidepressants | DDI: The concomitant use of Nuplazid with other drugs known to prolong the QT interval (e.g. Class IA antiarrhythmics, Class 3 antiarrhythmics, antipsychotics, and antibiotics such as gatifloxacin and moxifloxacin). Lybalvi: Step through aripiprazole and Latuda. If criteria is met then initial approval for 3 months. Subsequent approvals will be based on evidence of not gaining >= 10 % baseline body weight for ongoing approval. If weight gain >= 10 % of initial body weight, then criteria for ongoing use not met. Invega Hafyera: The patient is started and stabilized on the medication OR The patient has been adequately treated with Invega Sustenna (paliperidone palmitate 1-month) for at least four months or Invega Trinza (paliperidone palmitate 3- month) following at least one 3-month injection cycle. | |
| ANTIPSYCHOTICS - SPECIAL ATYPICALS | MC/DEL | | CLOZAPINE TABS | MC/DEL MC/DEL MC/DEL | | CLOZAPINE ODT CLOZARIL TABS VERSACLOZ SUSP | | Use PA Form# 20420 | Preferred generic drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred brand will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Patients previously stabilized on brand name drug will be approved. | |
| ANTIPSYCHOTICS - TYPICAL | MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL | | CHLORPROMAZINE HCL FLUPHENAZINE DECANOATE FLUPHENAZINE HCL HALDOL HALOPERIDOL HALOPERIDOL DECANOATE SOLN HALOPERIDOL LACTATE SOLN LOXAPINE SUCCINATE CAPS LOXITANE-C CONC MOBAN TABS PERPHENAZINE PROCHLORPERAZINE SERENTIL THIORIDAZINE HCL THIOTHIXENE TRIFLUOPERAZINE HCL TABS | MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL | | COMPAZINE COMPRO SUPP FLUPHENAZINE HCL CONC HALDOL DECANOATE LOXITANE CAPS MELLARIL NAVANE CAPS PROLIXIN STELAZINE TABS | | Use PA Form# 20420 If prescribing 2 or more antipsychotics, PA will be required for both drugs, except if one is Clozapine. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. If prescribing 2 or more antipsychotics, PA will be required for both drugs, except if one is Clozapine. | |
| LITHIUM | | | | | | | | | | |
| LITHIUM | MC/DEL MC/DEL | | LITHIUM CARBONATE LITHIUM CITRATE SYRP | MC/DEL MC/DEL | | ESKALITH CAPS ESKALITH CR TBCR | | Use PA Form# 20420 | | |
| COMBINATION - PSYCHOTHERAPEUTIC | | | | | | | | | | |
| PSYCHOTHERPEUTIC COMBINATION | MC/DEL MC/DEL | | CHLORDIAZEPOXIDE/AMITRIPT PERPHENAZINE/AMITRIPTYLIN | | | | | Use PA Form# 20420 | | |
| STIMULANTS | | | | | | | | | | |
| STIMULANT - AMPHETAMINES -SHORT ACTING | MC/DEL MC/DEL MC | | AMPHETAMINE SALT COMBO ^{1,4} DEXTROAMPHET SULF TABS PROCENTRA | MC/DEL MC MC/DEL MC | | ADDERALL TABS EVEKEO METHAMPHETAMINE HCL ZENZEDI | | 1. Preferred stimulants will be available without PA if diagnosis of ADHD or Narcolepsy. | | |

| | | | | | | | |
|--|--------------------|--|---|----------------|--|---|---|
| | | | | | | <p>2. As per recent FDA alert, Adderal & Dexedrine should not be used in patients with underlying heart defects since they may be at increased risk for sudden death.</p> <p>3. Dosing limits apply, please see dosing consolidation list.</p> <p>4. Max daily dose of 50mg.</p> <p>Use PA Form# 20420</p> | |
| STIMULANT - LONG ACTING AMPHETAMINES SALT | MC/DEL MC MC | | <p>AMPHETAMINE/DEXTROAMPHET ER^{3,4,7}</p> <p>ADDERALL XR CP24^{1,3,4,7}</p> <p>VYVANSE^{2,3,4}</p> | MC MC MC | | <p>MYDAYIS⁵</p> <p>VYVANSE CHEW^{2,3,4,6}</p> <p>XELSTRYM⁸</p> <p>Use PA Form# 20420</p> <p>1. As per recent FDA alert, Adderall should not be used in patients with underlying heart defects since they may be at increased risk for sudden death.</p> <p>2. FDA approval is currently for adults and children 6 or older. Will be available without PA for this age group if within dosing limits. Limit of one capsule daily. Max dose of 70MG daily.</p> <p>3. Preferred stimulants will be available without PA if diagnosis of ADHD.</p> <p>4. Dosing limits apply, please see dosing consolidation list.</p> <p>5. For the treatment of Attention Deficit Hyperactivity Disorder (ADHD) in patients 13 years and older</p> <p>6. Vyvanse chew grace period for current user through June 2022.</p> <p>7. FDA approval is currently for adults and children 6 or older. Will be available without PA for this age group if within dosing limits. Max dose of 50MG daily without a PA.</p> <p>8. For the treatment of patients 6 years of age and older.</p> | <p>DDI: The concomitant use of Mydayis® is contraindicated with monoamine oxidase inhibitors (MAOIs) or within 14 days after discontinuing MAOI treatment, as concomitant use can increase hypertensive crisis.</p> |
| LONG ACTING AMPHETAMINES | MC MC/DEL | | <p>DEXTROAMPHET SULF CPSR^{1,3}</p> <p>DEXTROAMPHETAMINE ER</p> | MC/DEL MC | | <p>ADZENYS ER³</p> <p>ADZENYS XR- ODT</p> <p>1. Preferred stimulants will be available without PA if diagnosis of ADHD.</p> | |

| | | | | | | | | |
|---|--|--|---|--|--|--|---|--|
| | MC | | DYANAVEL XR SUS | MC MC MC | | ADZENYS XR ³ DEXEDRINE CAP SR ^{2,3} DYANAVEL XR TAB | 2. As per recent FDA alert, Adderall & Dexedrine should not be used in patients with underlying heart defects since they may be at increased risk for sudden death. 3. Dosing limits apply, please see dosing consolidation list. Use PA Form# 20420 | DDI: : The concomitant use of Adzenys® XR is contraindicated with monoamine oxidase inhibitors (MAOIs) or within 14 days after discontinuing MAOI treatment. |
| STIMULANT - METHYLPHENIDATE | MC/DEL MC/DEL MC/DEL MC/DEL | | DEXMETHYLPHENIDATE IR TABS METHYLPHENIDATE SOL METHYLPHENIDATE TAB METHYLIN TABS ^{1,2} | MC/DEL MC/DEL MC MC MC/DEL MC/DEL | | FOCALIN IR TABS METADATE ER METHYLPHENIDATE HCL CHEW METHYLIN CHEWABLES METHYLIN SOL RITALIN | 1. Preferred stimulants will be available without PA if diagnosis of ADHD. Use PA Form# 20420 2. Dosing limits apply, please see dosing consolidation list. Maximum daily doses are as follows: 72mg daily for methylphenidate and 36mg daily for dexmethylphenidate. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please refer to General Criteria category E. |
| STIMULANT - METHYLPHENIDATE - LONG ACTING | MC MC/DEL MC MC MC/DEL | | CONCERTA TBCR DEXMETHYLPHENIDATE CAP ER 50/50 QUILLICHEW ER ^{5,1} QUILLIVANT XR SUS ^{1,5} RITALIN LA ⁴ | MC MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | 5 8 8 8 8 8 8 8 8 8 8 8 | METADATE CD CPR ADHANSIA XR ^{2,6} APTENSIO XR ² AZSTARYS ⁶ COTEMPLA XR ² COTEMPLA XR ODT ² DAYTRANA ^{2,3} FOCALIN XR ² JORNAY PM ^{2,6} METHYLPHENIDATE ER CAPS ^{2,4} METHYLPHENIDATE LA CAPS ² METHYLPHENIDATE ER ^{2,4} CAPS 50/50 METHYLPHENIDATE ER ² CAPS 40/60 METHYLPHENIDATE CD CAPS ² 30-70 | 1. Preferred stimulants will be available without PA if diagnosis of ADHD. 2. Non-preferred products must be used in specified step order. 3. FDA approval currently only for ages 6-16. Limit of one patch daily. Max dose of 30MG daily. 4. Dosing limits apply, please see dosing consolidation list 5. Quillivant XR and Quillichew ER are only indicated for use in patients 6 years of age and older. 6. For the treatment of patients ≥ 6 years of age. Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| STIMULANT - STIMULANT LIKE | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC | | ATOMOXETINE HCL ARMODAFINIL CLONIDINE ER GUANFACINE ER MODAFINIL TABS QELBREE ^{6,7} | MC/DEL MC MC MC/DEL MC MC/DEL MC | 7 7 8 8 8 8 8 8 | PROVIGIL TABS ³ STRATTERA ^{1,2} CAFICIT SOLN ³ INTUNIV KAPVAY SUNOSI WAKIX | 1. Failure of both an amphetamine and methylphenidate is required for consideration for approval of Strattera, unless history of substance abuse without current use of abusable medication(s). Additionally, for patients <17 years of age, a trial of | Provigil requests require diagnosis of Narcolepsy, ADHD, or Obstructive Sleep Apnea. Previous failures of methylphenidate and amphetamine is required for Narcolepsy and ADHD diagnosis, with additional Strattera trial needed with ADHD diagnosis. Please refer to detailed criteria on Provigil PA form Sunsosi is non-preferred and is indicated for to improve wakefulness in adult patients with excessive daytime sleepiness associated with narcolepsy or obstructive sleep apnea (OSA). Wakix is non-preferred and is indicated for the treatment of excessive daytime sleepiness (EDS) in adults with narcolepsy DDI: Sunosi® is contraindicated with MAO inhibitors or within 14 days after discontinuing the MAO inhibitor. |

| | | | | | | | | |
|--|--|--|--|--------|---|---------------------------|---|--|
| | | | | MC | 8 | XYREM SOL | <p>guanfacine in required before approval of Stratterra.</p> <p>2. Stratterra currently has dosing limitations allowing one tablet per day for all strengths if obtain approval. Max daily dose of Stratterra is 100mg. Please see dosing consolidation list.</p> | <p>Xywav: Diagnosis of cataplexy associated with narcolepsy OR excessive daytime sleepiness associated with narcolepsy. Diagnosis must be confirmed by submission of supporting documentation to include the specialist's interpretation of the Polysomnography (PSG) and Multiple Sleep Latency Test (MSLT) results</p> |
| | | | | MC | 8 | XYWAV ⁵ | | |
| | | | | MC/DEL | 9 | NUVIGIL ³ | 3. Non-preferred products must be used in specified | |
| | | | | MC | 9 | DESOXYN TABS ³ | 4. Please use generic Guanfacine. | FDA reminded healthcare professionals and patients that the combined use of Xyrem (sodium oxybate) with alcohol or central nervous system (CNS) depressant drugs can markedly impair consciousness and may lead to severe breathing problems (respiratory depression) |
| | | | | MC | 9 | DESOXYN CR ³ | 5. For patients 7 years of age and older with 6. For pediatric patients 6 years of age or older 7. Preferred with a trial and fail either Atomoxetine OR any 2 preferred ADHD agents. | DDI: Concomitant use of Qelbree® with an MAO inhibitor or within 2 weeks after discontinuing an MAO inhibitor is contraindicated DDI: Concomitant use of Qelbree® significantly increases the total exposure, but not peak exposure, of sensitive CYP1A2 substrates, which may increase the risk of adverse reactions associated with these CYP1A2 substrates. Coadministration of Qelbree® with sensitive CYP1A2 substrates or CYP1A2 substrates with a narrow therapeutic range (e.g. alosetron, duloxetine, ramelteon, tasimelteon, tizanidine, theophylline), is contraindicated. |
| | | | | | | | <p>Use PA Form# 20710 for Provigil, Nuvigil and Xyrem</p> <p>Use PA Form# 20420 for all others</p> | |

ANTI-CATAPLECTIC AGENTS

| | | | | | | | | |
|----------------------------------|--|--|--|----|--|----------|--|---|
| PSYCHOTHERAPEUTIC AGENTS - MISC. | | | | MC | | NUDEXTA | | |
| | | | | MC | | XENAZINE | | |
| | | | | | | | | Use PA Form# 20710 for Xenazine |

WEIGHT LOSS

| | | | | | | | | |
|-------------|--|--|--|--|--|--|--|---|
| WEIGHT LOSS | | | | | | | No longer covered: PHENTERMINE, XENICAL, DIDREX, and MERIDIA | Weight loss drugs are not covered as permitted by Federal Medicaid regulations and Maine Medicaid (MaineCare) Policy. |
|-------------|--|--|--|--|--|--|--|---|

ALZHEIMER DISEASE

| | | | | | | | | |
|------------------------------------|--------|--|---|--------|---|-----------------------------------|--|---|
| ALZHEIMER - Cholinomimetics/Others | MC/DEL | | DONEPEZIL HYDROCHLORIDE TABS ¹ | MC | 6 | ARICEPT TABS ² | 1. PA is required to establish dementia diagnosis and baseline mental status score. | Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | MC/DEL | | DONEPEZIL HYDROCHLORIDE ODT ¹ | MC | 6 | ARICEPT ODT ² | | |
| | MC/DEL | | EXELON DIS ¹ | MC/DEL | 7 | DONEPEZIL HYDROCHLORIDE TABS 23MG | | |
| | MC/DEL | | GALANTAMINE CAPS ¹ | MC | 8 | ADLARITY ³ | | |
| | MC/DEL | | GALANTAMINE TAB ¹ | MC/DEL | 8 | EXELON CAP | 2. Must fail all preferred products before moving to non-preferred. | Kisunla and Leqembi: Testing to rule out reversible causes of dementia (CBC, CMP, TSH, B12, urine drug screen, RPR/VDRL, (folate (if alcohol abuse is present), HIV (if risk present) and an assessment including a review of current medications as a cause of intellectual decline |
| | MC/DEL | | MEMANTINE ¹ | MC/DEL | 8 | GALANTAMINE HYDROBROMIDE SOL | | - Prescribed by or in consultation with a neurologist or geriatrician or geriatric psychiatrist. Diagnosis of Alzheimer's disease defined as: |
| | MC/DEL | | RIVASTIGMINE TARTRATE CAPS ¹ | MC | 8 | KISUNLA | 3. Approvals will require trials and failure or clinical rationale why preferred patches cant be used. | •Confirmed presence of amyloid pathology and mild cognitive impairment or mild dementia stage of disease, consistent with Stage 3 and Stage 4 Alzheimer's disease OR •Confirmed presence of amyloid pathology and prodromal or mild dementia stage of disease, consistent with Stage 3 and Stage 4 Alzheimer's disease |
| | | | | MC | 8 | LEQEMBI ^{1,2} | | -Testing: |
| | | | | MC/DEL | 8 | MEMANTINE HCL SOL | | •Clinical Dementia Rating (CDR) global score of 0.5 or 1.0 OR |
| | | | | MC/DEL | 8 | NAMENDA | | •Repeatable Battery for Assessment of Neuropsychological Status (RBANS) delayed memory index score ≤ 85 OR |
| | | | | MC/DEL | 8 | NAMENDA XR CAPS | | •Mini-Mental State Examination (MMSE) score of 20-30 OR |
| | | | | MC/DEL | 8 | NAMZARIC | | •Montreal Cognitive Assessment (MoCA) score ≤ 22 |
| | | | | MC | 8 | RAZADYNE ² | | - Member is age 50 or older - Obtain recent (within one year) brain magnetic resonance imaging (MRI) prior to initiating treatment - Provider attestation to obtain MRIs prior to the 7th infusion (first dose of 10 mg/kg) and 12th infusion (sixth dose of 10 mg/kg) |

| | | | | | | | | |
|---------------------------------|--|--|--|--|------------------|---|---|---|
| | | | | MC | 9 | COGNEX CAPS ² | | <p>Member does NOT have history or increased risk of amyloid related imaging abnormalities-edema (ARIA-E), which includes brain edema or sulcal effusions and amyloid related imaging abnormalities hemosiderin deposition (ARIA-H), which includes microhemorrhage and superficial siderosis</p> <p>- Member does NOT have hypersensitivity to any components of these drugs</p> <p>- Failure of or inability to tolerate at least two other preferred Alzheimer therapies for at least four months each, one of which should include a combination of a cholinesterase inhibitor with memantine</p> <p>*If the initial drug utilized is the combination of a cholinesterase inhibitor and memantine, then only that single trial of two drugs is required</p> |
| SMOKING CESSATION | | | | | | | | |
| NICOTINE PATCHES / TABLETS | MC/DEL MC/DEL MC/DEL MC/DEL | | CHANTIX TAB ¹ CHANTIX STARTER PACK NICOTINE DIS PT24 ¹ VARENICLINE TAB | MC/DEL | | NICODERM CQ PT24 ¹ | Use PA Form# 20420 1. See criteria section for exemptions | <p>As of July 1, 2014 per MaineCare policy, smoking cessation products will be covered without a copay(including MEDEL). No annual or lifetime limits, must follow FDA approved indications and therapy guidelines.</p> <p>Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>Note: MaineCare policy, smoking cessation product were "not covered" except for during pregnancy between 9/1/12 and 1/1/14, between 1/1/2014 and 7/1/14 smoking cessation products were covered with limitations</p> <p>Patients may qualify for the medication through The Maine Tobacco Helpline if they do not have MaineCare or MEDEL. Patients are encouraged to call The Maine Tobacco helpline at 1-800-207-1230.</p> |
| NICOTINE REPLACEMENT - OTHER | MC/DEL MC/DEL MC/DEL | | NICOTINE POLACRILEX GUM ¹ NICOTINE LOZENGE MINI NICOTINE LOZENGE | MC/DEL MC/DEL MC/DEL MC | 8 8 8 8 | NICOTROL INHALER ^{1,2} NICOTROL NASAL SPRAY ^{1,2} NICORETTE GUM ^{1,2} NICORETTE LOZENGES | Use PA Form# 20420 1. See criteria section for exemptions 2. Must use non-preferred products in specified step order. | <p>As of July 1, 2014 per MaineCare policy, smoking cessation products will be covered without a copay(including MEDEL). No annual or lifetime limits, must follow FDA approved indications and therapy guidelines.</p> <p>Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>Note: MaineCare policy, smoking cessation product were "not covered" except for during pregnancy between 9/1/12 and 1/1/14, between 1/1/2014 and 7/1/14 smoking cessation products were covered with limitations</p> <p>Patients may qualify for the medication through The Maine Tobacco Helpline if they do not have MaineCare or MEDEL. Patients are encouraged to call The Maine Tobacco helpline at 1-800-207-1230.</p> |
| ALCOHOL DETERRENTS | | | | | | | | |
| ALCOHOL DETERRENTS | MC/DEL MC MC MC/DEL | | ACAMPROSATE ANTABUSE TABS DISULFIRAM TABS NALTREXONE HCL TABS | MC/DEL | | ACAMPRO ¹ | 1. Should only be used in conjunction with formal structured outpatient detoxification program. Use PA Form# 20420 | Preferred generic drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| MISCELLANEOUS ANALGESICS | | | | | | | | |
| ANALGESICS - MISC. | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL | | ACETAMINOPHEN ASPIRIN ASPRIN/ APAP/ CAFF TAB BUTAL/ASA/CAFF BUTALBITAL COMPOUND BUTALBITAL/ACET TABS BUTALBITAL/APAP CAPS BUTALBITAL/APAP/CAFFEINE TABS CHOLINE MAGNESIUM TRISALI DIFLUNISAL TABS EXCEDRIN SALSALATE TABS | MC MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC MC MC | | AXOCET CAPS ESGIC-PLUS FIORICET TABS FIORINAL CAPS FIORTAL CAPS FORTABS TABS PHRENILIN TABS PHRENILIN FORTE CAPS TRILISATE LIQD TRILISATE TABS ZEBUTAL CAPS ZORPRIN TBCR | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| LONG ACTING NARCOTICS | | | | | | | | |
| NARCOTICS - LONG ACTING | MC/DEL MC/DEL | | FENTANYL PATCH ⁴ BUTRANS ⁴ | MC MC | 8 8 | ARYMO ER AVINZA | Use PA Form# 20510 Use PA form #10300 for | Preferred drugs (Fentanyl Patch, Morphine Sulfate ER tab, Butrans and Embeda) must be tried for at least 2 weeks each & failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Adequate clinical dose/ratio/treatment of non-preferred drug |

Beginning January 2017, all current opiate users who are above the maximum combined daily dose of 100 MME must titrate their total daily dose of opioid medications below 300 MME. Also, the maximum daily supply of an opiate prescription for acute pain will be limited to 7-day supplies. The maximum day supply of an opiate prescription for chronic pain will be limited to 30-day supplies. As of July 1, 2017 all users of opioid medications must comply with the maximum combined daily dose of 100 MME.

However, for MaineCare members, effective January 1, 2017, opioid prescription(s) for more than a 7-day supply and/or more than 30 MME/ day will require a prior authorization. Please note that MaineCare implemented a 30 MME limit January 1, 2013 that is still effective.

Post-surgical members may receive prior authorizations for opiates up to a 60 days in length if medical necessity is provided by the surgical provider.

An MME conversion chart is available at www.mainearepd.org. Click on "General Pharmacy Info."

Please see the Pain Management Policy tab for the complete criteria

MISCELLANEOUS NARCOTICS

| NARCOTICS - MISC. | | | | | | |
|-------------------|--|--------|---|----------------------------------|---|---|
| MC/DEL | ACETAMINOPHEN/CODEINE | MC/DEL | 8 | ABSTRAL | 1. Fentanyl OT loz (Barr) | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please refer to General Criteria category E. |
| MC/DEL | ASPIRIN/CODEINE TABS | MC/DEL | 8 | APADAZ | and Capital and codeine suspension products require PA for users over 18 years of age. PA is not required if under 18 years of age. | |
| MC/DEL | BUTAL/ASA/CAFF/COD CAPS | MC/DEL | 8 | ASCOMP/CODEINE CAPS | | |
| MC | BUTALBITAL/ASPIRIN/CAFFEI CAPS | MC/DEL | 8 | BUTALBITAL/APAP/CAFFEINE/ CAPS | | |
| MC | CAPITAL AND CODEINE SUSP ¹ | MC/DEL | 8 | BUTALBITAL COMPOUND- CODEINE CAP | | |
| MC | CAPITAL/CODEINE SUSP ¹ | MC | 8 | DEMEROL | | |
| MC/DEL | CODEINE PHOSPHATE SOLN | MC/DEL | 8 | DILAUDID | | Beginning January 2017, all current opiate users who are above the maximum combined daily dose of 100 MME must titrate their total daily dose of opioid medications below 300 MME. Also, the maximum daily supply of an opiate prescription for acute pain will be limited to 7-day supplies. The maximum day supply of an opiate prescription for chronic pain will be limited to 30-day supplies. As of July 1, 2017 all users of opioid medications must comply with the maximum combined daily dose of 100 MME. |
| MC/DEL | CODEINE SULFATE TABS | MC | 8 | DILAUDID-HP SOLN | 2. Oxycodone/acet 10/650 is 8 times more expensive. Use twice as many of oxycod/acet 5/325 instead. | |
| MC/DEL | ENDOCET TABS ³ | MC | 8 | FENTANYL CITRATE SOLN | You can mix andmatch preferred strengths of oxycodone and oxycodone/acet to minimize acet. dose similar to certain non-preferred drugs. | |
| MC/DEL | ENDODAN TABS | MC/DEL | 8 | FENTORA | | However, for MaineCare members, effective January 1, 2017, opioid prescription(s) for more than a 7-day supply and/or more than 30 MME/ day will require a prior authorization. Please note that MaineCare implemented a 30 MME limit January 1, 2013 that is still effective. |
| MC/DEL | FENTANYL OT LOZ ¹ | MC/DEL | 8 | FIORICET/CODEINE CAPS | | |
| MC/DEL | FENTANYL OT LOZ1 | MC | 8 | FIORINAL/CODEINE #3 CAPS | | |
| MC/DEL | HYDROCODONE/ACETAMINOPHEN | MC | 8 | FIORTAL/CODEINE CAPS | | |
| MC/DEL | HYDROMORPHONE HCL ³ | MC/DEL | 8 | HYDROCODONE/IBUPROFEN | | Post-surgical members may receive prior authorizations for opiates up to a 60 days in length if medical necessity is provided by the surgical provider. |
| MC | LORTAB ELX | MC/DEL | 8 | HYDROMORPHONE ER | | |
| MC/DEL | MEPERIDINE SOL | MC/DEL | 8 | HYDROMORPHONE RECTAL SUPP | | An MME conversion chart is available at www.mainearepd.org . Click on "General Pharmacy Info." |
| MC/DEL | NUCYNTA | MC | 8 | IBUDONE | | |
| MC/DEL | OXYCODONE TAB | MC/DEL | 8 | LEVORPHANOL TARTRATE TAB | | |
| MC/DEL | OXYCODONE/ACETAMINOPHEN ^{2,3} | MC/DEL | 8 | LORCET | 3. Only preferred manufacturer's products will be available without prior authorization. | |
| MC/DEL | ROXICET | MC | 8 | LORTAB | | |
| MC | ROXIPRIN TABS | MC | 8 | MAXIDONE TABS | | |
| | | MC/DEL | 8 | MEPERIDINE TABS | | Please see the Pain Management Policy for the complete criteria |
| | | MC/DEL | 8 | NORCO TABS | | |
| | | MC/DEL | 8 | ONSOLIS | | |
| | | MC/DEL | 8 | OXECTA | | |
| | | MC/DEL | 8 | OXYCODONE CAP | | |
| | | MC/DEL | 8 | OXYCODONE/APAP 10/650 | | |
| | | MC/DEL | 8 | OXYCODONE/APAP 7.5/500 | | |
| | | MC/DEL | 8 | PENTAZOCINE/ACET TABS | | |
| | | MC/DEL | 8 | PENTAZOCINE/NALOXONE TABS | | |
| | | MC | 8 | PERCOCET TABS | | |
| | | MC | 8 | PERCOCET TABS | | |
| | | MC | 8 | PHRENILIN W/CAFFEINE/CODE CAPS | | |
| | | MC/DEL | 8 | ROXICET 5/500 TABS | | |
| | | MC | 8 | ROXICODONE TABS | | |
| | | MC/DEL | 8 | ROXYBOND | | |
| | | MC | 8 | SYNALGOS-DC CAPS | | |
| | | MC | 8 | TALACEN TABS | | |

| | | | | | | | | |
|------------------------------|--------|--|--|--------|---|------------------------------|---|--|
| | | | | MC | 8 | TREZIX | | |
| | | | | MC | 8 | TYLENOL/CODEINE #3 TABS | | |
| | | | | MC | 8 | TYLOX CAPS | | |
| | | | | MC | 8 | XOLOX | Use PA Form# 20420 | |
| | | | | MC | 8 | VICODIN | | |
| | | | | MC | 8 | VICOPROFEN TABS | Use PA form #10300 for PAs over the opiate limit | |
| | | | | MC | 8 | ZYDONE TABS | | |
| | | | | MC | 9 | ACTIQ LPOP | | |
| | | | | MC | 9 | CONZIP | | |
| | | | | MC | 9 | OPANA | | |
| OPIOID DEPENDENCE TREATMENTS | MC | | SUBOXONE FILM ² | | | | | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | MC/DEL | | BUPRENORPHINE/NALOXONE TABS ² | MC | | BRIXADI | Use PA Form #20100 | Members will continue to be required to follow the criteria listed below: |
| | | | | MC/DEL | | BUPRENORPHINE ^{1,2} | | 1-Induction period for 30 days |
| | | | | MC | | SUBLOCADE | 1. Buprenorphine will only be approved for use during pregnancy. | 2-Members will be allowed multiple induction periods per year where they can receive max 24 mg Daily for up to 30-days, without a PA once they have been on a maintenance dose. |
| | | | | MC | | ZUBSOLV | 2. See Criteria Section | 3-Max dose of 24 mg for maintenance |
| | | | | | | | | 4-Max dose of 16 mg for maintenance |
| | | | | | | | | 5- Suboxone will not require a PA if patient requires concomitant use of an opioid for acute pain. |
| | | | | | | | | 7- Buprenorphine monotherapy is preferred if member is pregnant and dose not > 16 mg day and pregnancy diagnosis is noted on the prescription. |
| | | | | | | | Use PA form #20200 for Extended Release Buprenorphine | Brixadi and Sublocade: The prescriber can attest (and medical record should document) that: -member has a documented history of opioid use disorder (OUD), -XRB is being used for the treatment of OUD (rather than pain or any other non-FDA approved indication) and -member's total daily dose of sublingual buprenorphine is less than or equal to 24 mg daily. AND at least one of the following is true: -The member's previous use of sublingual buprenorphine has included misuse, overuse, or diversion. -The member is at high risk of overdose (e.g., individuals leaving incarceration or abstinence-based treatment programs; individuals who are unhoused; or those facing potential gaps in care due to delays in care or geographically limited treatment access). -The member has experienced significant medical complications of OUD and/or of injection drug use. Occurrence should be in the last 5 years, or it should be clearly documented that the risk indicated by this infection or complication is ongoing (Examples of medical complications of OUD include: threatened the function of organs or life or limb threatening and required medical and/or surgical therapy. Examples of medical complications of injection drug use include osteomyelitis, endocarditis, renal failure, joint infection or other serious medical complications directly related to OUD.) -The member has treatment-resistant OUD, including those with ongoing illicit substance use in the context of sublingual buprenorphine treatment as documented by positive urine drug screens or other clear objective evidence, and/or further functional decline with explicit documentation of the functional decline. -The member has a significant intolerance of, or documented allergy to, sublingual buprenorphine (either buprenorphine monotherapy or buprenorphine/naloxone combination therapy) that has resulted in the patient's inability to comply with continued treatment using the sublingual product. (A true allergy is usually accompanied by rash, respiratory symptoms, or anaphylaxis. Other complaints such as bad taste, mouth tingling, etc. do not constitute evidence of allergy or significant intolerance. Formulation preference or convenience are not, in and of themselves, indications for using XRB.) -The member is in ongoing treatment with XRB and would like to continue the medication. |
| OPIOID WITHDRAWAL AGENTS | | | | MC | | LUCEMYRA ¹ | 1. Clinical PA for appropriate approved use and patient has documented contraindication to clonidine Use PA Form#20420 | |
| NARCOTIC ANTAGONISTS | | | | | | | | |
| NARCOTIC - ANTAGONISTS | MC/DEL | | NALTREXONE HCL TABS | MC | | EVZIO | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| | MC | | NALOXONE INJ | MC | | OPVEE ² | | |
| | MC | | NARCAN NS | MC | | KLOXXADO | 1. Will only be approved for | |

| | | | | | | |
|---|--|--|--|--|---|---|
| | MC MC MC | NALOXONE SPRAY OTC VIVITROL INJ ZIMHI | MC/DEL | REVIA TABS ¹ | side effects experienced with generic that are not described in the literature as occurring with the brand version. 2. For the treatment of adult and pediatric patients 12 years of age and older. | |
| COX 2 / NSAIDS | | | | | | |
| COX 2 INHIBITORS - SELECTIVE / HIGHLY SELECTIVE | MC/DEL MC/DEL MC/DEL MC/DEL | CELECOXIB ^{4,5} KETOROLAC TROMETHAMINE ^{2,3,5} NABUMETONE TABS ⁵ MELOXICAM TABS ^{1,5} | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | CELEBREX CAPS ^{4,5} MELOXICAM CAPS ⁵ MOBIC ⁵ MOBIC SUSP ⁵ RELAFEN TABS ⁵ QMIIZ ODT VIVLODEX | <p>Use PA Form# 20420</p> <p>1. Meloxicam has dosing limits allowing one tablet daily of all strengths without PA.</p> <p>2. Ketorolac Tromethamine is indicated for the short term (up to 5 days) management of moderately severe acute pain that requires analgesic at the opioid level in adults. Not indicated for minor or chronic pain conditions.</p> <p>3. Ketorolac has dosing limits allowing 24 tablets for a 5 day supply every 30 days.</p> <p>4. Dosing limits will be set at a maximum of 400mg daily</p> <p>5. The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use.</p> | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| NSAIDS | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | CHILDRENS IBUPROFEN DICLOFENAC POTASSIUM TABS DICLOFENAC SODIUM TABS DICLOFENAC SODIUM 1% GEL ¹ ETODOLAC FENOPROFEN CALCIUM TABS FLURBIPROFEN TABS IBUPROFEN INDOMETHACIN KETOPROFEN MECLOFENAMATE SODIUM CAPS NAPROSYN SUSP NAPROXEN SUSP NAPROXEN TABS NAPROXEN SODIUM TABS NAPROXEN SODIUM CAPS NAPROXEN DR TBEC | MC MC MC MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC | ADVIL TABS ANAPROX TABS ANAPROX DS TABS CAMBIA CATAFLAM TABS CHILDRENS ADVIL SUSP CHILD'S IBUPROFEN SUSP CHILDREN'S MOTRIN SUSP CLINORIL TABS DAYPRO TABS DICLIFENAC GEL EC-NAPROSYN TBEC ETODOLAC ER 600MG FELDENE CAPS FLECTOR PATCH IBU-200 INDOCIN | <p>The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use.</p> <p>1. Dosing limits apply, please see Dosage Consolidation List.</p> <p>Use PA Form# 20420</p> | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approvals will be granted for other requests based on failure of at least one generic NSAID from at least 3 different NSAID classes as described in the COX-II PA form. DDI: Diclofenac will now be non-preferred and require prior authorization if it is currently being used in combination with Ilescol. The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use. |

[Use PA Form# 20420](#)

MISCELLANEOUS ARTHRITIS

| | | | | | | | | |
|-------------------|----------|--|----------------------------------|--------|--|------------------------|---|---|
| ARTHRITIS - MISC. | MC MC | | RIDAURA CAPS MYOCHRYSINE SOLN | MC/DEL | | ARTHROTEC ¹ | 1. The individual components of Arthrotec are available without PA. Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. The individual components of Arthrotec are available without PA. |
|-------------------|----------|--|----------------------------------|--------|--|------------------------|---|---|

LUPUS-SLE

| | | | | | | | | |
|-----------|--|--|--|----------------|--|---|--|---|
| LUPUS-SLE | | | | MC MC MC | | BENLYSTA ¹ LUPKYNIS SAPHNELO | Use PA Form# 20420 1. Approvals will require previous trial of corticosteroids, antimalarials, NSAIDS and immunosuppressives. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Lupkynis is a sensitive CYP3A4 substrate. Co-administration with strong or moderate CYP3A4 inhibitors increases voclosporin exposure, which may increase the risk of Lupkynis® adverse reactions. Co-administration of Lupkynis® with strong CYP3A4 inhibitors (e.g. ketoconazole, itraconazole, clarithromycin) is contraindicated. Reduce Lupkynis® dosage when co-administered with moderate CYP3A4 inhibitors (e.g. verapamil, fluconazole, diltiazem) |
|-----------|--|--|--|----------------|--|---|--|---|

PIK3CA-Related Overgrowth Spectrum (PROS)

| | | | | | | | | |
|---|--|--|--|----|--|----------------------|--|--|
| PIK3CA-Related Overgrowth Spectrum (PROS) | | | | MC | | VIJOICE ¹ | Use PA Form# 20420 1. PA required to confirm FDA approved indication. | Preferred drugs must be tried and failed, in step-order, due to lack of efficacy (failure to reach target IOP reduction) or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
|---|--|--|--|----|--|----------------------|--|--|

MIGRAINE THERAPIES

| | | | | | | | | |
|-----------------------------------|--|--|--|--------------|--|----------------------------|------------------------------------|--|
| MIGRAINE - ERGOTAMINE DERIVATIVES | | | | MC/DEL MC | | D.H.E. 45 SOLN TRUDHESA | Use PA Form# 10110 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
|-----------------------------------|--|--|--|--------------|--|----------------------------|------------------------------------|--|

| | | | | | | | | |
|--|----|--|--------------------|----|--|------------------|------------------------------------|--|
| MIGRAINE - CARBOXYLIC ACID DERIVATIVES | MC | | DIVALPROEX ER TB24 | MC | | DEPAKOTE ER TB24 | Use PA Form# 10110 | |
|--|----|--|--------------------|----|--|------------------|------------------------------------|--|

| | | | | | | | | |
|---|--|---------------------------------|---|--|--|---|--|--|
| MIGRAINE - SELECTIVE SEROTONIN AGONISTS (5HT)--Tabs/Nasal | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | 1 1 1 1 1 1 2 | MIGRANAL NASAL SPRAY RELPAX ¹ RIZATRIPTAN ODT RIZATRIPTAN TABS SUMATRIPTAN TABS ¹ ZOLMITRIPTAN TAB ¹ NARATRIPTAN HCI TABS ¹ | MC MC MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | | AMERGE TABS ^{1,2} AXERT TABS ^{1,2} FROVA TABS ^{1,2} IMITREX NASAL SPRAY ¹ IMITREX TABS ^{1,2} MAXALT ^{1,2,3} MAXALT MLT ^{1,2,3} ONZETRA XSAIL ² SUMATRIPTAN NASAL SPRAY ¹ ZOLMITRIPTAN ODT ZOLMITRIPTAN SPRAY ZOMIG TABS ^{1,2} ZOMIG NASAL SPARY ^{1,2} ZOMIG ZMT TBDP ^{1,2} | 1. All drugs in this category have dosing limits. Please refer to dose consolidation table. 2. Must fail all preferred products before non-preferred. 3. Established users will be grandfathered Use PA Form# 10110 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Quantity limit exceptions will require ongoing therapy with therapeutic doses of highly effective prophylactic medication as listed on the Triptan PA form. |
|---|--|---------------------------------|---|--|--|---|--|--|

| | | | | | | | | |
|--|------------------------|--|---|--------------------|--|---|---|--|
| MIGRAINE - SELECTIVE SEROTONIN AGONISTS (5HT)--Injectables | MC MC/DEL MC/DEL | | IMITREX CARTRIDGE ¹ SUMATRIPTAN SYRINGE ¹ SUMATRIPTAN PEN INJCTR ¹ | MC/DEL MC MC | | TOSYMRA ZEMBRACE ¹ IMITREX PEN INJCTR ¹ | Use PA Form# 10110 1. Dosing limits apply. Please refer to the dose consolidation table. | |
|--|------------------------|--|---|--------------------|--|---|---|--|

| | | | | | | | | |
|---|--|--|--|--------|--|-------------------------|---|--|
| MIGRAINE - SELECTIVE SEROTONIN AGONISTS (5HT)--Combinations | | | | MC/DEL | | TREXIMET ^{1,2} | Use PA Form# 10110 1. Dosing limits apply. Please see dose consolidation list. | |
|---|--|--|--|--------|--|-------------------------|---|--|

| | | | | | | | | |
|---|--|--|--|--|--|---|---|--|
| | | | | | | 2. Use preferred Sumatriptan and Naproxen separately. Treximet only available if component ingredients of sumatriptan and naproxen are unavailable. | | |
| MIGRAINE - MISC. | MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL | | AIMOVIG ¹ AJOVY ¹ AJOVY AUTO INJCT ¹ EMGALITY SYRINGE ¹ 200mg/ml EMGALITY PEN ¹ NURTEC ODT ² SPASTRIN TABS | MC MC MC/DEL MC/DEL MC MC MC MC/DEL | | BELCOMP-PB SUPP ELYXYB MIGRAZONE CAPS MIGERGOT SUP QULIPTA REYVOW ² UBRELVY ² VYEPTI ² ZAVZPRET ² | Use PA Form# 10110 1. See criteria section 2. Dosing limits apply, please see the dose consolidation list. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Aimovig, Ajovy and Emgality: The patient is 18 years of age or older AND patient has a diagnosis of episodic migraine (4-14 headache days per month with migraine lasting 4 hours or more) or chronic migraine (≥ 15 headache days per month, of which ≥ 8 are migraine days, for at least 3 months) AND patient has failed or has a contraindication to an adequate trial (≥ 60 days) of at least 2 medications for migraine prophylaxis from at least 2 different classes. Ubrely is non-preferred and is indicated for the acute treatment of migraine with or without aura in adults. This is not indicated for the preventive treatment of migraine. Reyvow is non-preferred and is indicated for the acute treatment of migraine with or without aura in adults. Reyvow® is not indicated for the preventive treatment of migraine. Zavzpret: The patient must have a documented side effect, allergy, or treatment failure to preferred oral CGRP Inhibitor and two non-preferred oral CGRP Inhibitors. Nurtec ODT will be preferred after 2 adequate trials of at least two preferred triptans |
| GOUT | | | | | | | | |
| GOUT | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | | ALLOPURINOL TABS COLCHICINE TAB FEBUXOSTAT TAB PROBENECID TABS PROBENECID/COLCHICINE TABS | MC/DEL MC MC MC/DEL MC MC | | COLCHICINE CAP COLCRYS GLOPERBA ULORIC ¹ MITIGARE ZYLOPRIM TABS | Use PA Form# 20420 1. Failure of therapeutic (300mg) dose of Allopurinol (failure define as not being able to get uric acid levels below 6mg/dl) or severe renal disease. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: The concomitant use of Gloperba® and CYP3A4 inhibitors (e.g. clarithromycin, ketoconazole, grapefruit juice, erythromycin, verapamil, etc.) should be avoided due to the potential for serious and life-threatening toxicity. |
| MISC. | | | | | | | | |
| ACID SPHINGOMYELINASE DEFICIENCY (ASMD) | | | | MC | | XENPOZYME ^{1,2} | 1.For treatment of non-central nervous system manifestations of acid sphingomyelinase deficiency (ASMD) in adult and pediatric patients 2. Clinical PA required for appropriate diagnosis and clinical parameters. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ANESTHETICS - MISC. | MC MC MC | | BUPIVACAINE HCL SOLN LIDOCAINE HCL SOLN MARCAINE SOLN | MC MC/DEL MC | | SENSORCAINE-MPF SOLN SYNVISC INJ XYLOCAINE SOLN | Use PA Form# 30130 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| COLD AGGLUTININ DISEASE (CAD) | | | | MC | | ENJAYMO ¹ | 1. Indicated to decrease the need for red blood cell transfusion due to hemolysis in adults with cold agglutinin disease (CAD). | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| PRIMARY HYPEROXALURIA TYPE 1 (PH1) | | | | | | OXLUMO ¹ RIVFLOZA | 1. PA is required to establish diagnosis and medical | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the |

| | | | | | | | | |
|--|----------------------------------|--|--|--------------------------------|------------------|--|---|---|
| | | | | | | | Use PA Form# 20420 | preferred drug(s) exists. |
| | | | | | | | | Rivfloza: The patient has a diagnosis of Primary Hyperoxaluria Type I (PH1) confirmed via genetic testing (identification of alanine: glyoxylate aminotransferase gene (AGXT) mutation) AND urinary oxalate excretion > 0.5mmol/1.73 m2 or urinary oxalate: creatinine ratio is above the upper limit of normal for age AND is at least 9 years of age AND medication is being prescribed by, or in consultation, with a nephrologist or urologist |
| SICKLE CELL DISEASE | MC/DEL MC | | HYDROXYUREA DROXIA | MC MC MC MC MC/DEL | | ADAKVEO CASGEVY ^{2,3} ENDAR ¹ LYFGENIA ^{2,3} OXBRYTA ² SIKLOS | 1.Evidence of other preferred L-glutamine products utilization and reason for failure. 2. For the treatment of patients ≥ 12 years of age. 3. PA required to confirm FDA approved indication. Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: The concomitant use of Oxbryta and strong CYP3A4 inhibitors or fluconazole may increase voxelotor plasma levels and may lead to increased toxicity. |
| HUTCHINSON- GILFORD PROGERIA SYNDROME (HGPS) | | | | MC | | ZOKINVY ^{1,2} | 1.In patients 12 months of age and older with a body surface area (BSA) of 0.39m2 and above 2. PA required to confirm FDA approved indication. Use PA Form# 20420 | ZOKINVY: To reduce the risk of mortality in Hutchinson-Gilford Progeria Syndrome (HGPS). For the treatment of processing-deficient Progeroid Laminopathies with either: Heterozygous LMNA mutation with progerin-like protein accumulation OR Homozygous or compound heterozygous ZMPSTE24 mutations |
| VACCINES | MC/DEL MC MC/DEL MC/DEL | | ABRYSVO AREXVY GARDASIL 9 SHINGRIX | | | | Use PA Form# 20420 | Gardasil 9 will be preferred by MaineCare for ages 19-45 for FDA approved indications. Under the Maine Immunization Program Gardasil 9 is covered under the Vaccine for Children Program for ages 9-18. Please contact 1-800-867-4775 or 207-287-3746 for assistance. Abrysvo will be a preferred vaccine indicated for active immunization for the prevention of lower respiratory tract disease (LRTD) caused by respiratory syncytial virus (RSV) in individuals 60 years of age and older. Active immunization of pregnant individuals at 32 through 36 weeks gestational age for the prevention of LRTD and severe LRTD caused by RSV in infants from birth through 6 months of age. Arexvy will be preferred for active immunization for the prevention of LRTD caused by respiratory syncytial virus (RSV) in individuals 60 years of age and older. SHINGRIX (>= 50yo) is preferred as of 11-20-20 with respective age edit. |
| APDS | | | | MC | | JOENJA ^{1,2,3} | Use PA Form# 20420 1.Clinical PA required for appropriate diagnosis 2. For the treatment of patients 2 years of age and older. 3. Avoid CYP3A drug drug interaction. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ALPHA- MANNOSIDOSIS | | | | MC | | LAMZEDE | Use PA Form# 20420 1.Clinical PA required for appropriate diagnosis | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ANTI-CONVULSANTS | | | | | | | | |
| ANTICONVULSANTS | MC/DEL MC MC/DEL MC/DEL | | CARBAMAZEPINE CARBAMAZEPINE ER CAP CARBATROL CP12 CELONTIN CAPS | MC MC MC/DEL MC | 8 8 8 8 | APTIOM BANZEL BRIVIACT ⁷ CARBAMAZEPINE SUS | Use PA Form# 20420 All non-preferred meds must be used in specified order | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |

| | | | | | | |
|--------|-----------------------------|--------|---|--|--|--|
| MC/DEL | CLOBAZAM | MC | 8 | DEPAKOTE | | |
| MC/DEL | CLONAZEPAM TABS | MC | 8 | DEPAKOTE ER | | |
| MC | DEPAKOTE SPRINKLES CPSP | MC | 8 | DIACOMIT | 1. Quantity limit. 5/month | |
| MC/DEL | DIASTAT ¹ | MC/DEL | 8 | DIVALPROEX SODIUM SPRINKLE CAPS | 2. Dosing limits apply, please see dose consolidation list. | Approvals will be for patients with a variety of drug-specific FDA-approved indications and for specific conditions supported by at least two published peer-reviewed double-blinded, placebo-controlled randomized trials that are not contradicted by other studies of similar quality after recommendation by the DUR Committee and as long as all first line therapies have been tried and failed at full therapeutic doses for adequate durations (at least two weeks). |
| MC/DEL | DIAZEPAM GEL ¹ | MC | 8 | ELEPSIA XR ¹⁰ | | |
| MC/DEL | DILANTIN | MC | 8 | EPRONTIA SOLN ¹¹ | | |
| MC/DEL | DIVALPROEX SODIUM | MC/DEL | 8 | FELBATOL | 3. Dosing limits apply per strength as well as a maximum daily dose of 600mg. Please see dose consolidation list. | |
| MC | DIVALPROEX SPRINKLE CAP | MC/DEL | 8 | FELBATOL SUS | | |
| MC/DEL | EPIDIOLEX ⁸ | MC/DEL | 8 | FELBAMATE SUS | | |
| MC/DEL | EPITOL TABS | MC | 8 | FINTEPLA ⁹ | | |
| MC/DEL | ETHOSUXIMIDE SYRP | MC | 8 | FYCOMPA ² | | |
| MC/DEL | EQUETRO | MC/DEL | 8 | HORIZANT | 4. Adjunctive therapy 17 and older. | *** SEE CHART AT END OF DOCUMENT |
| MC/DEL | GABAPENTIN ² CAP | MC | 8 | GRALISE | | |
| MC/DEL | GABAPENTIN ² TAB | MC/DEL | 8 | KEPPRA TABS | 5. Max dose 2400mg | |
| MC/DEL | GABAPENTIN SOL | MC/DEL | 8 | KEPPRA SOLN | 6. Clinical PA required for appropriate diagnosis | Topamax and Neurontin - Second line therapy for migraine prophalaxis after trial of at least three preferred preventive medications from Group 1 listed on page 2 of the Acute Migraine PA form. |
| MC/DEL | GABITRIL TABS | MC/DEL | 8 | KLONOPIN TABS | | |
| MC/DEL | LACOSAMIDE SOL | MC | 8 | LAMICTAL IR | | |
| MC/DEL | LACOSAMIDE TAB | MC | 8 | LAMICTAL ODT | | |
| MC | LAMICTAL CHEW | MC/DEL | 8 | LEVETIRACETAM INJ | | All non-preferred meds must be used in specified order. |
| MC | LAMICTAL XR | MC | 8 | LIBERVANT | 7. Adjunctive therapy in the treatment of partial-onset seizures in patient's ≥16 years of age with epilepsy. | Please use Drug-Drug Interaction PA form #10400 for this combination. |
| MC/DEL | LAMOTRIGINE ER ODT | MC/DEL | 8 | LYRICA CR | | |
| MC/DEL | LAMOTRIGINE IR ² | MC/DEL | 8 | LYRICA SOL ³ | | |
| MC/DEL | LEVETIRACETAM SOLN | MC | 8 | MOTPOLY XR | | |
| MC/DEL | LEVETIRACETAM TABS | MC/DEL | 8 | MYSOLINE TABS | | |
| MC/DEL | LEVETIRACETAM ER TABS | MC | 8 | ONFI | 8. Epidiolex is for the treatment of seizures associated with Lennox-Gastaut syndrome (LGS), Dravet syndrome (DS) or TS (Tuberous Sclerosis Complex) in patients 1 years of age and older. | Epidiolex Criteria for Lennox-Gastaut syndrome (LGS) and Dravet: a trial of two drugs (clobazam, levetiracetam, valproate derivatives, lamotrigine, topiramate, rufinamide, or felbamate). |
| MC/DEL | LYRICA ³ | MC/DEL | 8 | OXCARBAZEPINE SUS | | Diacomit is for the treatment of seizures associated with Dravet syndrome (DS) in patients 6 months of age and older and weighing 7kg or more There are no clinical data to support the use of Diacomit® as monotherapy in DS. |
| MC/DEL | NAYZILAM ¹ | MC | 8 | OXTELLAR XR ⁵ | | DDI: Concomitant use of Diacomit® with other CNS depressants, including alcohol, may increase the risk of sedation and somnolence. Concomitant use of strong inducers (CYP1A2, CYP3A4, or CYP2C19 inducers, such as rifampin, phenytoin, phenobarbital, and carbamazepine) should be avoided, or dosage adjustments should be made. |
| MC/DEL | OXCARBAZEPINE | MC/DEL | 8 | PHENYTEK CAPS | | |
| MC/DEL | PREGABALIN CAPS | MC/DEL | 8 | POTIGA | | |
| MC/DEL | PHENYTOIN | MC/DEL | 8 | PREGABALIN (ORAL) SOL | | |
| MC/DEL | PRIMIDONE TABS | MC | 8 | ROWEEPRA TAB | 9. For seizures associated with Dravet syndrome in patients 2 years of age and older | DDI: Avoid concomitant use of Nayzilam® with moderate or strong CYP3A inhibitors. |
| MC/DEL | QUDEXY XR | MC | 8 | SABRIL | | |
| MC/DEL | TEGRETOL SUS | MC | 8 | SEZABY | | |
| MC/DEL | TOPIRAMATE | MC | 8 | SPRITAM | | |
| MC/DEL | TOPIRAMATE SPRINKLE IR CAPS | MC | 8 | SYMPAZAN | | |
| MC/DEL | TRILEPTAL SUS | MC/DEL | 8 | TEGRETOL TAB | 10. Adjunctive therapy 12 and older. | Xcopri criteria: History of trials with at least 4 AEDs (2 generic, 2 branded or Uncontrolled seizures on three AEDs; or Uncontrolled on 2 AEDs given along with VNS. Uncontrolled defined as 3 or more TC seizures per year (increases risk of SUDEP); > 6 disabling seizures per year . Any patient who has gone to the ED 2 or more times in the prior 12 months (who has also tried and failed at least 3 other drugs). Ongoing use requires 50 percent reduction in seizure frequency after three months. |
| MC/DEL | VALPROIC ACID TABS | MC/DEL | 8 | TIAGABINE | | |
| MC/DEL | VALPROIC ACID SOL | MC | 8 | TOPAMAX | | |
| MC | VALTOCO ² | MC/DEL | 8 | TOPIRAMATE ER CAPS | | Motpoly XR: pediatric patient weight must be > 50kg and requires multiple preferred medication trials including generic lacosamide |
| MC/DEL | ZONISAMIDE | MC | 8 | TOPAMAX SPRINKLE ER CAPS ² | | |
| | | MC | 8 | TOPAMAX SPRINKLE IR CAPS ² | | |
| | | MC/DEL | 8 | TOPIRAMATE SPRINKLE ER CAPS ² | | |
| | | MC | 8 | TROKENDI ^{2,6} | | Libervant: For the acute treatment of intermittent, stereotypic episodes of frequent seizure activity (i.e., seizure clusters, acute repetitive seizures) that are distinct from a patient's usual seizure pattern in patients with epilepsy 2 to 5 years of age as long as all preferred therapies have been tried and failed at full therapeutic doses. |
| | | MC/DEL | 8 | VIMPAT ⁴ | 11. Initial monotherapy for the treatment of partial-onset or primary generalized tonic-clonic seizures in patients 2 years of age and older. | |
| | | MC/DEL | 8 | VIMPAT SOL ⁴ | Adjunctive therapy for the treatment of partial-onset seizures, primary generalized tonic-clonic seizures, and seizures associated with Lennox Gastaut syndrome in patients 2 years of age and older. The preventive treatment of migraine in patients 12 years and older. | |
| | | MC | 8 | XCOPRI | | |
| | | MC/DEL | 8 | ZARONTIN SYRP | | |
| | | MC/DEL | 8 | ZARONTIN CAP | | |
| | | MC/DEL | 8 | ZARONTIN SOL | | |
| | | MC | 8 | ZONISADE | | |
| | | MC | 8 | ZTALMY | | |
| | | MC/DEL | 9 | KEPPRA XR | | |
| | | MC/DEL | 9 | NEURONTIN | | |
| | | MC/DEL | 9 | TEGRETOL-XR TB12 | | |

| | | | | | | |
|--|--|--|--|--|---|--|
| | | | | | | <p>Will require a step though topiramate.</p> <p>SEE ANTICONVULSANT INDICATION CHART AT THE END OF THIS DOCUMENT</p> <p>M= Monotherapy A= Adjunctive 9= No Evidence</p> <p>The step orders show the relative strength of evidence for use in bi-polar and will guide prior authorization determinations.</p> <p>Step 4 drugs-no PA required.</p> |
| | | | | | <p>BIPOLAR DISORDER: STEP ORDER</p> <p>M ~ A</p> <p>4 ~ 4 LAMICTAL</p> <p>4 ~ 4 LITHIUM</p> <p>4 ~ 4 CARBAMAZEPINE</p> <p>4 ~ 4 VALPROATE</p> <p>4 ~ 4 ATYPICAL ANTIPSYCHOTICS EXC. CLOZAPINE</p> <p>5 ~ 5 TRILEPTAL</p> <p>9 ~ 6 TOPAMAX</p> <p>9 ~ 7 KEPBRA TABS</p> <p>9 ~ 8 GABITRIL TABS</p> <p>9 ~ 9 NEURONTIN</p> | |
| | | | | | <p>PEDIATRIC BIPOLAR1 DISORDER: STEP ORDER</p> <p>M ~ A</p> <p>(6-18 YEARS WITH OR WITHOUT PSYCHOSIS)</p> <p>4 ~ 4 LITHIUM</p> <p>4 ~ 4 CARBAMAZEPINE</p> <p>4 ~ 4 VALPROATE</p> <p>4 ~ 4</p> <p>ATYPICAL ANTIPSYCHOTICS EXC. CLOZAPINE</p> <p>4 ~ 4 LAMICTAL</p> <p>5 ~ 5 TRILEPTA</p> | <p>Two-step 1 preferred drugs must be tried before Trileptal.</p> <p>The step orders show the relative strength of evidence for use in bi-polar and will guide prior authorization determinations.</p> <p>Step 4 drugs-no PA required.</p> |

ANTI-PARKINSON DRUGS

| | | | | | | | | |
|--|------------------------|--|---|----------------------------------|------------------|--|---|---|
| PARKINSONS - ANTICHOLINERGICS | MC/DEL MC MC/DEL | | BENZTROPINE MESYLATE TABS COGENTIN SOLN TRIHEXYPHENIDYL | | | | Use PA Form# 20420 | |
| PARKINSONS - ADENOSINE RECEPTOR ANTAGONIST | | | | MC/DEL | | NOURIANZ | | <p>Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>DDI: Avoid use of Nourianz® with strong CYP3A4 inducers (e.g. carbamazepine, rifampin, phenytoin, St. John's wort).</p> <p>Use PA Form# 20420</p> |
| PARKINSONS - COMT INHIBITORS | | | | MC/DEL MC MC/DEL | | COMTAN TABS ONGENTYS TASMAR TABS | Use PA Form# 20420 | Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| PARKINSONS - SELECTED DOPAMIN AGONISTS | MC/DEL MC/DEL | | PRAMIPEXOLE ROPINIROLE | MC/DEL MC MC/DEL MC/DEL | 5 8 8 8 | MIRAPEX TABS ¹ REQUIP TABS MIRAPEX ER NEUPRO PATCH | Use PA Form# 20420 1. As of 12/08 users of Mirapex will be grandfathered if diagnosis is Parkinsons. | Preferred drug must be tried and failed in step-order due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| PARKINSONS- MAOIS | | | | MC | | XADAGO | | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered |

| | | | | | | | | |
|--|--|--|--|--|---|--|---|---|
| | | | | | | | on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. | |
| | | | | | | | Use PA Form# 20420 | |
| PARKINSONS - DOPAMINERGICS/CARBI/ LEVO | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL | | AMANTADINE HCLCAPS AMANTADINE HCL TABS BROMOCRIPTINE MESYLATE TABS BROMOCRIPTINE MESYLATE CAPS CARBIDOPA/LEVODOPA TABS ³ CARBIDOPA/LEVODOPA ER CARBIDOPA/LEVO/ENTACAPONE TAB LARODOPA TABS SELEGILINE CAPS HCL SELEGILINE TABS HCL | MC/DEL MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC/DEL MC MC MC MC | | APOKYN AZILECT ² CARBIDOPA/LEVODOPA RAPDIS ELDEPRYL CAPS GOCOVRI INBRIJA KYNMOBI LODOSYN TABS OSMOLEX ER PARLODEL CAPS PARLODEL TABS RYTARY SINEMET TABS SINEMET TBCR ZELAPAR ¹ | 1. Approvals will require concurrent therapy with Levodopa and failed trials of Selegiline, Comtan, and Stalevo. 2. Approvals will require trials of Carbidopa/Levodopa, Selegiline, Comtan, and Stalevo. 3. Only preferred manufacturer's products will be available without prior authorization. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Inbrija is recommended for the intermittent treatment of OFF episodes in patients with Parkinson's disease treated with carbidopa/levodopa. |
| PARKINSONS - COMBO. | | | | MC/DEL MC | | STALEVO ¹ CARBIDOPA/LEVODOPA/ENTACA ¹ | Use PA Form# 20420 1.Clinical PA is required to establish diagnosis and medical necessity. | |
| MUSCLE RELAXANTS | | | | | | | | |
| MUSCLE RELAXANTS | MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL | | BACLOFEN TABS CHLORZOXAZONE TABS CYCLOBENZAPRINE HCL 5mg & 10mg TABS LIORESAL INTRATHECAL KIT METHOCARBAMOL TABS TIZANIDINE HCL TABS | MC/DEL MC/DEL MC/DEL MC MC MC MC MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | 7 8 8 8 8 8 8 8 8 8 8 8 8 9 9 9 9 | ORPHENADRINE CITRATE CARISOPRODOL 350MG TABS AMRIX DANTRIUM CAPS FLEQSUVY LIORESAL TABS LORZONE LYVISPAH METAXALONE NORFLEX TBCR OZOBAX ROBAXIN-750 TABS VECUROMIUM INJ ZANAFLEX TABS CARISOPRODOL 250MG TABS CHLORZOXAZONE 250mg TABS SKELAXIN TAB SOMA TABS | At least 4 preferred drugs (including tizanidine) must be tried for at least 2 weeks and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an..... acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Elderly patients, over 65, will require written notice of the increased sedative risks and impaired driving.Prior Authorization will not be given for:1. frequent or persistent early refills of controlled drugs; 2. multiple instances of early refill overrides due to reports of misplacement, stolen, dropped in toilet or sink, distant travel, etc. Non-preferred drugs will not be approved if members circumventing MaineCare prior authorization requirements by paying (prescribers failed to submit prior authorization prior to cash narcotic scripts being filled by member). Non-preferred products must be used in specified step order. Lorzone is non preferred and requires at least 4 preferred drugs (including tizanidine) and step care therapy (orphenadrine), as well as reasons for why chlorzoxazone is not acceptable. | |
| MUSCLE RELAXANT - COMBO. | | | | MC/DEL MC/DEL MC MC/DEL MC/DEL MC | | CARISOPRODOL/ASPIRIN TABS CARISOPRODOL/ASPIRIN/CODE NORGESIC TABS ORPHENADRINE COMPOUND ORPHENADRINE/ASA/CAFF ORPHENGESIC | Use PA Form# 20420 | Individual components are available with PA described in the section above.1. frequent or persistent early refills of non-controlled drugs; 2. multiple instances of early refill overrides due to reports of misplacement stolen, dropped in toilet or sink, distant travel, etc. |
| PARATHYROID HORMONE | | | | | | | | |
| PARATHYROID HORMONE | | | | MC | | NATPARA ¹ | 1. Recommended only for those who cannot be well-controlled on calcium | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the |

| | | | | |
|--------|---|--------|---|---|
| MC | DAILY MULTI VIT/IRON | MC | FERRALET 90 | Click here for the OTC List |
| MC/DEL | DIALYVITE 1MG | MC | IBERET | |
| MC/DEL | DIALYVITE 800MG | MC | MATERNA TABS | |
| MC/DEL | FULL SPECTRUM B | MC | MAXARON | |
| MC | M.V.I.-12 INJ | MC | MULTIRET FOLIC -500 TBCR | |
| MC | MULTI-VIT/FLUORIDE | MC/DEL | NATAFORT TABS | |
| MC/DEL | NATALCARE RX TABS | MC/DEL | NATALCARE CFE 60 TABS ¹ | |
| MC/DEL | NEPHRONEX | MC/DEL | NATALCARE GLOSS TABS ¹ | |
| MC/DEL | NIVA-PLUS (ORAL) TABLET | MC | NATALCARE PIC TABS ¹ | |
| MC/DEL | ONE DAILY TABS | MC | NATALCARE PIC FORTE TABS ¹ | |
| MC/DEL | ONE-DAILY MULTIVITAMINS | MC/DEL | NATALCARE PLUS TABS ¹ | |
| MC/DEL | ONE-TABLET-DAILY | MC | NATALCARE THREE TABS ¹ | |
| MC/DEL | POLY-VIT/IRON/FLUORID SOLN | MC/DEL | NATACHEW CHEW | |
| MC/DEL | POLY-VITAMIN/FLUORIDE SOLN | MC | NATALFIRST TABS | |
| MC/DEL | POLY-VITAMINS/IRON SOLN | MC | NATATAB RX TABS | |
| MC | PRENATA (ORAL) TAB CHEW | MC/DEL | NEPHPLEX RX TABS | |
| MC/DEL | PRENATAL TABS ¹ | MC/DEL | NEPHROCAPS CAPS | |
| MC/DEL | PRENATAL FORMULA 3 TABS ¹ | MC/DEL | NEPHRO-VITE TABS | |
| MC/DEL | PRENATAL PLUS TABS ¹ | MC | NESTABS RX TABS | |
| MC/DEL | PRENATAL PLUS NF TABS ¹ | MC/DEL | NIFEREX | |
| MC | PRENATAL PLUS/27MG IRON ¹ | MC/DEL | OCUVITE TABS | |
| MC | PRENATAL PLUS/IRON TABS ¹ | MC | POLY-VI-FLOR SOLN | |
| MC | PRENATAL VITAMIN PLUS LOW IRON (ORAL) TAB | MC | POLY-VI-SOL SOLN | |
| MC/DEL | PRENATAL RX/BETA-CAROTENE ¹ | MC | POLY-VI-SOL/IRON SOLN | |
| MC/DEL | PREPLUS (ORAL) TABLET | MC | POLY-VITAMIN DROPS SOLN | |
| MC/DEL | RENAL CAPS | MC | PRECARE | |
| MC/DEL | RENAPHRO CAPS | MC | PREFERA OB | |
| MC | STRESS TAB NF TABS | MC | PREMESIS RX TABS | |
| MC | THERAPEUTIC-M TABS | MC | PRENATABS CBF TABS ¹ | |
| MC | THERAVITE LIQD | MC | PRENATAL CARE TABS ¹ | |
| MC/DEL | TRINATAL RX 1 (ORAL) TABLET | MC | PRENATAL MR 90 TBCR ¹ | |
| MC/DEL | TRIVEEN-DUO DHA (ORAL) COMBO. PKG | MC/DEL | PRENATAL MTR/SELENIUM TABS ¹ | |
| MC/DEL | TRI-VITAMIN/FLUORIDE SOLN | MC | PRENATAL OPTIMA ADVANCE TABS ¹ | |
| MC | VITA CON FORTE CAPS | MC | PRENATAL PC 40 TABS ¹ | |
| MC | VITAPLEX PLUS TABS | MC/DEL | PRENATAL RX TABS ¹ | |
| | | MC | PRENATE ¹ | |
| | | MC | PRENATE ELITE ¹ | |
| | | MC | PRIMACARE MISC | |
| | | MC | PROTEGRA CAPS | |
| | | MC | STUARTNATAL PLUS 3 TABS ¹ | |
| | | MC | TRI-VI-SOL SOLN | |
| | | MC | TRI-VI-SOL/IRON SOLN | |
| | | MC/DEL | ULTRA NATALCARE TABS | |
| | | MC | ULTRA-NATAL TABS ¹ | |
| | | MC | VICON FORTE CAPS | |
| | | MC | VINATAL FORTE TABS ¹ | |
| | | MC | VINATE ¹ | |
| | | MC/DEL | VINATE ADVANCED TABS ¹ | |

MISCELLANEOUS MINERALS

| | | | | | | |
|----------|--------|---------------------------|--------|------------------------------|---|---|
| MINERALS | MC | CALCARB | MC | ANEMAGEN | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. |
| | MC | CALCI-MIX CAPSULE CAPS | MC | CALCET TABS | Please refer to OTC list. | |
| | MC | CALCIQUID SYRP | MC/DEL | CALCIUM 600-D TABS | | |
| | MC | CALCITRATE/VITAMIN D TABS | MC | CALCIUM/VITAMIN D TABS | | |
| | MC/DEL | CALCIUM | MC | CALTRATE 600 PLUS/VIT D TABS | Click here for the OTC List | |
| | MC/DEL | CALCIUM CARBONATE | MC | CALTRATE PLUS TABS | | |
| | MC/DEL | CALCIUM CITRATE TABS | MC | CHROMAGEN | | |
| | | | | | | |

| | | | |
|--------|--------------------------------|--------|---------------------------------------|
| MC/DEL | CALCIUM GLUCONATE TABS | MC | CITRACAL PLUS TABS |
| MC/DEL | CALCIUM LACTATE TABS | MC | CONTRIN CAPS |
| MC | CALCIUM/MAGNESIUM TABS | MC | FEOGEN FORTE CAPS |
| MC/DEL | CALCIUM/VITAMIN D TABS | MC | FEROCON CAPS |
| MC | CALTRATE 600 TABS | MC/DEL | FERREX 150 CAPS |
| MC/DEL | CHEWABLE CALCIUM CHEW | MC | FERRO-SEQUELS TBCR |
| MC | CITRACAL TABS | MC | FE-TINIC CAPS |
| MC | CITRACAL + D TABS | MC | FE-TINIC 150 FORTE CAPS |
| MC | CITRUS CALCIUM TABS | MC/DEL | FLUOR-A-DAY SOLN |
| MC | CITRUS CALCIUM 1500 + D TABS | MC | HEMOCYTE TABS |
| MC | EFFERVESCENT POTASSIUM TBEF | MC/DEL | K-DUR TBCR |
| MC/DEL | FEOSTAT CHEW | MC | KLOR-CON PACK |
| MC | FERATAB TABS | MC | K-LYTE |
| MC/DEL | FER-GEN-SOL SOLN | MC/DEL | K-PHOS TABS NEUTRAL |
| MC | FER-IRON SOLN | MC | K-TABS TBCR |
| MC | FERRONATE TABS | MC | K-VESCENT PACK |
| MC/DEL | FERROUS SULFATE | MC | MICRO-K 10 MEG CPCR |
| MC/DEL | FLUOR-A-DAY CHEW | MC | NU-IRON 150 CAPS |
| MC | FLUORIDE CHEW | MC/DEL | OYSTER SHELL CALCIUM/VITA TABS |
| MC | FLUORIDE SODIUM CHEW | MC/DEL | POLY-IRON 150 CAPS |
| MC | FLUORITAB CHEW | MC/DEL | POLYSACCHARIDE IRON CAPS |
| MC | HM CALCIUM TABS | MC/DEL | POTASSIUM BICARB/CHLORIDE |
| MC | K+ POTASSIUM PACK | MC/DEL | POTASSIUM CHLORIDE 10MEQ CAPS |
| MC | KAON ELIX | MC/DEL | POTASSIUM CHLORIDE 8MEQ CAPS |
| MC | KAON-CL-10 TBCR | MC | TUMS 500 CHEW |
| MC | KCL 0.075%/D5W/NACL 0.2% SOLN | MC | VIACTIV CHEW |
| MC | K-EFFERVESCENT TBEF | | |
| MC | KLOR-CON | | |
| MC | KLOTRIX TBCR | | |
| MC/DEL | K-PHOS TABS | | |
| MC/DEL | K-VESCENT TBEF | | |
| MC/DEL | LURIDE CHEW | | |
| MC/DEL | MAGNESIUM GLUCONATE TABS | | |
| MC/DEL | MAGNESIUM SULFATE SOLN | | |
| MC | MAGTABS | | |
| MC | MICRO-K 8 MEG | | |
| MC/DEL | OS-CAL TABS | | |
| MC/DEL | OS-CAL 500 + D TABS | | |
| MC/DEL | OYSCO | | |
| MC/DEL | OYST-CAL TABS | | |
| MC/DEL | OYST-CAL D TABS | | |
| MC/DEL | OYST-CAL/VITAMIN D TABS | | |
| MC/DEL | OYSTER CALCIUM TABS | | |
| MC/DEL | OYSTER SHELL | | |
| MC | PHARMA FLUR | | |
| MC/DEL | PHOSPHA 250 NEUTRAL TABS | | |
| MC | POTASSIUM BICARBONATE TBEF | | |
| MC/DEL | POTASSIUM CHLORIDE 8MEQ | | |
| MC | POTASSIUM EFFERVESCENT | | |
| MC/DEL | SELENIUM TABS | | |
| MC | SLOW-MAG TBCR | | |
| MC/DEL | SODIUM FLUORIDE | | |
| MC | V-R CALCIUM | | |
| MC | V-R OYSTER SHELL CALCIUM | | |
| MC | ZINC SULFATE CAPS | | |

Please refer to OTC list.

Preferred products that used to require diag codes still require diag codes unless indicated otherwise.

| | | | | | | | | |
|---|--------|--|--------------------------------|--------|---|-----------------------------------|---|---|
| | | | | MC | 8 | STIMUFEND | | |
| | | | | MC/DEL | 8 | ZARXIO | | |
| | | | | MC | 9 | NEULASTA ¹ | Use PA Form# 20520 | |
| GAUCHER DISEASE | | | | | | | | |
| GAUCHER DISEASE | | | | MC | | CERDELGA ¹ | 1. Clinical PA for indication required. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Exceeding days supply limits for LMWH class requires PA. |
| | | | | MC | | YARGESA ¹ | | Yargesa: As monotherapy for the treatment of adult patients with mild to moderate type 1 Gaucher disease for whom enzyme replacement therapy is not a therapeutic option (e.g., due to allergy, hypersensitivity, or poor venous access). |
| | | | | | | | Use PA Form# 20420 | |
| ANTICOAGULANTS / PLATELET AGENTS | | | | | | | | |
| ANTICOAGULANTS | MC | | COUMADIN TABS | MC | | ARIXTRA SOLN | 1. Enoxaparin therapy durations greater than 7 days every 30 days require PA | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Exceeding days supply limits for LMWH class requires PA. |
| | MC/DEL | | ENOXAPARIN ¹ | MC/DEL | | FONDAPARINUX | 2. Use other strengths available to obtain desired dose. | |
| | MC | | ELIQUIS | MC/DEL | | FRAGMIN INJ | 3. Diagnosis required | DDI: Warfarin will require prior authorization if being used in combination with fluconazole, miconazole, or voriconazole. |
| | MC | | ELIQUIS STARTER PACK | MC/DEL | | FRAGMIN VIAL | 4. For the treatment of patients aged 3 months to less than 12 years of age. | DDI: Warfarin will require prior authorization if being used in conjunction with Gemfibrozil or Fenofibrate. |
| | MC | | HEPARIN SODIUM/NACL 0.9% SOLN | MC/DEL | | LOVENOX SOLN | | DDI: Rifampin will require prior authorization if being used in combination with Savaysa |
| | MC | | HEP-LOCK SOLN | MC/DEL | | LOVENOX 300 ² | | |
| | MC | | INNOHEP | MC/DEL | | LOVENOX SUBQ SYRINGE | | |
| | MC | | HEPARIN LOCK SOLN | MC/DEL | | PRADAXA ORAL PELLETS ⁴ | | |
| | MC/DEL | | HEPARIN LOCK FLUSH SOLN | MC | | IPRIVASK | | |
| | MC/DEL | | HEPARIN SODIUM SOLN | MC/DEL | | SAVAYSAS ³ | | |
| | MC/DEL | | HEPARIN SODIUM LOCK FLUSH SOLN | | | | | |
| | MC/DEL | | PRADAXA | | | | | |
| | MC/DEL | | JANTOVEN | | | | | |
| | MC/DEL | | WARFARIN SODIUM TABS | | | | | |
| | MC/DEL | | XARELTO | | | | | |
| | MC/DEL | | XARELTO STARTER PACK | | | | | |
| | | | | | | | Use PA form# 20420 | |
| ANTIHEMOPHILIC AGENTS | MC/DEL | | AFSTYLA | MC/DEL | | ADYNOVATE VIAL | 1. Only if other products unavailable. | Non-preferred will only be approved if other preferred products are unavailable. |
| | MC | | ALPHANATE | MC | | ADVATE ^{1,2,5} | | |
| | MC | | ALPHANINE SD | MC | | ALTUVIIIIO ⁴ | | Beqvez: FDA Approved Indication: An adeno-associated virus vector-based gene therapy indicated for the treatment of adults with moderate to severe hemophilia B (congenital factor IX deficiency) who: |
| | MC/DEL | | ALPROLIX VIAL | MC/DEL | | BEQVEZ | 2. Advate may be available with PA in cases of large volume dosing in patients with poor venous access. | · Currently use factor IX prophylaxis therapy, or |
| | MC/DEL | | BEBULIN VIAL | MC/DEL | | ESPEROCT | | · Have current or historical life-threatening hemorrhage, or |
| | MC/DEL | | BENEFIX SOLR | MC/DEL | | ELOCTATE | | · Have repeated, serious spontaneous bleeding episodes, and, |
| | MC/DEL | | HELIXATE FS KIT | MC/DEL | | HEMGENIX | | · Do not have neutralizing antibodies to adeno-associated virus serotype Rh74var (AAVRh74var) capsid as detected by an FDA- approved test. |
| | MC | | HEMLIBRA | MC/DEL | | IDELVION | | |
| | MC | | HEMOFIL - M | MC/DEL | | KOGENATE FS ⁵ | | Hemgenix® is an adeno-associated viral vector-based gene therapy for IV infusion after dilution. For treatment of adults with Hemophilia B (congenital Factor IX deficiency) who: Currently use Factor IX prophylaxis therapy, or have current or historical life-threatening hemorrhage, or Have repeated, serious spontaneous bleeding episodes. |
| | MC | | HUMATE-P SOLR | MC/DEL | | REBINYN | 3. Not indicated for use in children <12 years of age due to greater risk for hypersensitivity reactions and is not indicated for use in previously untreated patients. | |
| | MC/DEL | | IXINITY VIAL | MC | | RECOMBINATE VIAL ⁵ | | Altuviiiio is a von Willebrand Factor (VWF) independent recombinant DNA-derived, Factor VIII concentrate indicated for use in adults and children with hemophilia A (congenital factor VIII deficiency) for: Routine prophylaxis to reduce the frequency of bleeding episodes, On-demand treatment and control of bleeding episodes, Perioperative management of bleeding. |
| | MC/DEL | | JIVI ³ | MC | | ROCTAVIAN ⁴ | | |
| | MC | | KOATE-DVI | MC | | SEVENFACT | | |
| | MC | | KONYNE - 80 | | | | | |
| | MC/DEL | | KOVALTRY | | | | | Roctavian: For the treatment of adults with severe hemophilia A (congenital factor VIII deficiency with factor VIII activity <1 IU/dL) without antibodies to adeno-associated virus serotype |
| | MC | | MONARC - M | | | | | Inclusion: |
| | MC | | MONOCLATE - P | | | | | Severe factor VIII deficiency (less than 1% native factor VIII). |
| | MC | | MONONINE | | | | | Exclusion Criteria: |
| | MC/DEL | | NOVOEIGHT | | | | | Antibodies to the virus AAV5 |
| | MC | | NOVOSEVEN SOLR | | | | | Factor VIII inhibitors (or history of) |
| | MC | | NUWIQ | | | | | Known significant fibrosis of cirrhosis of the liver, or unexplained elevated LFTs |
| | MC/DEL | | PROFILNINE | | | | | History of inadequate compliance with prophylaxis, or regular bleeds despite adequate prophylaxis |
| | MC | | RECOMBINATE SOLR | | | | | Conditions in which high-dose steroids are contraindicated. |
| | MC | | REFACTO | | | | | |

| | | | | | | | | | |
|---|--|---|--|--|----------------------------|---|--|--|---|
| | MC/DEL MC MC/DEL | | RIXUBIS VIAL WILATE INJ XYNTHA | | | | | | -Inability to abstain from alcohol for one year Plan to impregnate a partner within 6 months of infusion -Hypersensitivity to mannitol -Active infections, either acute or uncontrolled chronic -HIV infection (limited information on use in this population) Use PA Form# 20420 |
| PLATELET AGGREGATION INHIBITORS | MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL | | ASPIRIN ASPIRIN-DIPYRIDAMOLE ER CPMP 12HR BRILINTA ¹ DIPYRIDAMOLE TABS CLOPIDOGREL 75MG PRASUGREL HCL TAB | MC/DEL MC MC MC/DEL MC/DEL MC/DEL | 7 8 8 8 8 8 | TICLOPIDINE HCL TABS DURLAZA EFFIENT PERSANTINE TABS PLAVIX TABS ZONTIVITY | | Use PA Form# 20715 for Plavix, Effent & Brilinta Use PA form# 20420 for other requests 1. Dosing limits apply, please see dose consolidation list. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. A special PA may be obtained at the pharmacy for members scheduled for "stent" placement or have had placement if in the last 12months. Please indicate on prescription date of stent placement. DDI: Plavix will require prior authorization if being used in combination with omeprazole, esomeprazole, cimetidine, fluconazole, ketoconazole, intelence, fluoxetine, ticlopidine, and fluvoxamine. DDI: exists for using maintenance ASA dose >100mg, as it reduces the effectiveness of Brilinta Brilinta- Concomitant use with strong CYP3A4 inhibitors should be avoided (including ketoconazole, itraconazole, atazanavir, and telithromycin). Doses of simvastatin and lovastatin >40mg should be avoided. |
| PLATELET AGGR. INHIBITORS / COMBO'S - MISC. | MC/DEL MC/DEL | | CILOSTAZOL PENTOXIFYLLINE ER TBCR | MC/DEL MC/DEL MC/DEL MC MC | | AGRYLIN CAPS ANAGRELIDE CAPS PLETAL TABS TRENAL TBCR YOSPRALA | | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| HEMATOLOGICALS | | | | | | | | | |
| MONOCLONAL ANTIBODY | | | | MC/DEL MC MC/DEL MC MC/DEL MC MC | | EMPAVELI ENSPRYNG FABHALTA GAMIFANT SOLIRIS ULTOMIRIS UPLIZNA VOYDEYA | | Use PA Form# 20420 | A diagnosis of Paroxysmal nocturnal hemoglobinuria (PNH) using the HAM test or flow cytometry is required. In addition, the patient must show evidence of having received a meningitis vaccine at least 2 weeks prior to the start of therapy. Gamifant is recommended for the treatment of adult and pediatric (newborn and older) patients with primary hemophagocytic lymphohistiocytosis (HLH) with refractory, recurrent, or progressive disease or intolerance with conventional HLH therapy. Fabhalta and Ultomiris are recommended for the treatment of adults with paroxysmal nocturnal hemoglobinuria (PNH). |
| IMMUNE GLOBULIN | MC MC/DEL MC MC MC/DEL MC/DEL MC | | BIVIGAM ¹ CUTAQUIG ¹ GAMUNEX-C GAMMAGARD S-D ¹ HIZENTRA ¹ PANZYGA ¹ PRIVIGEN ¹ | MC MC/DEL MC MC/DEL MC MC/DEL | | ASCENIV ² CUVITRU GAMMAPLEX INJ HYQVIA OCTAGAM INJ ¹ XEMBIFY | | Use PA Form# 20420 1. Clinical PA required 2. For the treatment of patients between 12 to 17 years of age. | Cutaquig is indicated as replacement therapy for primary humoral immunodeficiency (PI) in adults. Xembify is indicated for treatment of primary humoral immunodeficiency (PI) in patients 2 years of age and older. Asceniv indicated for the treatment of primary humoral immunodeficiency (PI) in adults and adolescents (12 to 17 years of age). PI includes but is not limited to the humoral immune defect in congenital agammaglobulinemia, common variable immunodeficiency (CVID), X-linked agammaglobulinemia, Wiskott-Aldrich syndrome, and severe combined immunodeficiencies (SCID). |
| HEREDITARY ANGIOEDEMA | MC MC MC MC/DEL | PROPHYLAXIS | | | | PROPHYHLAXIS | | 1. Clinical PA is required to establish diagnosis and medical necessity. 2. For the treatment of patients ≥ 12 years of age. | Haegarda is indicated for routine prophylaxis to prevent Hereditary Angioedema (HAE) attacks in adolescent and adult patients |
| | | CINRYZE ¹ HAEGARDA ¹ ORLADEYO ^{1,2} TAKHZYRO ¹ | | | | | | | |
| | | TREATMENT | | | | TREATMENT | | | |
| | | BERINERT KIT ¹ FIRAZYR ¹ RUCONEST VIAL ¹ | MC/DEL | | KALBITOR VIAL | | | | |
| | | | | | | | | Use PA Form# 20420 | |

| | | | | | | | |
|--|--|--|---|--|---|--|---|
| HEMATOLOGICAL AGENTS- THROMBOPOIETIN RECEPTOR AGONISTS | MC MC | | PROMACTA ¹ NPLATE ¹ | MC MC/DEL MC/DEL | ALVAIZ DOPTELET MULPLETA | Use PA Form# 20420 1. Clinical PA required. Must see prior trial with insufficient response to corticosteroids and immunoglobulins. | Doptelet and Mulpelta: For the treatment of thrombocytopenia in adults with chronic liver disease who are scheduled to undergo a procedure. |
| HEMATOLOGICAL AGENTS-IgAN | | | | MC/DEL MC | FILSPARI ¹ TARPEYO | Use PA Form# 20420 1. PA required to confirm FDA approved indication. | All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists |
| ANEMIA- BETA THALASSEMIA | | | | MC MC | REBLOZYL ZYNTEGLO | Use PA Form# 20420 | Reblozyl is indicated for the the treatment of anemia in adult patients with beta thalassemia who require regular red blood cell (RBC) transfusion. It is not indicated for use as a substitute for RBC transfusions in patients who require immediate correction of anemia. Zynteglo is indicated for the treatment of adult and pediatric patients with β-thalassemia who require regular red blood cell (RBC) transfusions. |
| HEMATOLOGIC DISORDER TREATMENT AGENTS | | | | MC/DEL MC | CABLIVI TAVALISSE | Use PA Form# 20420 | Tavalisse is recommended for patients at risk of bleeding when one line of therapy (steroids, IVIG, splenectomy) has failed. Cablivi is recommended for the treatment of adult patients with acquired thrombotic thrombocytopenic purpura (aTTP), in combination with plasma exchange and immunosuppressive therapy. |
| COMPLEMENT RECEPTOR ANTAGONIST | | | | MC | TAVNEOS | Use PA Form# 20420 | |
| WHIM SYNDROME AGENTS | | | | MC | XOLREMDI | Use PA Form#20420 | Xolremdi: In patients 12 years of age and older with WHIM syndrome (warts, hypogammaglobulinemia, infections, and myelokathexis) to increase the number of circulating mature neutrophils and lymphocytes. |
| HEMOSTATIC | | | | | | | |
| HEMOSTATIC | MC/DEL MC | | AMICAR AMINOCAPROIC ACID | MC MC | FIBRYGA RIASTAP | Use PA Form# 20420 | Fibryga and Riastap are indicated for the treatment of acute bleeding episodes in adults and adolescents with congenital fibrinogen deficiency, including afibrinogenemia and hypofibrinogenemia. Fibryga® is not indicated for dysfibrinogenemia. |
| ACUTE HEPATIC PORPHYRIA (AHP) | | | | | | | |
| ACUTE HEPATIC PORPHYRIA (AHP) | | | | MC | GIVLAARI | Use PA Form# 20420 | Givlaari is indicated for the treatment of adults with acute hepatic porphyria (AHP). |
| PYRUVATE KINASE DEFICIENCY AGENTS | | | | | | | |
| PYRUVATE KINASE DEFICIENCY AGENTS | | | | MC | PYRUKYND ¹ | Use PA Form# 20420 1.PA required to confirm FDA approved indication. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) |
| OP. - ANTIBIOTICS | | | | | | | |
| | MC MC MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL | | AK-SPORE OINT BACITRACIN/NEOMYCIN/POLYM BACITRACIN/POLYMYXIN B OINT CHLOROPTIC SOLN ERYTHROMYCIN OINT NEOSPORIN SOLN POLYSPORIN TRIMETHOPRIM SULFATE/POLY TOBRAMYCIN SULFATE SOLN | MC MC MC MC MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC MC MC | AK-POLY-BAC OINT AK-SULF OINT AK-TOB SOLN AZASITE BACITRACIN OINT BLEPH-10 SOLN GATIFLOXACIN DROPS GENTAMICIN SULFATE GENTAK ILOTYCIN OINT LEVOFLOXACIN DROPS NEOMYCIN/BACI/POLYM OINT NEOMYCIN/POLYMYXIN/GRAMIC NEOSPORIN OINT OCUSULF-10 SOLN OCUTRICIN SOLN | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |

| | | | | | | | |
|--|--|--|--|--|--|---|--|
| | | | | MC/DEL MC/DEL MC/DEL MC | POLYTRIM DROPS SULFACETAMIDE SODIUM DROPS SULFACETAMIDE SODIUM OINT TERAK OINT | | |
| OP. - ANTI-PARASITIC | | | | MC | XDEMZY ¹ | Use PA Form# 20420 1. For the treatment of Demodex biopharitis. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| OP. - RHO KINASE INHIBITORS | MC | | RHOPRESSA | | | | on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) |
| | | | | | | Use PA Form# 20420 | |
| OP. - QUINOLONES | MC/DEL MC/DEL MC/DEL MC/DEL | | CILOXAN OINT CIPROFLOXACIN SOL 0.3% OFLOXACIN QUIXIN SOLN | MC/DEL MC/DEL MC | BESIVANCE CILOXAN SOLN OCUFLOX SOLN | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| OP. - QUINOLONES-4TH GENERATION | MC/DEL | | MOXIFLOXACIN 0.5% SOLN (Generic Vigamox) | MC | ZYMAXID | Use PA Form# 20420 | |
| OP. - ARTIFICIAL TEARS AND LUBRICANTS | MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC MC MC MC MC | | ARTIFICIAL TEARS OINT ARTIFICIAL TEARS SOLN CELLUVISC SOLN EYE LUBRICANT OINT GENTEAL LIQUITEARS SOLN MAJOR TEARS SOLN PURALUBE OINT PURALUBE TEARS SOLN REFRESH SOLN OP REFRESH PLUS SOLN ¹ REFRESH PM OINT | MC/DEL MC MC MC MC/DEL MC MC MC MC MC/DEL MC MC MC MC/DEL MC/DEL MC MC/DEL MC | ARTIFICIAL TEARS SOLN OP BION TEARS SOLN DRY EYES OINT DURATEARS OINT HYPO TEARS ISOPTO TEARS SOLN LACRI-LUBE LUBRIFRESH P.M. OINT MURINE SOLN MUROCEL SOLN NATURE'S TEARS SOLN REFRESH SOLN REFRESH TEARS SOLN ¹ TEARGEN SOLN TEARISOL SOLN TEARS NATURALE TEARS PURE SOLN TEARS RENEWED OINT THERATEARS SOLN V-R ARTIFICIAL TEARS SOLN | Use PA Form# 20420 1. Dosing limits apply, please see dose consolidation list. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| OP. - BETA - BLOCKERS | MC/DEL MC/DEL MC/DEL MC/DEL | | BETOPTIC-S SUSP CARTEOLOL HCL SOLN LEVOBUNOLOL HCL SOLN METIPRANOLOL SOLN | MC MC/DEL MC MC/DEL MC MC/DEL MC/DEL | BETAGAN SOLN BETAXOLOL HCL SOLN ISTALOL OCUPRESS SOLN OPTIPRANOLOL SOLN TIMOPTIC SOLN TIMOLOL DROP TIMOLOL SOL-GEL TIMOPTIC-XE SOLG | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| OP. - ANTI-INFLAMMATORY / STEROIDS OPTH. | MC MC/DEL | | AK-SPORE HC OINT ALREX SUSP | MC MC | AK-TROL SUSP BAC/POLY/NEOMY/HC OINT | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |

| | | | | | | | |
|---|--|--|--|--|---|---|--|
| | MC/DEL MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL | DEXAMETH SOD PHOS SOLN FLAREX SUSP FLUOROMETHOLONE SUSP FML DROPS SUSP 1% FML FORTE SUSP FML S.O.P. OINT LOTEMAX OINT LOTEMAX SUSP LOTEMAX SM DROPS GEL 0.38% NEO/POLY/DEXAMETH OINT NEO/POLY/DEXAMETH SUSP PRED-G SUSP PRED FORTE SUSP 1% PRED MILD SUSP PREDNISOLONE TOBRADEX OINT TOBRADEX SUSP TOBEX OINT SULFACETAMIDE/PREDNISOLONE ZYLET SUSP | MC MC MC MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC | BLEPHAMIDE S.O.P. OINT BLEPHAMIDE SUSP BROMDAY EFLONE SUSP FLUOR-OP SUSP ILUVIEN IMPLANT INVELTYS LOTEMAX GEL MAXITROL OPTH OINT 0.1% NEO/POLY/BAC/HC OINT NEOM/POLY/DEX OPTH OINT 0.1% OMNIPRED DROPS SUSP OZURDEX PRED-G S.O.P. OINT PREDNISOLONE SODIUM PHOSPHATE SOL RETISERT IMPLANT SULFACET SOD/PRED SOLN TRIESENCE VIAL TOBRADEX ST TOBRAMYCIN SUSP DEXAMETHASONE VASOCIDIN SOLN VEXOL SUSP XIPERE | | preferred drug(s) exists. | |
| OP. - PROSTAGLANDINS | MC/DEL MC MC/DEL MC/DEL | LATANOPROST SOL 0.005% LUMIGAN SOLN ROCKLATAN TRAVATAN-Z | MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | 7 8 8 8 8 8 8 8 8 8 | ZIOPTAN BIMATOPROST 0.03% DROPS DURYSTA IYUZEH RESCULA ^{1,2,3} TRAVATAN SOLN TRAVOPROST VYZULTA XALATAN SOLN ¹ XELPROS | 1. All preferreds must be tried. 2. Dosing limits apply, please see dosing consolidation list. 3. Clinical PA is required to establish diagnosis and medical necessity. Use PA Form# 20420 | Preferred drugs must be tried and failed, in step-order, due to lack of efficacy (failure to reach target IOP reduction) or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| OP. - CYCLOPLEGICS | MC MC/DEL MC/DEL MC/DEL | AK-PENTOLATE SOLN ATROPINE SULFATE CYCLOPENTOLATE HCL SOLN ISOPTO HYOSCINE SOLN | MC/DEL MC MC/DEL MC | | CYCLOGYL SOLN ISOPTO ATROPINE SOLN ISOPTO HOMATROPINE SOLN MUROCOLL-2 SOLN | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| OP. - MIOTICS - DIRECT ACTING | MC/DEL MC MC MC/DEL MC/DEL | ISOPTO CARBACHOL SOLN ISOPTO CARPINE SOLN PILOCAR SOLN PILOCARPINE HCL SOLN PILOPINE HS GEL | | | | Use PA Form# 20420 | |
| OP. - SELECTIVE ALPHA ADRENERGIC AGONISTS | MC MC MC MC/DEL MC/DEL | ALPHAGAN SOLN ALPHAGAN P 0.1% SOLN ALPHAGAN P 0.15% SOLN BRIMONIDINE DROPS 0.2 % SIMBRINZA | MC/DEL MC/DEL | | BRIMONIDINE TARTRATE DROPS 0.15 % IOPIDINE SOLN | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| OP. - ANTI-ALLERGICS | MC/DEL MC MC/DEL MC/DEL MC | AZELASTINE HCL DROPS BEPREVE CROMOLYN SODIUM DROPS KETOTIFEN FUMARATE DROPS LASTACFT | MC MC/DEL MC/DEL MC MC/DEL | 8 8 8 8 8 | ALOCRIOL SOLN ALOMIDE SOLN EMADINE SOLN OPTICROM SOLN PATANOL SOLN | Use PA Form# 20420 | All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |

| | | | | | | | | |
|---------------------------|--|--|---|--|--|--|---|---|
| TOPICAL - ANTIVIRALS | | | | MC/DEL MC/DEL MC MC | | ACYCLOVIR OINT DENA VIR CREA ^{1,3} YCANTH ZOVIRAX OINT ^{1,2} | 1. Must fail oral treatment with Acyclovir or Valacyclovir. 2. Approvals limited to 1 tube per 180 days. 3. Dosing limits apply, please see dosing consolidation list. 4. For the topical treatment of molluscum contagiosum in adult and pediatric patients 2 years of age and older. | |
| TOPICAL - ANTINEOPLASTICS | MC | | EFUDEX | MC/DEL MC/DEL MC MC/DEL | | CARAC CREA FLUOROURACIL SOLARAZE GEL ZYCLARA | | Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| TOPICAL - BURN PRODUCTS | MC MC/DEL MC MC MC/DEL | | FURACIN CREA SILVER SULFADIAZINE CREA SSD AF CREA SSD CREA THERMAZENE CREA | MC/DEL | | SILVADENE CREA | | Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| TOPICAL - CORTICOSTEROIDS | MC MC/DEL MC MC MC MC MC MC MC MC MC/DEL MC MC MC MC MC MC MC/DEL | | LOW POTENCY DERMA-SMOOTH- FS BODY HYDROCORTISONE CREA HYDROCORTISONE LOTN HYDROCORTISONE LOTN TEXACORT SOLN MEDIUM POTENCY DESOXIMETASONE 0.05% CREA/GEL FLUTICASONE PROPIONATE CREA/OINT HYDROCORTISONE BUTYRATE HYDROCORTISONE OINT HYDROCORTISONE VALERATE MOMETASONE FUROATE OINT TRIAMCINOLONE ACETONIDE .025-.1% HIGH POTENCY DESONIDE ¹ | MC/DEL MC MC MC/DEL MC/DEL MC MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC/DEL | | LOW POTENCY ACLOVATE ANUSOL HC-1 OINT DESONATE GEL FLUOCINOLONE ACETONIDE FLUOCINOLONE HALOG HYDROCORTISONE POWD LIDA MANTLE HC CREA PROCTOCORT CREA VERDESO MEDIUM POTENCY BESER LOTION³ CLODERM CREA CORDRAN CUTIVATE CREA / OINT CUTIVATE LOTN DERMATOP ELOCON OINT KENALOG AERS LOCOID LUXIQ FOAM | 1. Dosing limits apply, please see dosing consolidation list. 2. Treatment beyond 4 weeks is not recommended. 3. For the treatment of patients ≥ 12 years of age. 4. For the treatment of patients ≥ 18 years of age. | At least 1 drug from each potency of preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |

| | MC/DEL | | LIDOCAINE PTCH 5% | | | CONSOLIDATION LIST | |
|--|--|--|---|--|--------------------------------------|---|---|
| TOPICAL - DEPIGMENTING AGENTS | | | | MC MC MC MC/DEL MC/DEL MC MC MC | 8 8 8 8 8 8 8 9 | ALUSTRA CREA EPIQUIN MICRO GLYQUIN CREA HYDROQUINONE CREA HYDROQUINONE/SUNSCREENS SOLAQUIN FORTE CREA TRI-LUMA CREA ELDOQUIN | Use PA Form# 20420 Use PA Form# 20420 As per Medicaid Policy, cosmetic drugs are not covered. Non-cosmetic clinical applications will be considered by prior authorization on a case by case basis. |
| TOPICAL - SCABICIDES AND PEDICULICIDES | MC/DEL MC MC/DEL MC/DEL MC | | ACTICIN CREA LICE KILLING SHAM LICE TREATMENT CREME RINS LIQD PERMETHRIN LOTN NATROBA ¹ | MC MC MC/DEL MC MC MC/DEL | | ELIMITE CREA EURAX LINDANE MALATHION OVIDE LOTN SPINOSAD SUSP | Use PA Form# 20420 1. Dosing limits apply, please refer to dosage consolidation list. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| TOPICAL - WOUND / DECUBITUS CARE | | | | MC MC MC | | FILSUVEZ REGRANEX GEL VYJUVEK | Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Regranex will be approved for diabetic patients in good control (HgbA1c <8), who are not smoking, with a stage III or IV WOCN AND NPUAP lower extremity diabetic ulcer and with an adequate blood supply (TcP 02 >30, ABI>0.7 or ASP> 70), and where the underlying cause has been corrected. The wound must be free of infection and have been previously treated with preferred standard therapies for at least 2 months. Maximum approval for 20 weeks. Vyjuvek: For the treatment of wounds in patients 6 months of age and older with dystrophic epidermolysis bullosa (DEB) with mutation(s) in the collagen type VII alpha 1 chain (COL7A1) gene. Filsuvez: The patient has a diagnosis of dystrophic or junctional epidermolysis bullosa. The patient is at least 6 months old and does not have current evidence or history of squamous cell carcinoma or active infection in the area requiring Filsuvez application. The patient has used standard wound care treatments, including silicone or foam dressings without wound resolution Accuzyme and Ethezyme products have been removed from the PDL due to FDA concerns regarding drugs containing Papain. |
| TOPICAL - ASTRINGENTS / PROTECTANTS | MC | | XERAC AC SOLN | MC MC MC MC | | LOWILA BAR MOISTURIN DRY SKIN CREA PROSHIELD PLUS SKIN PROTE CREA SURGILUBE GEL | Use PA Form# 20420 1. Dosing limits apply, please refer to dosage consolidation list. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| TOPICAL - ANTISEPTICS / DISINFECTANTS | MC/DEL | | POVIDONE-IODINE SOLN | MC MC MC MC | | BETADINE OINT FORMALYDE-10 AERS IODOSORB LAZERFORMALYDE SOLUTION SOLN | Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| MISCELLANEOUS EYE | | | | | | | |
| OP. - EYE | MC MC MC MC MC MC/DEL | | AK-DILATE SOLN EYE WASH SOLN NAPHAZOLINE HCL SOLN PHENYLEPHRINE HCL SOLN PONTOCAINE SOLN SODIUM CHLORIDE | MC MC/DEL MC | | LENS PLUS REWETTING DROPS MURO 128 NEO-SYNEPHRINE SOLN | Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| MISCELLANEOUS EAR | | | | | | | |
| EAR | MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL | | A/B OTIC SOLN ACETASOL SOLN ACETASOL HC SOLN ACETIC ACID ACETIC ACID/HYDROCORTISON ALLERGEN SOLN CARBAMIDE PEROXIDE 6.5% OTIC SOLN. CIPRO HC SUSP CORTISPORIN-TC SUSP | MC MC MC/DEL MC/DEL MC/DEL MC MC MC MC | | ANTIBIOTIC EAR SOLN ANTIBIOTIC EAR SUSP CIPRODEX CIPROFLOXACIN HCL DEBROX SOLN FLOXIN FLUOCINOLONE ACETONIDE OIL DROPS 0.01% OTIPRIO OTOVEL | Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |

| | | | | | | | | |
|-------------------------------------|--|--|---|--|--|---|---|---|
| | MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL | | CORTOMYCIN COLY-MYCIN-S SUSP DERMOTIC EAR DROPS SOLN EAR DROPS RX SOLN EAR WAX REMOVAL DROPS NEOMYCIN/POLYMYXIN/HC OFLOXACIN 0.3% OTIC | | | | | |
| MOUTH ANTISEPTICS | | | | | | | | |
| MOUTH ANTI-INFECTIVES | MC MC/DEL | | NILSTAT SUSP NYSTATIN SUSP | MC MC | | MYCELEX TROC ORAVIG | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| MOUTH ANTISEPTICS | MC/DEL MC/DEL MC MC | | CHLORHEXIDINE GLUCONATE LIDOCAINE VISCOUS SOLN TRIAMCINOLONE IN ORABASE PSTE TRIAMCINOLONE ORADENT PSTE | MC MC MC MC | | APHTHASOL PSTE ¹ PERIOGARD SOLN ¹ TRIAMCINOLONE ACETONIDE PSTE ¹ | Use PA Form# 20420 1. Must fail all preferred products before non-preferred. | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| DENTAL PRODUCTS | | | | | | | | |
| DENTAL PRODUCTS | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC | | ETHEDENT CREA GEL-KAM CONC GEL-KAM GEL 0.4% PHOS FLUR SOLN SF 5000 PLUS CREA SF GEL STANNOUS FLUORIDE ORAL RI CONC | MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC | | APF GEL GEL DENTAGEL GEL PHOS-FLUR GEL THERA-FLUR-N GEL | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| ARTIFICIAL SALIVA/STIMULANTS | | | | | | | | |
| ARTIFICIAL SALIVA/STIMULANTS | MC | | SALIVA SUBSTITUTE SOLN | MC MC MC | | EVOXAC CAPS RADIACARE SOLR SALAGEN TABS | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| MISCELLANEOUS ANORECTAL | | | | | | | | |
| ANORECTAL - MISC. | MC MC MC/DEL MC/DEL MC/DEL | | CORTENEMA ENEM ELA-MAX 5 CREA HYDROCORTISONE ENEM PROCTOSOL HC CREA PROCTOZONE-HC CREA | MC/DEL MC/DEL MC/DEL MC/DEL MC | | ANUSOL-HC CREA CORTIFOAM FOAM PROCTOFOAM HC FOAM PROCTO-KIT CREA 2.5% RECTIV OINT | Use PA Form# 20420 | |
| T-CELL ACTIVATION INHIBITOR | | | | | | | | |
| PSORIASIS BIOLOGICALS | MC MC MC MC MC | | ENBREL ^{1,5} ENBREL SURECLICK ¹ HUMIRA ^{1,5} OTEZLA TALTZ ² | MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC MC MC MC MC | | AMJEVITA BIMZELX ³ COSENTYX ⁴ CYLTEZO HADLIMA HULIO HYRIMOZ IDACIO ILUMYA ³ SKYRIZI SOTYKTU SPEVIGO SILIQ STELARA TREMIFYA YUFLYMA YUSIMRY | 1. Dosing limits apply, please refer to dosage consolidation list. 2. Clinical PA required and will be preferred for the indication of plaque psoriasis, psoriatic arthritis and ankylosing spondylitis. 3. For the treatment of adults with moderate-to-severe plaque psoriasis who are candidates for systemic therapy or phototherapy. 4. Please see criteria section | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Cosentyx approvals for 300mg dose(s) must use "300DOSE" package (containing 2 x 150mg pens or syringes). It is recommended to assess for TB infection prior to starting treatment with Taltz®. |

5. Will not require a PA if at least one systemic drug such as methotrexate, cyclosporine, methoxsalen or acitretin is in members drug profile.

[Use PA Form# 20910.](#)

ALTERNATIVE MEDICINES

| | | | | | | | | |
|-----------------------|----------|--|--------------------------------------|--------|--|----------------|-------------------------------------|---|
| ALTERNATIVE MEDICINES | MC MC | | DIMETHYL SULFOXIDE SOLN MELATONIN | MC/DEL | | CO-ENZYME Q-10 | Use PA Form# 20420. | Will only be approved for specific conditions supported by at least two double-blinded, placebo-controlled randomized trials that are not contradicted by other studies of similar quality. |
|-----------------------|----------|--|--------------------------------------|--------|--|----------------|-------------------------------------|---|

CHELATING AGENTS

| | | | | | | | | |
|------------------|--------|--|----------------|------------------------------------|--|--|---|---|
| CHELATING AGENTS | MC/DEL | | CUPRIMINE CAPS | MC MC MC/DEL MC MC/DEL | | CLOVIQUE DEPEN TITRATABS TABS EXJADE ¹ SYPRINE TRIENTINE CAPS | Use PA Form# 20420 1. FDA indication of treatment of chronic iron overload due to blood transfusions in membes 2 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Clovique® should be used when continued treatment with penicillamine is no longer possible because of intolerable or life endangering side effects. |
|------------------|--------|--|----------------|------------------------------------|--|--|---|---|

ANTILEPROTIC

| | | | | | | | | |
|--------------|--|--|--|----|--|----------------------------|--|---|
| ANTILEPROTIC | | | | MC | | THALOMID CAPS ¹ | 1. All PA requests for 150mg dosing will require use of Thalomid 100mg and 50mg capsules. Use PA Form# 20420. | Approved for indications of leprosy, treatment-resistant multiple myeloma and AIDS. |
|--------------|--|--|--|----|--|----------------------------|--|---|

ANTINEOPLASTIC AGENTS

| | | | | | | | | |
|--|--------------------------------------|--|--|--|--|---|--|--|
| ANTINEOPLASTIC AGENTS - ANTIADNDROGENS | MC/DEL | | BICALUTAMIDE | MC/DEL | | CASODEX | Use PA Form# 20420. | |
| ANTINEOPLASTIC AGENTS- LHRH ANALOGS | MC/DEL MC/DEL MC/DEL MC/DEL | | LUPRON DEPOTSYPHNGEKIT ¹ LUPRON DEPOT- PED KIT ¹ (1-month) LUPRON DEPOT-PED SYRINGEKIT (3-month) TRIPTODUR VIAL | MC/DEL MC/DEL MC/DEL MC/DEL MC | | LUPRON DEPOT SYRINGEKIT FIRMAGON ² SUPPRELIN LA (IMPLANT) KIT TRELSTAR VANTAS ² | 1. Dosing limits apply, please refer to dosage consolidation list. 2. PA required to confirm FDA approved indication. Use PA Form# 20420. | |
| ANTINEOPLASTIC AGENTS - TYROSINE KINASE INHIBITORS | | | | MC MC/DEL MC | | SPRYCEL ¹ TYKERB ² GLEEVEC ¹ | Use PA Form# 20420. 1. Verification of diagnosis is required. 2. PA required to confirm FDA approved indication and to monitor for potential drug-drug interactions. | |
| ANTINEOPLASTICS-MISCELLANEOUS | MC MC/DEL MC/DEL | | AMIFOSTINE MERCAPTOPYRINE OXALIPLATIN | MC MC/DEL MC/DEL MC MC/DEL MC/DEL | | DOCEFREZ ELOXATIN ETHYOL LEUPROLIDE PURINETHOL ZOLINZA | Use PA Form# 20420. | |

| | | | | | |
|--|--------|-----------|--------|-----------|------------------------------------|
| ANTINEOPLASTICS- MONOCLONAL ANTIBODIES | MC/DEL | TRAZIMERA | MC/DEL | ENHERTU | |
| | | | MC/DEL | HERCEPTIN | |
| | | | MC/DEL | HERZUMA | |
| | | | MC | KANJINTI | |
| | | | MC | OGIVRI | |
| | | | MC/DEL | ONTRUZANT | Use PA Form# 20420 |

CANCER

| | | | | | | |
|--------|--------|-------------------|-------------------------|----------------------------|---|---|
| CANCER | MC | ALIMTA | MC | ABECMA | <p>1. PA required to confirm appropriate diagnosis and testing.</p> <p>2. Avoid CYP3A drug drug interaction.</p> <p>3. Clinical PA required for appropriate diagnosis</p> <p>4. Re-approval will require documentation of response without disease progression and tolerance to treatment</p> <p>5. Dosing limits apply, please see dosage consolidation list.</p> <p>6. Max daily dose of 300mg.</p> <p>7. Monitor liver enzymes periodically and stop treatment upon Grade 3 or higher elevation of liver enzymes approved indication</p> <p>8. For patients ≥ 12 years of age</p> <p>9. For the treatment of patients up to 25 years of age with B-cell acute lymphoblastic leukemia (ALL) that is refractory or in second or later relapse.</p> <p>Use PA Form# 20420</p> | <p>All non-preferred: A clinical PA is required to confirm appropriate clinical indication for the individual drug request. Specific to each drug all age, clinical testing requirements, previous step therapies, adjunctive drug therapy requirements, and response without disease progression will be also be evaluated for clinical appropriateness. The standard for the appropriate indication will include the FDA label as well as current NCCN guidelines</p> <p>Scemblix is for the treatment of adult patients with: Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP), previously treated with two or more</p> |
| | MC/DEL | ANASTROZOLE TABS | MC | AKEEGA | | |
| | MC | ERBITUX | MC | ALECENSA | | |
| | MC | IMATINIB MESYLATE | MC/DEL | ALIQOPA ³ | | |
| | MC/DEL | LETROZOLE | MC | ALUNBRIG ¹ | | |
| | MC | RUXIENCE | MC | ALYMSYS | | |
| | MC/DEL | VIDAZA | MC/DEL | ARIMIDEX | | |
| | MC | ZIRABEV | MC | AUGTYRO | | |
| | | | MC | AYVAKIT | | |
| | | | MC/DEL | AVASTIN | | |
| | | | MC/DEL | BALVERSA | | |
| | | | MC | BAVENCIO ^{1,8} | | |
| | | | MC/DEL | BENDEKA ³ | | |
| | | | MC/DEL | BESPONS ³ | | |
| | | | MC | BESREMI ¹ | | |
| | | | MC | BLENREP | | |
| | | | MC/DEL | BOSULIF | | |
| | | | MC/DEL | BRAFTOVI ¹ | | |
| | | | MC | BREYANZI | | |
| | | | MC | BRUKINSA | | |
| | | | MC | CABOMETYX ³ | | |
| | | | MC | CAMCEVI | | |
| | | | MC/DEL | CALQUENCE ³ | | |
| | | | MC | COMETRIQ ^{3,4,5} | | |
| | | | MC | COTELLIC | | |
| | | | MC/DEL | COPIKTRA | | |
| | | | MC | DARZALEX ³ | | |
| | | | MC/DEL | DAURISMO | | |
| | | | MC/DEL | ELREXFIO | | |
| | | | MC/DEL | EMPLICITI(IV) ⁸ | | |
| | | | MC | EPKINLY | | |
| | | | MC/DEL | ERLEADA | | |
| | | | MC/DEL | ERIVEDGE | | |
| | | | MC | EXKIVITY | | |
| | | | MC | FARYDAK | | |
| | | MC/DEL | FEMARA | | | |
| | | MC | FOLOTYN | | | |
| | | MC | FOTIVDA | | | |
| | | MC | FRUZAQLA | | | |
| | | MC | GAVRETO | | | |
| | | MC/DEL | GILOTRIF ^{4,5} | | | |
| | | MC/DEL | IBRANCE | | | |
| | | MC | ICLUSIG ³ | | | |
| | | MC/DEL | IDHIFA ³ | | | |
| | | MC | IMBRUVICA | | | |
| | | MC | IMDELLTRA | | | |
| | | MC/DEL | IMFINZI | | | |
| | | MC/DEL | IMJUDO | | | |

| | |
|--------|-------------------------|
| MC | IMLYGIC |
| MC/DEL | INLYTA |
| MC/DEL | INREBIC |
| MC | INQOVI |
| MC | IWILFIN |
| MC | JAKAFI |
| MC | JAYPIRCA ^{1,2} |
| MC | JEMPERLI |
| MC/DEL | KEYTRUDA ¹ |
| MC | KIMMTRAK |
| MC | KISQALI ¹ |
| MC/DEL | KOSELUGO |
| MC | KRAZATI ³ |
| MC | KYMRIAH ^{3,9} |
| MC | KYPROLIS ¹ |
| MC | LARTRUVO ¹ |
| MC | LENVIMA |
| MC/DEL | LIBTAYO ¹ |
| MC | LONSURF |
| MC/DEL | LORBRENA |
| MC | LOQTORZI |
| MC | LUMAKRAS |
| MC/DEL | LUMOXITI ¹ |
| MC | LUNSUMIO ¹ |
| MC | LYNPARZA ¹ |
| MC | LYTGOBI |
| MC | NEXAVAR ¹ |
| MC | NERLYNX ³ |
| MC | NINLARO(PO) |
| MC/DEL | NUBEQA |
| MC | MARGENZA |
| MC/DEL | MEKINIST ^{3,4} |
| MC/DEL | MEKTOVI ¹ |
| MC | MONJUVI |
| MC/DEL | MYLOTARG ³ |
| MC/DEL | MVASI |
| MC | ODOMZO ^{1,2,5} |
| MC | OGSIVEO |
| MC | OJEMDA |
| MC | OJJAARA |
| MC | OMISIRGE |
| MC | ONUREG |
| MC/DEL | OPDIVO ³ |
| MC | OPDUALAG |
| MC | ORGOVYX |
| MC | ORSERDU ^{2,3} |
| MC | PADCEV |
| MC | PEMAZYRE |
| MC | PEPAXTO |
| MC | PHESGO |
| MC/DEL | PIQRAY |
| MC | POLIVY |
| MC | POMALYST |
| MC | PORTRAZZA ³ |
| MC | QINLOCK |
| MC | RETEVMO |

| | |
|--------|-----------------------------|
| MC | REZLIDHIA |
| MC/DEL | ROZLYTREK |
| MC | RUBRACA |
| MC | RITUXAN |
| MC | RYBREVANT |
| MC | RYDAPT |
| MC | RYLAZE |
| MC | RYTELO |
| MC/DEL | SARCLISA |
| MC | SCEMBLIX ¹ |
| MC/DEL | STIVARGA |
| MC/DEL | SUTENT ^{1,2} |
| MC/DEL | SYLATRON |
| MC | TABRECTA |
| MC | TALVEY |
| MC/DEL | TAFINLAR ^{3,4,5,6} |
| MC | TAZVERIK |
| MC/DEL | TALZENNA ¹ |
| MC/DEL | TAGRISSO |
| MC | TECARTUS |
| MC | TECENTRIQ ¹ |
| MC | TEPMETKO |
| MC/DEL | TIBSOVO ¹ |
| MC | TIVDAK |
| MC | TRODELVY |
| MC | TRUSELTIQ |
| MC/DEL | TRUXIMA |
| MC/DEL | TRUQAP |
| MC | TUKYSA |
| MC | UKONIQ |
| MC/DEL | VANFLYTA |
| MC | VEGZELMA |
| MC | VENCLEXTA ³ |
| MC | VERZENIO ³ |
| MC/DEL | VITRAKVI |
| MC/DEL | VIZIMPRO ¹ |
| MC | VONJO |
| MC/DEL | WELIREG |
| MC/DEL | XALKORI |
| MC/DEL | XPOVIO |
| MC/DEL | XOSPATA |
| MC/DEL | XTANDI |
| MC/DEL | YERVOY |
| MC | YESCARTA ³ |
| MC/DEL | ZALTRAP |
| MC | ZEJULA ¹ |
| MC/DEL | ZELBORAF |
| MC | ZEPZELCA |
| MC | ZYDELIG |
| MC/DEL | ZYKADIA |
| MC | ZYNLONTA |
| MC | ZYNYZ ¹ |
| MC | ZYTIGA |

| | | | | | | |
|--------------------|--------|-----------------------|--------|----------|-------------------------------|--|
| IMMUNOSUPPRESSANTS | MC/DEL | CYCLOSPORINE MODIFIED | MC/DEL | CELLCEPT | 1. For the treatment of adult | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered |
|--------------------|--------|-----------------------|--------|----------|-------------------------------|--|

| | | | | | | | | |
|---------------------------|--|--|--|--|--|--|---|---|
| | MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL | | GENGRAF CAPS MYCOPHENOLATE MYFORTIC NEORAL SOL RAPAMUNE SANDIMMUNE TACROLIMUS CAPS | MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL | | CYCLOSPORINE CAPS CYCLOSPORINE SOL. MODIFIED ENVARUSUS XR NEORAL CAP PROGRAF CAPS REZUROCK ¹ ZORTRESS | and pediatric patients 12 years and older with chronic graft-versus-host disease (chronic GVHD) after failure of at least 2 prior lines of systemic therapy Use PA Form# 20420 | on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Cyclosporine will now be non-preferred and require prior authorization if it is currently being used in combination with either Lipitor (doses greater than 20mg/day), Crestor, or lovastatin (doses greater than 20mg). DDI: Cyclosporine will require prior authorization when used with Livalo. DDI: All preferred immunosuppressants will require clinical PA for patients over 60 that are currently on fluoroquinolone therapy. |
| IMMUNOSUPPRESSANTS- Misc. | | | | MC | | HYFTOR ^{1,2} | 1. For the treatment of patients ≥ 6 years of age. 2. Clinical PA required for appropriate diagnosis and clinical parameters. Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| PURINE ANALOG | | | | | | | | |
| PURINE ANALOG | MC MC/DEL | | AZASAN TABS AZATHIOPRINE TABS | MC/DEL | | IMURAN TABS | Use PA Form# 20420 | Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. |
| K REMOVING RESINS | | | | | | | | |
| K REMOVING RESINS | MC/DEL MC/DEL | | LOKELMA SODIUM POLYSTYRENE SULFON | MC/DEL MC/DEL MC | | SPS SUSP SPS 30GM/120ML ENEMA SUSP VELTASSA | Use PA Form# 20420 | |

New drugs are initially non-preferred until reviewed by the DUR Committee and the State. According to State policy, any drug requiring specific diagnosis still requires the specific diagnosis unless otherwise noted within this document.

Last update 01/17

PDL DOSAGE CONSOLIDATION LIST

Tabs/Caps/Patches: Quantities in units

Shaded areas are non-preferred agents - Quantities of these

Sprays/Inhalers/Nebulizers: Quantities in GM, ML, OR MCG

non-preferred agents are available up the limit only with

Injectibles: Quantities in ML

prior authorization

| Drug Name | Strength | Limit/Day | Limit/Days | Drug Name | Strength | Limit/Day | Limit/Days |
|-----------------------------|---------------|---------------|------------|-----------------------|---------------|-------------------|------------|
| ABILIFY SOLUTION | 1MG/ML | 30ML | 1020/34 | ATROVENT HFA | 17MCG | 12 INHALATIONS | 25.8/34 |
| ACCUPRIL | 5MG | 1 | 35/35 | ATROVENT 30ML | 0.03% | 12 SPRAYS | 30/30 |
| ACCUPRIL | 10MG | 1 | 35/35 | ATROVENT 15ML | 0.06% | 16 SPRAYS | 45/30 |
| ACCUPRIL | 20MG | 1 | 35/35 | AVANDIA | 2MG | 1.5 | 53/35 |
| ACEON | 2MG | 1 | 35/35 | AVANDIA | 4MG | 1 | 35/35 |
| ACEON | 4MG | 1 | 35/35 | AVAPRO | 75MG | 1.5 | 53/35 |
| ACTONEL | 5MG | 1 | 35/35 | AVAPRO | 150MG | 1 | 35/35 |
| ACTONEL | 35MG | 1/WK | 5/35 | AXERT (Step 8) | 6.25MG | | 12/30 |
| ACTOS | All Strengths | 1 | 35/35 | AXERT (Step 8) | 12.5MG | | 12/30 |
| ADDERALL XR | 5MG | 3 | 90/30 | AZELEX | 20% | | 1 TUBE/18 |
| ADDERALL XR | 10MG | 3 | 90/30 | AZILECT | All Strengths | 1 | 35/35 |
| ADDERALL XR | 15MG | 3 | 90/30 | BACTROBAN CREAM | | | 1 TUBE/30 |
| ADDERALL XR | 20MG | 2 | 60/30 | BECONASE AQ | 42MCG | 8 INHALATIONS | 50/30 |
| ADDERALL XR | 30MG | 1 | 35/35 | BENICAR-HCT | All Strengths | 1 | 30/30 |
| ADEMPAS | All Strengths | 1 | 35/35 | BENAZEPRIL | 5MG | 1 | 35/35 |
| ADVAIR DISKUS | All Strengths | 2 | 60/30 | BENAZEPRIL | 10MG | 1.5 | 53/35 |
| ADVAIR HFA | All Strengths | 4 | 120/30 | BENAZEPRIL | 20MG | 1 | 35/35 |
| ADZENYS XR | All Strengths | 1 | 30/30 | BENAZEP/HCTZ | 5-6.25 | 1 | 35/35 |
| AEROBID | 250MCG | 8 INHALATIONS | 21/35 | BENAZEP/HCTZ | 10/12.5 | 1 | 35/35 |
| AEROBID-M | 250MCG | 8 INHALATIONS | 21/35 | BEVESPI AERO | | 4 INHALATIONS | 120/30 |
| ALAVERT-NON DROW | TAB | 1 | 96/96 | BONIVA | 2.5MG | 1 | 35/35 |
| ALENDRONATE | All Strengths | 1/WK | 35/35 | BOTOX (ADULTS) | 100U/ML | 1 session/90 days | 600U/90 |
| ALTABAX | 5GM | | 1 TUBE/30 | BOTOX (CHILDREN>12) | 100U/ML | 1 session/90 days | 400U/90 |
| ALTABAX | 15GM | | 1 TUBE/30 | BREO ELLIPTA | 100/25MCG | 1 INHALATIONS | 60/60 |
| ALTABAX | 30GM | | 1 TUBE/30 | BRILINTA | All Strengths | 2 | 70/35 |
| ALTACE | 1.25MG | 1 | 35/35 | BRINTELLIX | All Strengths | 1 | 35/35 |
| ALTACE | 2.5MG | 1 | 35/35 | BUTRANS | | 1 patch/WK | 4/28 |
| ALTACE | 5MG | 1 | 35/35 | BYETTA | 5mcg inj | 0.04ML | 1.2ML/30 |
| AMARYL | 1MG | 1 | 35/35 | BYETTA | 10mcg inj | 0.08ML | 2.4ML/30 |
| AMARYL | 2MG | 1 | 35/35 | CALAN SR | 120MG | 1 | 35/35 |
| AMBIEN | 5MG | | 12/34 | CALAN SR | 180MG | 2 | 70/35 |
| AMBIEN | 10MG | | 12/34 | CALAN SR | 240MG | 2 | 70/35 |
| AMBIEN CR | 6.25MG | | 12/34 | CARDIZEM CD | 120MG/24 | 1 | 35/35 |
| AMBIEN CR | 12.5MG | | 12/34 | CARDIZEM CD | 180MG/24 | 1 | 35/35 |
| AMERGE (Step 8) | 1MG | | 12/30 | CARDIZEM CD | 240MG/24 | 1 | 35/35 |
| AMERGE (Step 8) | 2.5MG | 2.5MG | 12/30 | CARDIZEM CD | 300MG/24 | 1 | 35/35 |
| AMLODIPINE | 2.5MG | 1.5 | 53/35 DAYS | CARDIZEM CD | 360MG/24 | 1 | 35/35 |
| AMLODIPINE | 5MG | 1.5 | 53/35 DAYS | CARDIZEM LA | 120MG/24 | 1 | 35/35 |
| AMMONIUM LACTATE CREA | 12% | | 1 TUBE/10 | CARDIZEM LA | 180MG/24 | 1 | 35/35 |
| AMMONIUM LACTATE LOTN | 12% | | 1TUBE/8 | CARDIZEM LA | 240MG/24 | 1 | 35/35 |
| AMPHETAMINE/DEXTROAMPHET ER | 5MG | 3 | 90/30 | CARDIZEM LA | 300MG/24 | 1 | 35/35 |
| AMPHETAMINE/DEXTROAMPHET ER | 10MG | 3 | 90/30 | CARDIZEM LA | 360MG/24 | 1 | 35/35 |
| AMPHETAMINE/DEXTROAMPHET ER | 15MG | 3 | 90/30 | CARDURA | 1MG | 1 | 35/35 |
| AMPHETAMINE/DEXTROAMPHET ER | 20MG | 2 | 60/30 | CARDURA | 2MG | 1.5 | 53/35 |
| AMPHETAMINE/DEXTROAMPHET ER | 30MG | 1 | 90/90 | CARDURA | 4MG | 1.5 | 53/35 |
| AMPHETAMINE SALT | 5,10,15MG | 3 | 105/35 | CARTIA XT | 120MG | 1 | 90/90 |
| AMPHETAMINE SALT | 20MG | 2 | 70/35 | CARTIA XT | 180MG | 1 | 90/90 |
| AMPHETAMINE SALT | 30MG | 1 | 35/35 | CARTIA XT | 240MG | 1 | 90/90 |
| ANDRODERM | 2.5MG | 2 | 60/30 | CARTIA XT | 300MG | 1 | 90/90 |
| ANDRODERM | 5MG | 1 | 30/30 | CATAPRES-TTS1 | 0.1 MG/24HR | | 5/35 |
| ARAVA | 10MG | 1 | 35/35 | CATAPRES-TTS2 | 0.2 MG/24HR | | 5/35 |
| ARCAPTA | 75MCG | 1 INHALATION | 35/35 | CATAPRES-TTS3 | 0.3 MG/24HR | | 5/35 |
| ARICEPT | 5MG | 1 | 35/35 | CEFIXIME | 400MG | 2 | 2/7 |
| ARICEPT | 10MG | 1 | 35/35 | CELEBREX | 100MG | 1 | 35/35 |
| ARIPIRAZOLE | 2MG | 2 | 180/90 | CELEBREX | 200MG | 2 | 70/35 |
| ARIPIRAZOLE | 5MG | 2 | 180/90 | CELEBREX | 400MG | 1 | 35/35 |
| ARIPIRAZOLE | 10MG | 2 | 180/90 | CELEXA | 20mg | 0.5 | 17/34 |
| ARIPIRAZOLE | 15MG | 2 | 180/90 | CELEXA | 40mg | 1 | 51/34 |
| ARIPIRAZOLE | 20MG | 1.5 | 135/90 | CITALOPRAM | 10MG | 2 | 180/90 |
| ARIPIRAZOLE | 30MG | 1 | 90/90 | CITALOPRAM | 20MG | 2 | 180/90 |
| ARIXTRA INJECTION | 2.5MG/0.5ML | | 7/30 | CITALOPRAM | 40MG | 1 | 90/90 |
| ARIXTRA INJECTION | 5MG/0.4ML | | 7/30 | CLARINEX | REDI TAB | 1 | 35/35 |
| ARIXTRA INJECTION | 7.5MG/0.6ML | | 7/30 | CLEOCIN-T | | 1 PACKAGE | 1/30 |
| ARIXTRA INJECTION | 10MG/0.8ML | | 7/30 | CLINDAMYCIN PHOSPHATE | | 1 PACKAGE | 1/30 |
| ARONAIR | All Strengths | 1 INHALATION | 60U/30 | COMBIVENT | 103-18MCG | 12 INHALATIONS | 30/35 |
| ASMANEX 30 UNITS | 220MCG | 1 INHALATION | 30U/30 | Drug Name | Strength | Limit/Day | Limit/Days |
| ASMANEX 60 UNITS | 220MCG | 2 INHALATIONS | 60U/30 | EFFEXOR XR | 37.5MG | 1 | 35/35 |
| ASMANEX 120 UNITS | 220MCG | 4 INHALATIONS | 120U/30 | EFFEXOR XR | 75MG | 1 | 35/35 |
| ATACAND | 4MG | 1.5 | 53/35 | EMSAM | All Strengths | 1 | 34/34 |
| ATACAND | 8MG | 1.5 | 53/35 | ENALAPRIL | 2.5 | 1 | 90/90 |

| Drug Name | Strength | Limit/Day | Limit/Days |
|-------------------|-------------------|-----------|-----------------|
| ATACAND | 16MG | 1 | 35/35 |
| ATRIPLA | 600MG | 1 | 35/35 |
| Drug Name | Strength | Limit/Day | Limit/Days |
| COMETRIQ | 80MG | 1 | 35/35 |
| COMETRIQ | 20MG | 3 | 105/35 |
| CONCERTA | 18MG | 1 | 30/30 |
| CONCERTA | 27MG | 1 | 30/30 |
| CONCERTA | 36MG | 2 | 60/30 |
| COPAXONE INJ | 20MG | | 1/32 |
| COPAXONE KIT | 20MG/ML | | 1/30 |
| COREG CR | All Strengths | 1 | 34/34 |
| COSENTYX | 150MG | 1 | 1/30 |
| CRESTOR | 5MG | 1 | 35/35 |
| CRESTOR | 10MG | 1 | 35/35 |
| CRESTOR | 20MG | 1 | 35/35 |
| CRESTOR | 40MG | 1 | 35/35 |
| CYMBALTA | All Strengths | 1 | 35/35 |
| DALMANE | 15MG | | 10/30 |
| DALMANE | 30MG | | 10/30 |
| DAYPRO | 600MG | 2 | 70/35 |
| DAYTRANA | 10mg/9hr (27.5mg) | 1 | 34/34 |
| DAYTRANA | 15mg/9hr (41.3mg) | 1 | 34/34 |
| DAYTRANA | 20mg/9hr (55.0mg) | 1 | 34/34 |
| DAYTRANA | 30mg/9hr (82.5mg) | 1 | 34/34 |
| DDAVP | 5ML | | 15/34 |
| DENAVIR CREAM | | | 2gm/30 |
| DEPO-PROVERA | 150MG/ML | | 1/90 |
| DEPO-PROVERA | 400MG/ML | | 2.5/90 |
| DEPO-TESTOSTERONE | 200MG/ML | | 20/90 |
| DESMOPRESSIN | 0.1MG | 12 | 420/35 |
| DESMOPRESSIN | 0.2MG | 6 | 210/35 |
| DESONIDE | 0.05% | | 2 TUBES/30 |
| DESOWEN | 0.05% | | 2 TUBES/30 |
| DETROL LA | 2MG | 1 | 35/35 |
| DEXEDRINE | All Strengths | 3 | 90/30 |
| DEXILANT | All Strengths | 1 | 35/35 |
| DEXTROAMPHETAMINE | All Strengths | 3 | 90/30 |
| DICLOFENAC 1% GEL | 1% GEL | | 2 TUBES/30 |
| DIFLUCAN | 150MG | | 1/7 |
| DILACOR XR | 240MG/24 | 1 | 35/35 |
| DILACOR XR | 120MG/24 | 1 | 35/35 |
| DILACOR XR | 180MG/24 | 1 | 35/35 |
| DILTIA - XT | 120MG/24 | 1 | 90/90 |
| DILTIA - XT | 180MG | 1 | 90/90 |
| DILTIA - XT | 240MG/24 | 1 | 90/90 |
| DILTIAZEM CAP ER | 120MG | 1 | 90/90 |
| DILTIAZEM CAP XR | 120MG | 1 | 90/90 |
| DILTIAZEM CAP | 120MG/24 | 1 | 90/90 |
| DILTIAZEM CAP | 180MG/24 | 1 | 90/90 |
| DILTIAZEM CAP ER | 240MG | 1 | 90/90 |
| DILTIAZEM CAP XR | 240MG | 1 | 90/90 |
| DILTIAZEM XR CAP | 240MG/24 | 1 | 90/90 |
| DILTIAZEM CAP | 240MG/24 | 1 | 90/90 |
| DILTIAZEM CAP | 300MG/24 | 1 | 90/90 |
| DILTIAZEM CAP | 360MG/24 | 1 | 90/90 |
| DIOVAN | 80MG | 1 | 35/35 |
| DIOVAN - HCT | 80 - 12.5 | 1 | 35/35 |
| DITROPAN XL | 5MG | 1 | 35/35 |
| DITROPAN XL | 10MG | 2 | 70/35 |
| DORAL | 7.5MG | | 10/30 |
| DOXAZOSIN | 1MG | 1 | 90/90 |
| DOXAZOSIN | 2MG | 1.5 | 135/90 |
| DOXAZOSIN | 4MG | 1.5 | 135/90 |
| DRYSOL SOL | 20% | | 1 BOTTLE/30DAYS |
| DURAGESIC PATCHES | 12.5MCG/HR | | 11/33 |
| DURAGESIC PATCHES | 25MCG/HR | | 11/33 |
| DURAGESIC PATCHES | 50MCG/HR | | 11/33 |
| DURAGESIC PATCHES | 75MCG/HR | | 11/33 |
| DURAGESIC PATCHES | 100MCG/HR | | 22/33 |
| DULOXETINE | 20MG | 3 | 270/90 |
| DULOXETINE | 30MG | 3 | 270/90 |
| DULOXETINE | 60MG | 2 | 180/90 |
| EDEX | All Strengths | | 1/30 |
| Drug Name | Strength | Limit/Day | Limit/Days |
| ILARIS | | | 2/28 |

| Drug Name | Strength | Limit/Day | Limit/Days |
|--|---------------|-------------------|-------------------|
| ENALAPRIL | 5MG | 1.5 | 135/90 |
| ENALAPRIL | 10MG | 1.5 | 135/90 |
| ENALAPR/HCTZ | 5-12.5 | 1 | 90/90 |
| ENBREL | 25MG/ML | | 8/28 |
| ENBREL SURECLICK | | | 8/28 |
| ESTAZOLAM | 1MG | | 10/30 |
| ESTAZOLAM | 2MG | | 10/30 |
| ESTRING MIS | 2MG | | 1/90 |
| EVENITY | | 12 DOSES/LIFETIME | 12 DOSES/LIFETIME |
| EVOTAZ | All Strengths | 1 | 30/30 |
| FELODIPINE | 2.5MG | 1 | 90/90 |
| FELODIPINE | 5MG | 1.5 | 135/90 |
| FENTANYL | 25MCG/HR | | 11/33 |
| FENTANYL | 50MCG/HR | | 11/33 |
| FENTANYL | 75MCG/HR | | 11/33 |
| FENTANYL | 100MCG/HR | | 22/33 |
| FETZIMA | All Strengths | 1 | 35/35 |
| FINASTERIDE | 5MG | 1 | 90/90 |
| FLONASE | 50MCG | 4 SPRAYS | 32/34 |
| FLOVENT HFA 44MCG | 44MCG | 4 INHALATIONS | 10.6/30 |
| FLOVENT HFA 110MCG | 110MCG | 4 INHALATIONS | 12/30 |
| FLOVENT HFA 220MCG | 220MCG | 8 INHALATIONS | 24/30 |
| FLOVENT DISKUS | 50MCG, 100MCG | 4 INHALATIONS | 60/30 |
| FLOVENT DISKUS | 250MCG | 3 INHALATIONS | 120/30 |
| FLUCONAZOLE | 150MG | | 1/7 |
| FLUNISOLIDE SOLN | 0.025% | 16 SPRAYS | 75/30 |
| FLUOXETINE CAP | 40MG | 2 | 180/90 |
| FLUOXETINE CAP | 20MG | 4 | 360/90 |
| FLUOXETINE CAP | 10MG | 3 | 270/90 |
| FLURAZEPAM | 15MG | | 10/30 |
| FLURAZEPAM | 30MG | | 10/30 |
| FLUTICASONE SPR | | 4 SPRAYS | 48/90 |
| FLUVOXAMINE | 25MG | 3 | 270/90 |
| FLUVOXAMINE | 50MG | 3 | 270/90 |
| FOCALIN | All Strengths | 3 | 105/35 |
| FOCALIN XR | All Strengths | 1 | 35/35 |
| FORFIVO XL | All Strengths | 1 | 35/35 |
| FOSAMAX | 5MG | 1 | 35/35 |
| FOSAMAX | 10MG | 1 | 35/35 |
| FOSAMAX | 70MG | 1/WK | 5/35 |
| FOSAMAX | 40MG | 2/WK | 10/35 |
| FOSINOPRIL | 10MG | 1.5 | 135/90 |
| FOSINOPRIL | 20MG | 2 | 180/90 |
| FRAGMIN INJ | 10000U/ML | 2ML | 14/7 |
| FRAGMIN INJ | 2500U/.2ML | 0.4ML | 2.80/7 |
| FRAGMIN INJ | 25000U/ML | 0.8ML | 5.6/7 |
| FRAGMIN INJ | 5000U/.2ML | 0.4ML | 2.80/7 |
| FRAGMIN INJ | 7500U/.3ML | 0.6ML | 4.2/7 |
| FROVA TAB (Step 8) | 2.5MG | | 12/30 |
| FULYZAQ | 125MG | 2 | 70/35 |
| FUZEON | KIT | 1 | 1/30 |
| FYCOMPA | All Strengths | 1 | 35/35 |
| GABAPENTIN | 300MG | 9 | 810/90 |
| GABAPENTIN | 400MG | 9 | 810/90 |
| GABAPENTIN | 600MG | 6 | 540/90 |
| GABAPENTIN | 800MG | 4 | 360/90 |
| GEODON | 20MG | 2 | 70/35 |
| GEODON | 40MG | 2 | 70/35 |
| GEODON | 60MG | 2 | 70/35 |
| GEODON | 80MG | 2 | 70/35 |
| GEODON | INJ | 2 | 70/35 |
| GILOTRIF | All Strengths | 1 | 35/35 |
| GLIMEPIRIDE | 1MG | 1 | 90/90 |
| GLIMEPIRIDE | 2MG | 1 | 90/90 |
| GLUCOSE TES STRP | | 12 | 420/35 |
| GLUCAGEN INJ. HYPOKIT | | | 2/30 |
| GLYCOLAX* | 255GM | | 255GM/90 |
| * Available for once daily dosing to members under the age of 18 years | | | |
| Drug Name | Strength | Limit/Day | Limit/Days |
| LUNESTA | 2MG | | 12/34 |
| LUNESTA | 3MG | | 12/34 |
| LUPRON DEPOT INJ | 11.25MG | KIT | 1/90 |
| LUPRON DEPOT INJ | 22.5 | KIT | 1/90 |
| LUPRON DEPOT INJ | 30MG | | 1/90 |

| | | | |
|---------------------|---------------|---------------|-----------------|
| HALCION | 0.125MG | | 10/35 |
| HALCION | 0.25 | | 10/35 |
| HUMIRA | 40mg/0.8ml | | 4/28 |
| HYDROXYZINE TAB | All Strengths | 3 | 270/90 |
| HYTRIN | 1MG | 1 | 35/35 |
| HYTRIN | 5MG | 1 | 35/35 |
| HYZAAR | 50-12.5 | 1 | 35/35 |
| IMDUR | 30MG | 1.5 | 53/35 |
| IMDUR | 60MG | 1.5 | 53/35 |
| IMITREX (step 8) | 25MG | | 12/30 |
| IMITREX (step 8) | 50MG | | 12/30 |
| IMITREX (step 8) | 100MG | | 12/30 |
| IMITREX VIAL | All Strengths | | 6 boxes/30 |
| IMITREX CARTRIDGE | All Strengths | | 12/30 |
| IMITREX NASAL SPRAY | All Strengths | | 12/30 |
| IMITREX PEN INJCTR | All Strengths | | 12/30 |
| IMIQUIMOD | 5% | | 12/30 |
| IMIQUIMOD | 5% | | 12/30 |
| INTAL | 800MCG | 8 INHALATIONS | 28.4/34 |
| INVOKANA | All Strengths | 1 | 35/35 |
| IPRATROPIUM 30ML | 0.03% | 12 SPRAYS | 90/90 |
| IPRATROPIUM 15ML | 0.06% | 16 SPRAYS | 135/90 |
| ISOPTIN SR | 180MG | 2 | 70/35 |
| IRBESARTAN | All Strengths | 1 | 90/90 |
| ISOPTIN SR | 240MG | 2 | 70/35 |
| ISOSORBIDE MONO | 30MG | 2 | 180/90 |
| ISOSORBIDE MONO | 60 MG | 1.5 | 135/90 |
| JANUMET | All Strengths | 2 | 70/35 |
| JANUVIA | All Strengths | 1 | 35/35 |
| JUVISYNC | All Strengths | 1 | 35/35 |
| KETOPROFEN | 100MG | 2 | 180/90 |
| KETOPROFEN | 200MG | 1 | 90/90 |
| KETOROLAC | 10MG | 4.8 | 24/30 |
| KHEDEZLA | All Strengths | 1 | 35/35 |
| LAC-HYDRIN CREAM | 12% | | 1TUBE/30 |
| LAMICTAL | 25MG | 6 | 210/35 |
| LAMICTAL | 25MG CHW | 6 | 210/35 |
| LAMICTAL | 100MG | 2 | 70/35 |
| LAMISIL | 250MG | 1 | 35/35 |
| LAMOTRIGINE | 25MG | 6 | 540/90 |
| LAMOTRIGINE | 100MG | 2 | 180/90 |
| LANSOPRAZOLE CAPS | All Strengths | 1 | 90/90 |
| LATUDA | All Strengths | 1 | 17/34 |
| LESCOL | 20MG | 1 | 35/35 |
| LEVAQUIN | 250MG | 1 | 35/35 |
| LEXAPRO | 5MG | 0.5 | 15/30 |
| LIPITOR | 10MG | 1 | 35/35 |
| LIPITOR | 20MG | 1 | 35/35 |
| LIPITOR | 40MG | 1.5 | 53/35 |
| LISINOP/HCTZ | 10/12.5MG | 1 | 90/90 |
| LINEZOLID | 600mg | | 14/60 |
| LOSARTAN | All Strengths | 1 | 90/90 |
| LOSARTAN- HCT | All Strengths | 1 | 90/90 |
| LOTENSIN | 5MG | 1 | 35/35 |
| LOTENSIN | 10MG | 1.5 | 35/35 |
| LOTENSIN | 20MG | 1 | 53/35 |
| LOTENSIN - HCT | 5 - 6.25 | 1 | 35/35 |
| LOTENSIN - HCT | 10 - 12.5 | 1 | 35/35 |
| LOVASTATIN | 10MG | 1.5 | 135/90 |
| LOVASTATIN | 20MG | 1.5 | 135/90 |
| LOVENOX INJ | 30MG/.3ML | 0.6 | 14 injections/7 |
| LOVENOX INJ | 40MG/.4ML | 0.8 | 14 injections/7 |
| LOVENOX INJ | 60MG/.6ML | 1.2 | 14 injections/7 |
| LOVENOX INJ | 80MG/.8ML | 1.6 | 14 injections/7 |
| LOVENOX INJ | 100MG/ML | 2 | 14 injections/7 |
| LOVENOX INJ | 120MG/.8ML | 1.6 | 14 injections/7 |
| LOVENOX INJ | 150MG/ML | 2 | 14 injections/7 |
| LUNESTA | 1MG | | 12/34 |
| Drug Name | Strength | Limit/Day | Limit/Days |
| NIFEDIPINE ER | 90MG | 1 | 90/90 |
| NIFEDIPINE ER,CR | 30MG | 1 | 90/90 |
| NORVASC | 2.5MG | 1.5 | 53/35 DAYS |
| NORVASC | 5MG | 1.5 | 53/35 DAYS |
| NURTEC ODT | All Strengths | | 8/30 |
| NUVARING | | 1/MO | 1/28 |

| | | | |
|----------------------|---------------|----------------|---------------|
| LUPRON DEPOT INJ | 30MG | KIT | 1/90 |
| LYRICA | 25,50,75MG | 3 | 102/35 |
| LYRICA | 100,150,200MG | 3 | 102/35 |
| LYRICA | 225,300MG | 2 | 70/35 |
| MAVIK | 1MG | 1 | 35/35 |
| MAVIK | 2MG | 1 | 35/35 |
| MAXAIR AUTO | 200MCG | 12 INHALATIONS | 14/30 |
| MAXALT (step 8) | 5MG | | 12/30 |
| MAXALT (step 8) | 10MG | | 12/30 |
| MAXALT MLT (step 1) | 5MG | | 12/30 |
| MAXALT MLT (step 1) | 10MG | | 12/30 |
| MEDROXYPR AC | 150MG/ML | | 1/90 |
| MELOXICAM TABS | All Strengths | 1 | 90/90 |
| METADATE ER | 10,20MG | 3 | 90/30 |
| METFORMIN ER | 500MG | 4 | 360/90 |
| METHYLIN | All Strengths | 3 | 90/30 |
| METHYLPHENIDATE ER | 36mg | 2 | 180/90 |
| METHYLPHENIDATE | All Strengths | 3 | 90/30 |
| METROCREAM | | 1 PACKAGE | 1/30 |
| METROGEL | | 1 PACKAGE | 1/30 |
| METROLOTION | | 1 PACKAGE | 1/30 |
| METRONIDAZOLE CREAM | | 1 PACKAGE | 1/30 |
| METRONIDAZOLE GEL | | 1 PACKAGE | 1/30 |
| METRONIDAZOLE LOTION | | 1 PACKAGE | 1/30 |
| MEVACOR | 10MG | 1.5 | 53/35 |
| MEVACOR | 20MG | 1.5 | 53/35 |
| MIACALCIN | | 3.75ml | 1 bottle/34 |
| MICARDIS | All Strengths | 1 | 30/30 |
| MICARDIS-HCT | All Strengths | 1 | 30/30 |
| MIGRANAL NASAL SPRAY | All Strengths | | 12/30 |
| MIRALAX | 255G | 8.5G | 1 bottle/30 |
| MIRALAX | 17G/PACKET | 0.5 packet | 15 packets/30 |
| MIRTAZAPINE | 15mg | 3 | 270/90 |
| MOBIC | 7.5 MG | 1 | 35/35 |
| MOBIC | 15MG | 1 | 35/35 |
| MOEXIPRIL | 7.5 | 1.5 | 135/90 |
| MONOPRIL | 10MG | 1.5 | 53/35 |
| MONOPRIL | 20MG | 2 | 70/35 |
| MUPIROCIN | | | 1 TUBE/30 |
| NABUMETONE | 500MG | 2 | 180/90 |
| NABUMETONE | 750MG | 2 | 180/90 |
| NARATRIPTAN | | | 12/30 |
| NASACORT AERS | 55 MCG | 4 SPRAYS | 9.3/25 |
| NASONEX | 50MCG | 4 SPRAYS | 17/30 |
| NATROBA | | 120ML | 1 bottle/30 |
| NAYZILAM | All Strengths | | 5/30 |
| NEUPOGEN INJ | 300MCG/ML | | 10/30 |
| NEUPOGEN INJ | 480MCG/1.6 | | 16/30 |
| NEUPOGEN INJ | 300MCG/.5ML | | 5/30 |
| NEUPOGEN INJ | 480MCG/.8ML | | 8/30 |
| NEURONTIN | 300MG | 9 | 315/35 |
| NEURONTIN | 600MG | 9 | 315/35 |
| NEXIUM | 20MG | 1 | 35/35 |
| NEXIUM | 40MG | 2 | 70/35 |
| NEXIUM SUS | All Strengths | 1 | 30/30 |
| NIFEDIPINE CR | 90MG | 1 | 90/90 |
| NIFEDIPINE ER | 60MG | 1 | 90/90 |
| NIFEDIPINE ER | 30MG | 1 | 90/90 |
| NIFEDIPINE ER | 60MG | 1 | 90/90 |
| Drug Name | Strength | Limit/Day | Limit/Days |
| RELPAX | All Strengths | | 12/30 |
| REMODULIN | All Strengths | | 1 MDV/30 |
| RESTORIL | 7.5MG | | 10/30 |
| RESTORIL | 15MG | | 10/30 |
| RESTORIL | 30MG | | 10/30 |
| RETIN-A | | 1 TUBE | 1 TUBE/30 |
| REVLIMID | All Strengths | 1 | 35/35 |
| REYVOW | All Strengths | | 4/30 |
| RHINOCORT AQ | 32MCG | 8 SPRAYS | 18/30 |
| REFRESH PLUS | | 15 ML | 1 bottle/30 |
| REFRESH PLUS | | 30 ML | 2 bottles/30 |
| REFRESH TEARS | | 15 ML | 1 bottle/30 |
| REFRESH TEARS | | 30 ML | 2 bottles/30 |
| RESCULA | | | 2 bottles/35 |

| | | | |
|-------------------------|---------------|----------------|--------------|
| ODOMZO | 200mg | 1 | 30/30 |
| OLMESARTAN | All Strengths | 1 | 90/90 |
| OLANZAPINE | 2.5MG | 3 | 270/90 |
| OLANZAPINE | 5MG | 3 | 270/90 |
| OLANZAPINE | 7.5MG | 3 | 270/90 |
| OLANZAPINE | 10MG | 3 | 270/90 |
| OLANZAPINE | 15MH | 2 | 180/90 |
| OLANZAPINE | 20MG | 1.5 | 135/90 |
| OLANZAPINE ODT | All Strengths | 1 | 90/90 |
| OMEPRAZOLE | 10MG | 1 | 90/90 |
| OMEPRAZOLE | 20MG | 1 | 90/90 |
| OMEPRAZOLE | 40MG | 1 | 90/90 |
| OMNARIS | 50MCG | 4 sprays | 12.5/30 |
| ONGLYZA | All Strengths | 1 | 35/35 |
| OPSUMIT | All Strengths | 1 | 35/35 |
| ORUVAIL | 100MG | 2 | 70/35 |
| ORUVAIL | 200MG | 1 | 35/35 |
| OXAPROZIN | 600MG | 2 | 180/90 |
| OXYCODONE ER | 10,20,40MG | 2 | 70/35 |
| OXYCODONE ER | 80MG | 4 | 140/35 |
| OXYCONTIN** | 10,20,30,40MG | 2 | 70/35 |
| OXYCONTIN** | 80MG | 4 | 140/35 |
| PANTOPRAZOLE | All Strengths | 1 | 90/90 |
| PAROXETINE | 10MG | 2 | 180/90 |
| PAROXETINE | 20MG | 2 | 180/90 |
| PAXIL | 10MG | 1.5 | 53/35 |
| PAXIL | 20MG | 1 | 35/35 |
| PEGASYS KIT | | KIT | 1/28 |
| PLAN B | | | 2/15 or 4/30 |
| PLENDIL | 2.5MG | 1 | 35/35 |
| PLENDIL | 5MG | 1.5 | 53/35 |
| PRAVACHOL | 10MG | 1 | 35/35 |
| PRAVACHOL | 20MG | 1 | 35/35 |
| PRAVACHOL | 40MG | 1 | 35/35 |
| PRAVACHOL | 80MG | 1 | 35/35 |
| PRAVASTATIN | 10MG | 1 | 35/35 |
| PRAVASTATIN | 20MG | 1 | 35/35 |
| PRAVASTATIN | 40MG | 2 | 180/90 |
| PRAVASTATIN | 80MG | 1 | 35/35 |
| PREVPAC MIS | 500MG-30MG | | 14/30 |
| PRILOSEC OTC | 20MG | 2 | 168/84 |
| PRINIVIL | 2.5MG | 1 | 35/35 |
| PRINIVIL | 5MG | 1 | 35/35 |
| PRINIVIL | 10MG | 1.5 | 53/35 |
| PRINIVIL | 20MG | 1.5 | 53/35 |
| PRINZIDE | 10-12.5 | 1 | 35/35 |
| PROAIR HFA | 90mcg | 12 INHALATIONS | 17/34 |
| PROTONIX | 20MG | 2 | 70/35 |
| PROTONIX | 40MG | 2 | 70/35 |
| PROZAC | 10MG | 1.5 | 53/35 |
| PULMICORT | 200MCG | 8 INHALATIONS | 1/25 |
| PULMICORT FLEX | All Strengths | 8 Inhalations | 2/30 |
| QUETIAPINE | 25MG | 3 | 270/90 |
| QUETIAPINE | 50MG | 3 | 270/90 |
| QUETIAPINE | 100MG | 3 | 270/90 |
| QUETIAPINE | 200MG | 3 | 270/90 |
| QUINAPRIL | 5MG | 1 | 90/90 |
| QUINAPRIL | 10MG | 1 | 90/90 |
| QUINAPRIL | 20MG | 1 | 90/90 |
| QVAR AERS | All Strengths | 8 Inhalations | 14.6/25 |
| RANITIDINE SYRUP*** | 15MG/ML | 20ML | 700ML/35 |
| RELAFEN | 500MG | 2 | 70/35 |
| RELAFEN | 750MG | 2 | 70/35 |
| REMERON | 15MG | 1.5 | 53/35 |
| Drug Name | Strength | Limit/Day | Limit/Days |
| SULAR | 10MG | 1.5 | 53/35 |
| SULAR | 20MG | 1 | 35/35 |
| SUMATRIPTAN PEN INJ | All Strengths | | 12/30 |
| SUMATRIPTAN NASAL SPRAY | All Strengths | | 12/30 |
| SUMATRIPTAN SYRINGE | All Strengths | | 12/30 |
| SUMATRIPTAN TAB | All Strengths | | 12/30 |
| SYNVISC INJ | 8MG/ML | | 2/30 |
| SYRINGES | | 10 | 1000/100 |
| TAFINLAR | 50MG | 6 | 210/35 |
| TAFINLAR | 75MG | 4 | 140/35 |

| | | | |
|-------------------|---------------|---------------|----------|
| REYATAZ | All Strengths | 1 | 35/35 |
| RISPERDAL | 0.5MG | 1.5 | 53/35 |
| RISPERDAL | 0.25MG | 1.5 | 53/35 |
| RISPERDAL | 1MG | 1.5 | 53/35 |
| RISPERDAL | 2MG | 1.5 | 53/35 |
| RISPERDAL | 3MG | 2 | 70/35 |
| RISPERDAL | 4MG | 2 | 70/35 |
| RISPERDAL INJ | 25MG | | 2/28 |
| RISPERDAL INJ | 37.5 | | 2/28 |
| RISPERDAL INJ | 50MG | | 2/28 |
| RISPERDAL M-TAB | 0.5MG | 1.5 | 53/35 |
| RISPERDAL M-TAB | 1MG | 1.5 | 53/35 |
| RISPERDAL M-TAB | 2MG | 4 | 140/35 |
| RISPERDAL SOL. | 1MG/ML | 8ML | 280/35 |
| RISPERIDONE | 0.5MG | 3 | 270/90 |
| RISPERIDONE | 0.25MG | 3 | 270/90 |
| RISPERIDONE | 1MG | 3 | 270/90 |
| RISPERIDONE | 2MG | 3 | 270/90 |
| RISPERIDONE | 3MG | 2 | 180/90 |
| RISPERIDONE | 4MG | 2 | 180/90 |
| RISPERIDONE SOL. | 1MG/ML | 8ML | 280/35 |
| RITALIN LA | All Strengths | 1 | 35/35 |
| RITALIN LA | 30mg | 2 | 70/35 |
| SAVELLA | All Strengths | 2 | 70/35 |
| SEREVENT DISKUS | 50MCG | 2 INHALATIONS | 60/30 |
| SEROQUEL | 100MG | | 45/30 |
| SEROQUEL XR | 150MG | 1 | 35/35 |
| SEROQUEL XR | 200MG | 1 | 35/35 |
| SEROQUEL XR | 300MG | 2 | 70/35 |
| SEROQUEL XR | 400MG | 2 | 70/35 |
| SERTRALINE | 25MG | 3 | 270/90 |
| SERTRALINE | 50MG | 3 | 270/90 |
| SERTRALINE | 100MG | 3 | 270/90 |
| SIMVASTATIN | 5MG | 1 | 35/35 |
| SIMVASTATIN | 10MG | 1.5 | 53/35 |
| SIMVASTATIN | 20MG | 1.5 | 53/35 |
| SIMVASTATIN | 40MG | 1.5 | 53/35 |
| SIMVASTATIN | 80MG | 1 | 35/35 |
| SINGULAIR | 4MG | 1 | 35/35 |
| SINGULAIR | 5MG | 1 | 35/35 |
| SINGULAIR | 10MG | 1 | 35/35 |
| SONATA | 5MG | | 12/34 |
| SONATA | 10MG | | 12/34 |
| SPIRIVA | HANDIHLR | 1 INHALTION | 30/30 |
| SPORANOX SOL | 10MG/ML | 10ML/ML | 300cc/30 |
| SPORANOX PULSEPAK | F | | 30/30 |
| SPORANOX | 100MG | | 30/30 |
| STADOL INJ | 1MG/ML | | 9/35 |
| STADOL INJ | 2MG/ML | | 9/35 |
| STRATTERA | All Strengths | 1 | 35/35 |
| SUPRAX | 400MG | 1 | 1/7 |

| Drug Name | Strength | Limit/Day | Limit/Days |
|------------------|---------------|----------------|---------------|
| XOPENEX HFA | | 12 INHALATIONS | 2 INHALERS/34 |
| XOPENEX NEB | | 12CC | 408/34 |
| ZALEPLON | All Strengths | | 30/30 |
| ZECUITY | 6.5 | | 4/28 |
| ZEMBRACE | All Strengths | | 3boxes/30 |
| ZESTORETIC | 10-12.5 | 1 | 35/35 |
| ZESTRIL | 2.5MG | 1 | 35/35 |
| ZESTRIL | 5MG | 1 | 35/35 |
| ZESTRIL | 10MG | 1.5 | 53/35 |
| ZESTRIL | 20MG | 1.5 | 53/35 |
| ZETONNA | 37MCG | 2 | 60/30 |
| ZIPRASIDONE | 20MG | 3 | 270/90 |
| ZIPRASIDONE | 40MG | 3 | 270/90 |
| ZOCOR | 5MG | 1 | 35/35 |
| ZOCOR | 10MG | 1.5 | 53/35 |
| ZOCOR | 20MG | 1.5 | 53/35 |
| ZOCOR | 40MG | 1.5 | 53/35 |
| ZOFRAN* | 4MG | 3 | 90/30 |
| ZOFRAN* | 8MG | 1.5 | 45/30 |
| ZOFRAN* | 24MG | 0.5 | 15/30 |
| ZOFRAN* | 4MG/5ML | 15ML | 450/30 |
| ZOLMITRIPTAN TAB | All Strengths | | 12/30 |

Pain Management Policy

[Back to Index](#)

Beginning January 2017, all current opiate users who are above the maximum combined daily dose of 100 MME must titrate their total daily dose of opioid medications below 300 MME. Also, the maximum daily supply of an opiate prescription for acute pain will be limited to 7-day supplies. The maximum day supply of an opiate prescription for chronic pain will be limited to 30-day supplies. As of July 1, 2017 all users of opioid medications must comply with the maximum combined daily dose of 100 MME.

However, for MaineCare members, effective January 1, 2017, opioid prescription(s) for more than a 7-day supply and/or more than 30 MME/ day will require a prior authorization. Please note that MaineCare implemented a 30 MME limit January 1, 2013 that is still effective.

The following are general exceptions: pain associated with cancer treatment, end-of-life and hospice care, palliative care, and symptoms related to HIV/AIDS. Per MaineCare criteria, the diagnosis of cancer must be written on the prescription. A palliative care exception for any MaineCare opioid prescription will require prior authorization (PA) with appropriate clinical documentation.

Post-surgical members may receive prior authorizations for opiates up to a 60 days in length if medical necessity is provided by the surgical provider.

An MME conversion chart is available at www.mainearepd.org. Click on "General Pharmacy Info."